Centre Number			Candidate Number		
Surname					
Other Names					
Candidate Signature					



General Certificate of Education Advanced Subsidiary Examination June 2015

Mathematics

MS1B

Unit Statistics 1B

Wednesday 3 June 2015 9.00 am to 10.30 am

For this paper you must have:

• the blue AQA booklet of formulae and statistical tables. You may use a graphics calculator.

Time allowed

• 1 hour 30 minutes

Instructions

- Use black ink or black ball-point pen. Pencil should only be used for drawing.
- Fill in the boxes at the top of this page.
- Answer all questions.
- Write the question part reference (eg (a), (b)(i) etc) in the left-hand margin.
- You must answer each question in the space provided for that question. If you require extra space, use an AQA supplementary answer book; do not use the space provided for a different question.
- Do not write outside the box around each page.
- Show all necessary working; otherwise marks for method may be lost.
- Do all rough work in this book. Cross through any work that you do not want to be marked.
- The **final** answer to questions requiring the use of tables or calculators should normally be given to three significant figures.


Information

- The marks for questions are shown in brackets.
- The maximum mark for this paper is 75.
- Unit Statistics 1B has a written paper only.

Advice

- Unless stated otherwise, you may quote formulae, without proof, from the booklet.
- · You do not necessarily need to use all the space provided.





Answer all questions.

Answer each question in the space provided for that question.

The table shows the annual gas consumption, x kWh, and the annual electricity consumption, y kWh, for a sample of 10 bungalows of similar size and occupancy.

x	21 371	18 521	15 222	17312	19854	23 561	20 738	22 111	17 897	24 523
y	2281	2327	2221	2378	2787	2856	3078	2647	2566	2559

$$S_{xx} = 76581640$$
 $S_{yy} = 694250$ $S_{xy} = 3629670$

- (a) Calculate the value of r_{xy} , the product moment correlation coefficient between x and y. [2 marks]
- **(b)** Interpret your value of r_{xy} in the context of this question.

[2 marks]

QUESTION PART REFERENCE	Answer space for question 1



QUESTION PART REFERENCE	Answer space for question 1



2 The table summarises the diameters, d millimetres, of a random sample of 60 new cricket balls to be used in junior cricket.

Diameter (d mm)	Number of cricket balls
$65 < d \leqslant 66$	5
$66 < d \leqslant 67$	9
$67 < d \leqslant 68$	12
68 < d ≤ 69	15
$69 < d \leqslant 70$	10
$70 < d \leqslant 71$	7
$71 < d \leqslant 72$	2
Total	60

(a) Calculate estimates of the mean and the variance of these 60 diameters.

[4 marks]

(b) David, a retired professional cricketer, requests that the values calculated in part (a) are expressed in inches, rather than in millimetres.

Given that 1 inch is equivalent to approximately $25.4\,\mathrm{mm}$, calculate new values for the mean and the variance in response to David's request.

[2 marks]

PART	Answer space for question 2
REFERENCE	



QUESTION PART REFERENCE	Answer space for question 2



A ferry sails once each day from port D to port A. The ferry departs from D on time or late but never early. However, the ferry can arrive at A early, on time or late.

The **probabilities** for some combined events of departing from D and arriving at A are shown in the table below.

(a) Complete the table.

[2 marks]

- **(b)** Write down the probability that, on a particular day, the ferry:
 - (i) both departs and arrives on time;
 - (ii) departs late.

[2 marks]

- **(c)** Find the probability that, on a particular day, the ferry:
 - (i) arrives late, given that it departed late;
 - (ii) does **not** arrive late, given that it departed on time.

[5 marks]

(d) On three particular days, the ferry departs from port D on time.

Find the probability that, on these three days, the ferry arrives at port A early once, on time once and late once. Give your answer to three decimal places.

Arrive at A

[4 marks]

QUESTION PART REFERENCE (a)

Answer space for question 3

			Early	On time	Late	Total
	Depart	On time	0.16	0.56	0.08	
	from D	Late				
	<u> </u>	Total	0.22	0.65		1.00
						-
.						



QUESTION PART REFERENCE	Answer space for question 3



QUESTION PART REFERENCE	Answer space for question 3



QUESTION PART REFERENCE	Answer space for question 3



Stephan is a roofing contractor who is often required to replace loose ridge tiles on house roofs. In order to help him to quote more accurately the prices for such jobs in the future, he records, for each of 11 recently repaired roofs, the number of ridge tiles replaced, x_i , and the time taken, y_i hours. His results are shown in the table.

Roof (i)	1	2	3	4	5	6	7	8	9	10	11
x_i	8	11	14	14	16	20	22	22	25	27	30
y_i	5.0	5.2	6.3	7.2	8.0	8.8	10.6	11.0	11.8	12.1	13.0

(a) The pairs of data values for roofs 1 to 7 are plotted on the scatter diagram shown on the opposite page.

Plot the 4 pairs of data values for roofs 8 to 11 on the scatter diagram.

[2 marks]

(b) (i) Calculate the equation of the least squares regression line of y_i on x_i , and draw your line on the scatter diagram.

[6 marks]

- (ii) Interpret your values for the gradient and for the intercept of this regression line.

 [3 marks]
- (c) Estimate the time that it would take Stephan to replace 15 loose ridge tiles on a house roof.

[1 mark]

- (d) Given that r_i denotes the residual for the point representing roof i:
 - (i) calculate the value of r_6 ;


[2 marks]

(ii) state why the value of $\sum_{i=1}^{11} r_i$ gives no useful information about the connection between the number of ridge tiles replaced and the time taken.

[1 mark]

PART REFERENCE	Answer space for question 4







QUESTION PART REFERENCE	Answer space for question 4



QUESTION PART REFERENCE	Answer space for question 4



Wooden lawn edging is supplied in $1.8\,\mathrm{m}$ length rolls. The actual length, X metres, of a roll may be modelled by a normal distribution with mean 1.81 and standard deviation 0.08.

Determine the probability that a randomly selected roll has length:

- (i) less than 1.90 m;
- (ii) greater than 1.85 m;
- (iii) between 1.81 m and 1.85 m.

[6 marks]

(b) Plastic lawn edging is supplied in $9\,\mathrm{m}$ length rolls. The actual length, Y metres, of a roll may be modelled by a normal distribution with mean μ and standard deviation σ .

An analysis of a batch of rolls, selected at random, showed that

$$P(Y < 9.25) = 0.88$$

(i) Use this probability to find the value of z such that

$$9.25 - \mu = z \times \sigma$$

where z is a value of $Z \sim N(0, 1)$.

[2 marks]

(ii) Given also that

$$P(Y > 8.75) = 0.975$$

find values for μ and σ .

[4 marks]

REFERENCE	



QUESTION PART REFERENCE	Answer space for question 5



QUESTION PART REFERENCE	Answer space for question 5



QUESTION PART REFERENCE	Answer space for question 5



6 (a)	In a particular country, 35 per cent of the population is estimated to have at least one mobile phone.
		A sample of 40 people is selected from the population.
		Use the distribution $B(40,0.35)$ to estimate the probability that the number of people in the sample that have at least one mobile phone is:
	(i)	at most 15;
	(ii)	more than 10;
	(iii)	more than 12 but fewer than 18;
	(iv)	exactly equal to the mean of the distribution. [9 marks]
(b)	In the same country, 70 per cent of households have a landline telephone connection.
		A sample of 50 households is selected from all households in the country.
		Stating a necessary condition regarding this selection, estimate the probability that fewer than 30 households have a landline telephone connection.
		[4 marks]
QUESTION PART REFERENCE	Ans	wer space for question 6



QUESTION PART REFERENCE	Answer space for question 6



QUESTION PART REFERENCE	Answer space for question 6



QUESTION PART REFERENCE	Answer space for question 6



7 (a) The weight of a sack of mixed dog biscuits can be modelled by a normal distribution with a mean of $10.15 \, \text{kg}$ and a standard deviation of $0.3 \, \text{kg}$.

A pet shop purchases 12 such sacks that can be considered to be a random sample.

Calculate the probability that the mean weight of the 12 sacks is less than $10 \, \mathrm{kg}$.

[4 marks]

(b) The weight of dry cat food in a pouch can also be modelled by a normal distribution.

The contents, x grams, of each of a random sample of 40 pouches were weighed. Subsequent analysis of these weights gave

$$\bar{x} = 304.6$$
 and $s = 5.37$

(i) Construct a 99% confidence interval for the mean weight of dry cat food in a pouch. Give the limits to one decimal place.

[4 marks]

(ii) Comment, with justification, on each of the following two claims.

Claim 1: The mean weight of dry cat food in a pouch is more than 300 grams.

Claim 2: All pouches contain more than 300 grams of dry cat food.

[4 marks]

QUESTION PART REFERENCE	Answer space for question 7



QUESTION PART REFERENCE	Answer space for question 7



QUESTION PART REFERENCE	Answer space for question 7	
END OF QUESTIONS		
Copyright © 2015 AQA and its licensors. All rights reserved.		

