


5.

Figure 2



A parcel of weight 10 N lies on a rough plane inclined at an angle of 30° to the horizontal. A horizontal force of magnitude P newtons acts on the parcel, as shown in Figure 2. The parcel is in equilibrium and on the point of slipping up the plane. The normal reaction of the plane on the parcel is 18 N. The coefficient of friction between the parcel and the plane is μ . Find

(a) the value of P , (4)

(b) the value of μ . (5)

The horizontal force is removed.

(c) Determine whether or not the parcel moves. (5)



6. [In this question the horizontal unit vectors \mathbf{i} and \mathbf{j} are due east and due north respectively.]

A model boat A moves on a lake with constant velocity $(-\mathbf{i} + 6\mathbf{j}) \text{ m s}^{-1}$. At time $t = 0$, A is at the point with position vector $(2\mathbf{i} - 10\mathbf{j}) \text{ m}$. Find

(a) the speed of A , (2)

(b) the direction in which A is moving, giving your answer as a bearing. (3)

At time $t = 0$, a second boat B is at the point with position vector $(-26\mathbf{i} + 4\mathbf{j}) \text{ m}$.

Given that the velocity of B is $(3\mathbf{i} + 4\mathbf{j}) \text{ m s}^{-1}$,

(c) show that A and B will collide at a point P and find the position vector of P . (5)


Given instead that B has speed 8 m s^{-1} and moves in the direction of the vector $(3\mathbf{i} + 4\mathbf{j})$,

(d) find the distance of B from P when $t = 7 \text{ s}$. (6)



7.

Figure 3



A fixed wedge has two plane faces, each inclined at 30° to the horizontal. Two particles A and B , of mass $3m$ and m respectively, are attached to the ends of a light inextensible string. Each particle moves on one of the plane faces of the wedge. The string passes over a small smooth light pulley fixed at the top of the wedge. The face on which A moves is smooth. The face on which B moves is rough. The coefficient of friction between B and this face is μ . Particle A is held at rest with the string taut. The string lies in the same vertical plane as lines of greatest slope on each plane face of the wedge, as shown in Figure 3.

The particles are released from rest and start to move. Particle A moves downwards and B moves upwards. The accelerations of A and B each have magnitude $\frac{1}{10}g$.

- (a) By considering the motion of A , find, in terms of m and g , the tension in the string. (3)
- (b) By considering the motion of B , find the value of μ . (8)
- (c) Find the resultant force exerted by the string on the pulley, giving its magnitude and direction. (3)



