Write your name here Surname		Other names	
Pearson Edexcel Certificate Pearson Edexcel International GCSE	Centre Number	Candidate Number	
Chemistry Unit: KCH0/4CH0 Paper: 2C	/		
Tuesday 9 June 2015 – Afte Time: 1 hour	rnoon	Paper Reference KCH0/2C 4CH0/2C	J
You must have: Calculator, ruler		Total Mar	rks

Instructions

- Use **black** ink or ball-point pen.
- Fill in the boxes at the top of this page with your name, centre number and candidate number.
- Answer **all** questions.
- Answer the questions in the spaces provided there may be more space than you need.
- Show all the steps in any calculations and state the units.
- Some questions must be answered with a cross in a box ⊠. If you change your mind about an answer, put a line through the box ₩ and then mark your new answer with a cross ⊠.

Information

- The total mark for this paper is 60.
- The marks for each question are shown in brackets
 use this as a guide as to how much time to spend on each question.

Advice

- Read each question carefully before you start to answer it.
- Keep an eye on the time.
- Write your answers neatly and in good English.
- Try to answer every question.
- Check your answers if you have time at the end.



Turn over 🕨

PEARSON



P44269A ©2015 Pearson Education Ltd. 1/1/1/1/

		E		5	Ę	c	_ =			
	0	2 Helium 2 Helium			84 Krypton 36		B8 B6 B6 B6			
	7				80 Bromine 35	127 127 53 53	Astatine B5			
	ဖ		16 Oxygen 8	32 Sultur 16	79 Selenium 34	1	Polonium 84			
	Ŋ		14 Nitrogen 7	31 Phosphorus 15	75 AS Arsenic 33	122 Sb 51 51	83 Uth 83 Uth 83 Uth			
	4		12 Carbon 6	28 Silicon 14	73 Ge Germanium 32	119 50 50	20/ PD 20/ 82 dd 20			
	ო		5 Baran ⊐	27 Al Aluminium 13		115 Indium 49	Thattium 81			
			L		65 Zinc 30	112 Cadmium 48				
TABLE					63.5 Copper 29	Ag Silver 47				
NODIC					28 Nickel 28	106 Pd A6 46	Platinum 78			
THE PERIODIC TABLE					59 Cobalt 27		192 Iridium 77			
È					26 FG 56	Hathenium Brand	OS OSmium 76		1	nic Der
	Group	Hydrogen			55 Mn Manganese 25	96 99 MO Molybdenum Technetium 42 43	186 Rhenium 75		Key	Relative atomic mass Symbol Name Atomic number
					52 Chromium 24	96 MO 42 42	184 Tungsten 74		I	
					51 Vanadium 23	3	181 Tantatum 73			
						91 Zriconium 40	Hafnium 72 72			
					45 Scandium 21	33 33 ^{Yttrium} ← 89	139 La anthanur 57	227 Actinium 89		
	N		9 Beryllium 4	24 Mg 12	40 Calcium 20	ε	13/ Barium 56	Radium 88 88		
	-		- 7 Lithium 3	23 Sodium 11	39 Potassium 19		133 CS 55	223 Fr 87 87		
		1	N	က	4	ا ى	9	~		

P 4 4 2 6 9 A 0 2 2 0



BLANK PAGE

Answer ALL questions.

The table shows the numbers of protons, neutrons and electrons in some atoms and ions. 1 Atom or ion **Electrons Protons Neutrons** Ρ 6 8 6 Q 5 6 5 R 9 10 10 S 3 4 2 Т 6 6 6 (a) (i) Which particles have the same mass? (1) A electrons and protons **B** electrons and neutrons \times **C** neutrons and protons \times **D** electrons, neutrons and protons (ii) What is the atomic number of P? (1) **A** 6 \times \times В 8 $\boldsymbol{\times}$ **C** 12 **D** 14 (iii) What is the mass number of Q? (1) A 5 \mathbf{X} **B** 6 **C** 10 \times X **D** 11





2 The equation for the thermal decomposition of copper(II) carbonate is

 $CuCO_{_3}(s) \rightarrow CuO(s) + CO_{_2}(g)$

A student investigates the decomposition of copper(II) carbonate using this apparatus.



00.0

She uses this method.

- weigh the crucible, lid and copper(II) carbonate
- heat the crucible, lid and contents for 2 minutes
- allow to cool and then reweigh
- heat for a second period of 2 minutes
- allow to cool and then reweigh
- heat for a third period of 2 minutes
- allow to cool and then reweigh

The table shows the student's results.

	Mass of crucible, lid and contents in grams				
Experiment	before heating	after heating for 2 minutes	after heating for 4 minutes	after heating for 6 minutes	
1	26.3	23.0	21.9	21.4	
2	25.8	22.7	21.5	21.5	
3	26.0	23.0	21.2	21.2	
4	26.1	23.2	21.8	21.8	

(a) Why does the mass decrease during heating?

(1)



(b) State the colours of the solids in the reaction. (2)	
CuCO ₃ (s)	
CuO(s)	
(c) (i) In which experiment might the decomposition not be complete?(1)	
(ii) Give a reason for your choice. (1)	
(iii) Which statement could explain why the decomposition might not be complete? (1)	
 A The student used a higher temperature than in the other experiments. D The student used last some or (1) contained to be a start of the start of t	
 B The student used less copper(II) carbonate than in the other experiments. C The student heated the crucible without a lid on. 	
 D The student used a spirit burner instead of a Bunsen burner. 	
(d) In another experiment, the student calculates that she should obtain a mass of 3.7 g of CuO(s) after completely decomposing a sample of CuCO ₃ (s).	
She actually obtains a mass of 3.4 g of CuO(s).	
Calculate the percentage yield in her experiment. (2)	
percentage yield =	%
(Total for Question 2 = 8 marks)	

7



BLANK PAGE





(c) A student carries out some experiments to investigate displacement reactions.

She adds some halogen solutions to halide solutions and observes whether a reaction occurs.

The table shows her results.

	Halogen solution added				
Halide solution	bromine	chlorine	iodine		
lithium chloride	no reaction	(not done)	no reaction		
sodium bromide	(not done)	reaction occurs	no reaction		
potassium iodide	reaction occurs	reaction occurs	(not done)		

(i) The table shows that she did not do three experiments.

Suggest why she did not do these experiments.

(1)

(ii) The table shows that there was no reaction in three experiments.

Why was there no reaction in these experiments?

(1)



(iii) The student writes this word equation for one of the experiments in which a reaction occurs.	9
bromine + potassium iodide \rightarrow potassium bromine + iodine	
The name of one of the substances is incorrect.	
Write the correct name of this substance.	(1)
	(1)
(iv) A reaction occurs when the student adds chlorine solution to potassium iodide solution.	
Complete the chemical equation for this reaction.	(2)
Cl_2 + $Kl \rightarrow$ +	
(v) All displacement reactions are examples of redox reactions.	
State the meaning of the term redox .	
	(1)
(vi) The ionic equation for another reaction is	
$Br_2 + 2l^- \rightarrow 2Br^- + l_2$	
Explain which species is oxidised in this reaction.	(2)
	(2)
(Total for Question 3 = 10 r	marks)



11

$\begin{array}{c} \text{glucose} & \begin{array}{c} \text{reaction 1} \\ \text{ethanol} \\ \end{array} \\ \begin{array}{c} \text{freaction 3} \\ \end{array} \\ \begin{array}{c} \text{ethene} \\ \end{array} \\ \begin{array}{c} \text{ethene} \\ \end{array} \\ \begin{array}{c} \text{(a) (i) Two conditions used in reaction 1 are} \\ \end{array} \\ \begin{array}{c} \text{(a) a temperature of about 30 °C} \\ \text{(b) the use of water as a solvent for the glucose} \\ \end{array} \\ \begin{array}{c} \text{State the name of the catalyst used in this reaction.} \\ \begin{array}{c} \text{(1)} \\ \\ \text{(ii) Complete the equation for reaction 1.} \\ \\ \\ \text{(a) (b) Ethanol can also be manufactured by reaction 3, which uses steam, a catalyst of phosphoric acid and a pressure of about 65 atm. \\ \end{array} \\ \begin{array}{c} \text{State the temperature used in reaction 3.} \\ \end{array} \\ \begin{array}{c} \text{(1)} \\ \text{(c) State the type of reaction that occurs in} \\ \end{array} \\ \begin{array}{c} \text{(2)} \\ \text{reaction 1} \\ \end{array} \\ \end{array} \\ \end{array}$	4 The scheme shows some reactions involving ethanol.	
• a temperature of about 30 °C • the use of water as a solvent for the glucose State the name of the catalyst used in this reaction. (1) (ii) Complete the equation for reaction 1. $C_6H_{12}O_6 \rightarrow 2C_2H_5OH + \dots$ (b) Ethanol can also be manufactured by reaction 3, which uses steam, a catalyst of phosphoric acid and a pressure of about 65 atm. State the temperature used in reaction 3. (1) (c) State the type of reaction that occurs in (2) reaction 1	glucose reaction 1 ethanol ethene	
(ii) Complete the equation for reaction 1. $C_{6}H_{12}O_{6} \rightarrow 2C_{2}H_{5}OH + \dots \qquad (1)$ (b) Ethanol can also be manufactured by reaction 3, which uses steam, a catalyst of phosphoric acid and a pressure of about 65 atm. State the temperature used in reaction 3. (1) (c) State the type of reaction that occurs in (2) reaction 1	 a temperature of about 30 °C the use of water as a solvent for the glucose 	
(1) $C_{6}H_{12}O_{6} \rightarrow 2C_{2}H_{5}OH + \dots$ (b) Ethanol can also be manufactured by reaction 3, which uses steam, a catalyst of phosphoric acid and a pressure of about 65 atm. State the temperature used in reaction 3. (1) (c) State the type of reaction that occurs in (2) reaction 1		(1)
 (b) Ethanol can also be manufactured by reaction 3, which uses steam, a catalyst of phosphoric acid and a pressure of about 65 atm. State the temperature used in reaction 3. (1) (c) State the type of reaction that occurs in (2) 		(1)
(c) State the type of reaction that occurs in (2) reaction 1	(b) Ethanol can also be manufactured by reaction 3, which uses steam, a catalyst of	
(2) reaction 1	State the temperature used in reaction 3.	(1)
	(c) State the type of reaction that occurs in	(2)
reaction 3	reaction 1	
	reaction 3	

(d) State two advantages of using reaction 3 to manufacture ethanol rather than reaction 1.	(2)
	x #
(e) Give a reason why some countries use reaction 1 to manufacture ethanol.	
	(1)
(f) Reaction 2 may be used in the future to manufacture ethene.	
(i) Write an equation for this reaction.	(1)
(ii) What type of reaction is this?	
(ii) What type of reaction is this:	(1)
(Total for Question 4 = 10 r	narks)



5 A student uses this apparatus to measure the temperature change when lithium iodide dissolves in water.



He measures the steady temperature of the water before adding the lithium iodide.

He then adds the lithium iodide, stirs the mixture until all the solid dissolves and records the maximum temperature reached.

The diagram shows the thermometer readings before and after dissolving the lithium iodide.



(a) Use the readings to complete the table.

(3)

Temperature in °C after adding lithium iodide	
Temperature in °C before adding lithium iodide	
Temperature change in °C	



 (b) In a second experiment, using the same mass of water, the student records a temperature increase of 4.9 °C. (i) Use this expression to calculate the heat energy change in this experiment. heat energy change = mass of water × 4.2 × temperature change (in joules) (in grams) (in °C) 	(2)
heat energy change =	J
(ii) In this experiment, 6.3 g of lithium iodide were used.	
Calculate the amount, in moles, of lithium iodide in 6.3 g.	
[<i>M</i> _r of lithium iodide = 134]	(2)
amount of Lil =	mol



(c) In a third exp	eriment the studen	t obtains these result	-S.	
	heat energy chan		2400	
	amount of lithium		0.048	
(i) Calculate t	he molar enthalpy	change, in kJ/mol, in	this experiment.	(2)
		molar	enthalpy change =	kJ/mol
		nis experiment shows le solution is an exoth	s that dissolving lith	
Complete solution.	the energy level dia	agram to show the po	osition of the lithiur	n iodide
Label the o	diagram to show Δt	H, the molar enthalpy	/ change.	(2)
		Δ		
	Energy	lithium iodide +	water	
		(To	otal for Question 5	= 11 marks)

P 4 4 2 6 9 A 0 1 6 2 0



BLANK PAGE

6	Magnesium	Magnesium and its compounds have many uses.		
	Magnesium is never found as an element in the Earth's crust, but its compounds occur naturally in rocks and seawater.			
	(a) Suggest why magnesium is not found as an element in the Earth's crust.		(1)	
	(b) Magnesium can be extracted from seawater by a multi-stage process.			
	stage 1	calcium hydroxide reacts with magnesium chloride in seawater to fo a precipitate of magnesium hydroxide	rm	
	stage 2 the magnesium hydroxide is filtered off and converted ir chloride solution by reacting it with hydrochloric acid		um	
	stage 3	the magnesium chloride solution is converted into solid magnesium	chloride	
	stage 4	the solid magnesium chloride is melted and electrolysed		
	(i) Which stage involves a neutralisation reaction?		(1)	
	🖾 A st	tage 1		
	B s	tage 2		
	C s	tage 3		
	D s	tage 4		
	(ii) Sugg	est the name of the other product formed in stage 1.	(1)	
	(iii) What	t happens to the ions in magnesium chloride during melting?	(1)	

P 4 4 2 6 9 A 0 1 8 2 0



(d) Magnesium oxide can be used to make magnesium sulfate by this reaction.

 $MgO(s) \ + \ H_{_2}SO_{_4}(aq) \ \rightarrow \ MgSO_{_4}(aq) \ + \ H_{_2}O(I)$

A student is provided with a beaker of dilute sulfuric acid.

Outline the steps she should use to obtain a pure sample of hydrated magnesium sulfate crystals using this reaction.

(5)

(Total for Question 6 = 14 marks)

TOTAL FOR PAPER = 60 MARKS

Every effort has been made to contact copyright holders to obtain their permission for the use of copyright material. Pearson Education Ltd. will, if notified, be happy to rectify any errors or omissions and include any such rectifications in future editions.

