

**Paper 3B/4B: Further Statistics 1 Mark Schemes**

Question	Scheme	Marks	AOs	
<b>1</b>	$H_0 : \lambda = 5$ ( $\lambda = 2.5$ ) $H_1 : \lambda > 5$ ( $\lambda > 2.5$ )	B1	2.5	
	$X \sim \text{Po}(2.5)$	B1	3.3	
	<b>Method 1:</b>	<b>Method 2:</b>		
	$P(X \geq 7) = 1 - P(X \leq 6)$ $= 1 - 0.9858$	$P(X \geq 5) = 0.1088$ $P(X \geq 6) = 0.042$	M1	1.1b
	$= 0.0142$	CR $X \geq 6$	A1	1.1b
	$0.0142 < 0.05$ $7 \geq 6$ or 7 is in critical region or 7 is significant Reject $H_0$ . There is evidence at the 5% significance level that the level of pollution has increased. <b>or</b> There is evidence to support the scientists claim is justified		A1cso	2.2b
<b>(5 marks)</b>				
<b>Notes:</b>				
<b>B1:</b> Both hypotheses correct using $\lambda$ <b>or</b> $\mu$ <b>and</b> 5 <b>or</b> 2.5 <b>B1:</b> Realising that the model $\text{Po}(2.5)$ is to be used. This may be stated or used <b>M1:</b> Using or writing $1 - P(X \leq 6)$ <b>or</b> $1 - P(X < 7)$ a correct CR <b>or</b> $P(X \geq 5) = \text{awrt } 0.109$ <b>and</b> $P(X \geq 6) = \text{awrt } 0.042$ <b>A1:</b> awrt 0.0142 <b>or</b> CR $X \geq 6$ <b>or</b> $X > 5$ <b>M1:</b> A fully correct solution and drawing a correct inference in context				

Question	Scheme	Marks	AOs
<b>2(a)</b>	$P(X \geq 1) = 1 - P(X = 0)$ $1 - P(X = 0) = 0.049$	B1	3.1b
	$P(X = 0) = 0.951$	B1	1.1b
	$x^5 = 0.951$ $x = 0.99$	M1	3.1b
	$p = 0.01$	A1	1.1b
	$X \sim B(1000, 0.01)$	M1	3.3
	Mean = $np = 10$	A1ft	1.1b
	Variance = $np(1 - p) = 9.9$	A1ft	1.1b
		(7)	
<b>(b)</b>	$X \sim \text{Po}(\text{"10"})$ then require: $P(X > 6) = 1 - P(X \leq 6)$	M1	3.4
	$= 1 - 0.1301$		
	$= 0.870$	A1	1.1b
		(2)	
<b>(c)</b>	The approximation is valid as : the number of calls is large	B1	2.4
	The probability of connecting to the wrong agent is small	B1	2.4
		(2)	
<b>(d)</b>	The answer is accurate to 2 decimal place	<b>B1</b>	3.2b
		(1)	
<b>(12 marks)</b>			
<b>Notes:</b>			
<b>(a)</b>			
<b>B1:</b> Realising that the $P(\text{at least 1 call}) = 1 - P(X = 0)$			
<b>B1:</b> Calculating $P(X = 0) = 0.951$			
<b>M1:</b> Forming the equation $x^5 = \text{"their 0.951"}$ may be implied by $p = 0.01$			
<b>A1:</b> 0.01 only			
<b>M1:</b> Realising the need to use the model $B(1000, 0.01)$ This may be stated or used			
<b>A1:</b> Mean = 10 or ft their $p$ but only if $0 < p < 1$			
<b>A1:</b> Var = 9.9 or ft their $p$ but only if $0 < p < 1$			
<b>(b)</b>			
<b>M1:</b> Using the model $\text{Po}(\text{"their 10"})$ (this may be written or used) and $1 - P(X \leq 6)$			
<b>A1:</b> awrt 0.870 Award M1 A1 for awrt 0.870 with no incorrect working			
<b>(c)</b>			
<b>B1:</b> Explaining why approximation is valid - need the context of number and calls			
<b>B1:</b> Need the context connecting, wrong agent			
<b>(d)</b>			
<b>B1:</b> Evaluating the accuracy of their answer in (b). Allow 2 significant figures			

Question	Scheme	Marks	AOs	
<b>3(a)</b>	Expected value for 2 = $150 \times P(X = 2)$	M1	3.4	
	= 28.3015...	A1	1.1b	
	Expected value for 4 or more = $150 - (53.8 + 56.6 + 28.3 + 8.9)$ = 2.4	A1ft	1.1b	
	H <sub>0</sub> : Bin(20, 0.05) is a suitable model H <sub>1</sub> : Bin(20, 0.05) is not a suitable model	B1	2.5	
	Combining last two groups			
		$\geq 3$	M1	2.1
	<b>Observed frequency</b>	19		
	<b>Expected frequency</b>	11.3		
	$\nu = 4 - 1 = 3$		B1	1.1b
	Critical value, $\chi^2(0.05) = 7.815$		B1	1.1a
	Test statistic = $\frac{(43 - 53.8)^2}{53.8} + \frac{(62 - 56.6)^2}{56.6} + \dots$		M1	1.1b
	= 8.117		A1	1.1b
	In critical region, sufficient evidence to reject H <sub>0</sub> , accept H <sub>1</sub> Significant evidence at 5% level to reject the manager's model		A1	3.5a
		<b>(10)</b>		
<b>(b)</b>	$\nu = 4 - 2 = 2$			
	4 classes due to pooling	B1	2.4	
	2 restrictions (equal total and mean/proportion)	B1	2.4	
			<b>(2)</b>	
<b>(c)</b>	H <sub>0</sub> : Binomial distribution is a good model H <sub>1</sub> : Binomial distribution is not a good model	B1	3.4	
	Critical value, $\chi^2(0.05) = 5.991$ Test statistic is not in critical region, insufficient evidence to reject H <sub>0</sub> There is evidence that the Binomial distribution is a good model	B1	3.5a	
			<b>(2)</b>	
	<b>(14 marks)</b>			

**Question 3 notes:****(a)****M1:** Using the binomial model  $150 \times p^2 \times (1-p)^{18}$  may be implied by 28.3**A1:** awrt 28.3**A1:** awrt 2.4 or ft their “28.3”**B1:** Both hypotheses correct using the correct notation or written out in full**M1:** For recognising the need to combine groups**B1:** Number of degrees of freedom = 3 may be implied by a correct CV**B1:** awrt 7.82**M1:** Attempting to find  $\sum \frac{(O_i - E_i)^2}{E_i}$  or  $\sum \frac{O_i^2}{E_i} - N$  may be implied by awrt 8.12**A1:** awrt 8.12**A1:** Evaluating the outcome of a model by drawing a correct inference in context**(b)****B1:** Explaining why there are 4 classes**B1:** Explanation of why 2 is subtracted**(c)****B1:** Correct hypotheses for the refined model**B1:** The CV awrt 5.99 and drawing the correct inference for the refined model

Question	Scheme	Marks	AOs
4	Po(2.3) $n = 100$ $\mu = 2.3$ $\sigma^2 = 2.3$		
	CLT $\Rightarrow \bar{X} \approx N\left(2.3, \frac{2.3}{100}\right)$	M1 A1	3.1a 1.1b
	$P(\bar{X} > 2.5) = P\left(Z > \frac{2.5 - 2.3}{\sqrt{0.023}}\right)$	M1	3.4
	$= P(Z > 1.318..)$		
	$= 0.09632\dots$	A1	1.1b
		(4)	
<b>(4 marks)</b>			
<b>Notes:</b>			
<p><b>M1:</b> For realising the need to use the CLT to set <math>\bar{X} \approx</math> normal with correct mean May be implied by using the correct normal distribution</p> <p><b>A1:</b> For fully correct normal stated or used</p> <p><b>M1:</b> Use of the normal model to find <math>P(\bar{X} &gt; 2.5)</math>. Can be awarded for <math>\frac{2.5 - 2.3}{\sqrt{0.023}}</math> or awrt 1.32</p> <p><b>A1:</b> awrt 0.0963</p>			

Question	Scheme	Marks	AOs
<b>5(a)</b>	$\binom{7}{1} \times 0.15^2 \times (0.85)^6$	M1	3.3
	= 0.05940... = awrt <b>0.0594</b>	A1	1.1b
		<b>(2)</b>	
<b>(b)</b>	The model is only valid if:		
	the games (trials) are <b>independent</b>	B1	3.5b
	the probability of winning a prize, 0.15, is <b>constant</b> for each game	B1	3.5b
		<b>(2)</b>	
<b>(c)</b>	$18 = \frac{r}{p}$ and $6^2 = \frac{r(1-p)}{p^2}$	M1 A1	3.1b 1.1b
	Solving: $2p = 1 - p$	M1	1.1b
	$p = \frac{1}{3}$ ( $> 0.15$ ) so Mary has the greater chance of winning a prize	A1	3.2a
		<b>(4)</b>	
	<b>(8 marks)</b>		
	<b>Notes:</b>		
<b>5(a)</b>			
<b>M1:</b> For selecting an appropriate model negative binomial or B(7, 0.15) with an extra success in 8 <sup>th</sup> trial e.g.			
$\binom{7}{1} 0.15 \times (0.85)^6 \times 0.15$ Allow $\binom{7}{1} 0.85 \times (0.15)^6 \times 0.85$ may be implied by awrt 0.0594			
<b>A1:</b> awrt 0.0594			
<b>(b)</b>			
<b>B1:</b> Stating the first assumption that games are independent			
<b>B1:</b> Stating the second assumption that the probability remains constant			
<b>(c)</b>			
<b>M1:</b> Forming an equation for the mean or for the standard deviation			
<b>A1:</b> Both equations correct			
<b>M1:</b> Solving the 2 equations leading to $2p = 1 - p$			
<b>A1:</b> For $p = \frac{1}{3}$ followed by a correct deduction			

Question	Scheme	Marks	AOs
<b>6(a)</b>	$G_X(1) = 1$ gives	M1	2.1
	$k \times 6^2 = 1$ so $k = \frac{1}{36}$ *	A1*cs0	1.1b
		(2)	
<b>(b)</b>	$P(X=3) = \text{coefficient of } t^3$ so $G_X(t) = k(\dots + 4t^3 \dots)$	M1	1.1b
	[ $P(X=3) = ] \frac{1}{9}$	A1	1.1b
		(2)	
<b>(c)</b>	$G'_X(t) = 2k(3+t+2t^2) \times (1+4t)$	M1	2.1
	$E(X) = G'_X(1) = 2k(3+1+2) \times (1+4)$	M1	1.1b
	$= \frac{5}{3}$	A1	1.1b
	$G''_X(t) = 2k \left[ (3+t+2t^2) \times 4 + (1+4t)^2 \right]$	M1 A1	2.1 1.1b
	$G''_X(1) = 2k[6 \times 4 + 5^2] \quad \left\{ = \frac{49}{18} \right\}$	M1	1.1b
	$\text{Var}(X) = G''_X(1) + G'_X(1) - [G'_X(1)]^2 = \frac{49}{18} + \frac{5}{3} - \frac{25}{9}$	M1	2.1
	$= \frac{29}{18}$ *	A1*cs0	1.1b
		(8)	
<b>(d)</b>	$G_{2X+1}(t) = \frac{t}{36} (3+t^2+2(t^2)^2)^2$ [ $\times t$ or sub $t^2$ for $t$ ]	M1	3.1a
	$= G_{2X+1}(t) = \frac{t}{36} (3+t^2+2t^4)^2$	A1	1.1b
		(2)	
<b>(14 marks)</b>			
<b>Notes:</b>			
<b>(a)</b>			
<b>M1:</b> Stating $G_X(1) = 1$			
<b>A1*:</b> Fully correct proof with no errors cs0			
<b>(b)</b>			
<b>M1:</b> Attempting to find the coefficient of $t^3$ . May be implied by obtaining $\frac{1}{9}$ or awrt 0.11			
<b>A1:</b> $\frac{1}{9}$ , allow awrt 0.111			

**Question 6 notes continued:****(c)****M1:** Attempting to find  $G'_X(t)$ . Allow Chain rule or multiplying out the brackets and differentiating**M1:** Substituting  $t = 1$  into  $G'_X(t)$ **A1:**  $\frac{5}{3}$ , allow awrt 1.67**M1:** Attempting to find  $G''_X(t)$ **A1:**  $2k\left[(3+t+2t^2)\times 4+(1+4t)^2\right]$  or  $k(48t^2+24t+26)$  o.e.**A1:**  $2k[6\times 4+5^2]$  o.e.**M1:** Using  $G''_X(1)+G'_X(1)-[G'_X(1)]^2$  to find the Variance**A1\*:**  $\frac{29}{18}$  cso**(d)****M1:** Realising the need to  $\times t$  or sub  $t^2$  for  $t$ **A1:**  $\frac{t}{36}(3+t^2+2t^4)^2$ , or  $\frac{t}{36}(9+6t^2+13t^4+4t^6+4t^8)$  o.e.



Question	Scheme	Marks	AOs
<b>7(a)</b>	$X \sim B(20, 0.2)$ and seek $c$ such that $P(X \leq c) < 0.10$	M1	3.3
	$[P(X \leq 1) = 0.0692]$ CR is $X \leq 1$	A1	1.1b
		<b>(2)</b>	
<b>(b)</b>	Size = <b><u>0.0692</u></b>	B1ft	1.2
		<b>(1)</b>	
<b>(c)</b>	$Y =$ no. of spins until red obtained so $Y \sim \text{Geo}(0.2)$	M1	3.3
	$\mu = \frac{1}{p}$ so if $p < 0.2$ then mean is <u>larger</u> so seek $d$ so that $P(Y \geq d) < 0.10$	M1	2.4
	$P(Y \geq d) = (0.8)^{d-1}$	M1	3.4
	$(0.8)^{d-1} < 0.10 \Rightarrow d - 1 > \frac{\log(0.1)}{\log(0.8)}$	M1	1.1b
	$d > 11.3..$	A1	1.1b
	CR is $Y \geq 12$	A1	2.2b
		<b>(6)</b>	
<b>(d)</b>	Size = $[0.8^{11} = 0.085899\dots] = \mathbf{0.0859}$	B1	1.1b
		<b>(1)</b>	
<b>(e)(i)</b>	Power = $P(\text{reject } H_0 \text{ when it is false}) = P(X \leq 1 \mid X \sim B(20, p))$	M1	2.1
	$= (1-p)^{20} + 20(1-p)^{19} p$	M1	1.1b
	$= (1-p)^{19} (1+19p) *$	A1*cso	1.1b
<b>(ii)</b>	Power = $(1-p)^{11}$	B1	1.1b
		<b>(4)</b>	
<b>(f)</b>	Sam's test has smaller $P(\text{Type I error})$ (or size) so is better	B1	2.2a
	Power of Sam's test = 0.1755...	B1	1.1b
	Power of Tessa's test = $0.85^{11} = 0.1673\dots$	B1	1.1b
	So for $p = 0.15$ <b>Sam's test</b> is recommended	B1	2.2b
		<b>(4)</b>	
			<b>(18 marks)</b>

<b>Question 7 notes:</b>	
<b>(a)</b>	<p><b>M1:</b> Realising the need to use the model Using B(20,0.2) with method for finding the CR or implied by a correct CR</p> <p><b>A1:</b> <math>X \leq 1</math> or <math>X &lt; 2</math></p>
<b>(b)</b>	<p><b>B1:</b> awrt 0.0692</p>
<b>(c)</b>	<p><b>M1:</b> Realising that the model Geo(0.2) is needed. This may be written or used</p> <p><b>M1:</b> Realising the key step that they need to find <math>P(Y \geq d) &lt; 0.10</math></p> <p><b>M1:</b> Using the model <math>(0.8)^{d-1}</math></p> <p><b>M1:</b> Using the model <math>(0.8)^{d-1} &lt; 0.10</math> and finding a method to solve leading to a value/range of values for <math>d</math></p> <p><b>A1:</b> For <math>d &gt; 11.3..</math></p> <p><b>A1:</b> For <math>Y \geq 12</math> or <math>Y &gt; 11</math> (a correct inference)</p>
<b>(d)</b>	<p><b>B1ft:</b> awrt 0.0692. ft their answer to part (c)</p>
<b>(e)(i)</b>	<p><b>M1:</b> Using B(20, <math>p</math>) and realizing they need to find <math>P(X \leq 1)</math> o.e. This may be used or written</p> <p><b>M1:</b> Using <math>P(X=0) + P(X=1)</math></p> <p><b>A1*:</b> Fully correct proof ( no errors) cso</p>
<b>(ii)</b>	<p><b>B1:</b> For <math>(1-p)^{11}</math></p>
<b>(f)</b>	<p><b>B1:</b> Making a deduction about the tests using the answers to part(b) and (d)</p> <p><b>B1:</b> awrt 0.0176</p> <p><b>B1:</b> awrt 0.167</p> <p><b>B1:</b> A correct inference about which test is recommended</p>