







3. The curve  $C$  has equation

$$x = 2 \sin y.$$

(a) Show that the point  $P\left(\sqrt{2}, \frac{\pi}{4}\right)$  lies on  $C$ . (1)

(b) Show that  $\frac{dy}{dx} = \frac{1}{\sqrt{2}}$  at  $P$ . (4)

(c) Find an equation of the normal to  $C$  at  $P$ . Give your answer in the form  $y = mx + c$ , where  $m$  and  $c$  are exact constants. (4)

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---







5.

Figure 1




Figure 1 shows an oscilloscope screen.

The curve shown on the screen satisfies the equation

$$y = \sqrt{3} \cos x + \sin x.$$

(a) Express the equation of the curve in the form  $y = R \sin(x + \alpha)$ , where  $R$  and  $\alpha$  are constants,  $R > 0$  and  $0 < \alpha < \frac{\pi}{2}$ . (4)

(b) Find the values of  $x$ ,  $0 \leq x < 2\pi$ , for which  $y = 1$ . (4)

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---



**Question 5 continued**

Lined writing area for the question response.

(Total 8 marks)

**Q5**





6. The function  $f$  is defined by

$$f : x \mapsto \ln(4 - 2x), \quad x < 2 \quad \text{and} \quad x \in \mathbb{R}.$$

(a) Show that the inverse function of  $f$  is defined by

$$f^{-1} : x \mapsto 2 - \frac{1}{2}e^x$$

and write down the domain of  $f^{-1}$ .

**(4)**

(b) Write down the range of  $f^{-1}$ .

**(1)**

(c) In the space provided on page 16, sketch the graph of  $y = f^{-1}(x)$ . State the coordinates of the points of intersection with the  $x$  and  $y$  axes.

**(4)**

The graph of  $y = x + 2$  crosses the graph of  $y = f^{-1}(x)$  at  $x = k$ .

The iterative formula

$$x_{n+1} = -\frac{1}{2}e^{x_n}, \quad x_0 = -0.3$$

is used to find an approximate value for  $k$ .

(d) Calculate the values of  $x_1$  and  $x_2$ , giving your answers to 4 decimal places.

**(2)**

(e) Find the value of  $k$  to 3 decimal places.

**(2)**

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---



**Question 6 continued**





**Question 7 continued**

**Q7**

**(Total 13 marks)**



8. (i) Prove that

$$\sec^2 x - \operatorname{cosec}^2 x \equiv \tan^2 x - \cot^2 x. \quad (3)$$

(ii) Given that

$$y = \arccos x, \quad -1 \leq x \leq 1 \text{ and } 0 \leq y \leq \pi,$$

(a) express  $\arcsin x$  in terms of  $y$ . (2)

(b) Hence evaluate  $\arccos x + \arcsin x$ . Give your answer in terms of  $\pi$ . (1)

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---



**Question 8 continued**

*[The answer area contains horizontal lines for writing but is currently blank.]*

**(Total 6 marks)**

**Q8**

**TOTAL FOR PAPER: 75 MARKS**

**END**

