

Please check the examination details below before entering your candidate information

Candidate surname

Other names

Centre Number

Candidate Number

--	--	--	--	--

--	--	--	--	--

**Pearson Edexcel Level 3 GCE**

**Tuesday 20 June 2023**

Afternoon

Paper  
reference

**9MA0/31**

**Mathematics**

**Advanced**

**PAPER 31: Statistics**

**You must have:**

Mathematical Formulae and Statistical Tables (Green), calculator

Total Marks

**Candidates may use any calculator allowed by Pearson regulations. Calculators must not have the facility for symbolic algebra manipulation, differentiation and integration, or have retrievable mathematical formulae stored in them.**

### Instructions

- Use **black** ink or ball-point pen.
- If pencil is used for diagrams/sketches/graphs it must be dark (HB or B).
- **Fill in the boxes** at the top of this page with your name, centre number and candidate number.
- Answer **all** questions and ensure that your answers to parts of questions are clearly labelled.
- Answer the questions in the spaces provided  
– *there may be more space than you need.*
- You should show sufficient working to make your methods clear. Answers without working may not gain full credit.
- Values from statistical tables should be quoted in full. If a calculator is used instead of tables the value should be given to an equivalent degree of accuracy.
- Inexact answers should be given to three significant figures unless otherwise stated.

### Information

- A booklet 'Mathematical Formulae and Statistical Tables' is provided.
- The total mark for this part of the examination is 50. There are 6 questions.
- The marks for **each** question are shown in brackets  
– *use this as a guide as to how much time to spend on each question.*

### Advice

- Read each question carefully before you start to answer it.
- Try to answer every question.
- Check your answers if you have time at the end.

Turn over ►


P72819A

©2023 Pearson Education Ltd.  
N:1/1/1/



Pearson

1. The Venn diagram, where  $p$  and  $q$  are probabilities, shows the three events  $A$ ,  $B$  and  $C$  and their associated probabilities.



- (a) Find  $P(A)$

(1)

The events  $B$  and  $C$  are independent.

- (b) Find the value of  $p$  and the value of  $q$

(3)

- (c) Find  $P(A|B')$

(2)

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---





2. A machine fills packets with sweets and  $\frac{1}{7}$  of the packets also contain a prize.

The packets of sweets are placed in boxes before being delivered to shops.  
There are 40 packets of sweets in each box.

The random variable  $T$  represents the number of packets of sweets that contain a prize in each box.

- (a) State a condition needed for  $T$  to be modelled by  $B(40, \frac{1}{7})$  (1)

A box is selected at random.

- (b) Using  $T \sim B(40, \frac{1}{7})$  find (2)
- (i) the probability that the box has exactly 6 packets containing a prize,
  - (ii) the probability that the box has fewer than 3 packets containing a prize.

Kamil's sweet shop buys 5 boxes of these sweets.

- (c) Find the probability that exactly 2 of these 5 boxes have fewer than 3 packets containing a prize. (2)

Kamil claims that the proportion of packets containing a prize is less than  $\frac{1}{7}$

A random sample of 110 packets is taken and 9 packets contain a prize.

- (d) Use a suitable test to assess Kamil's claim. (4)
- You should
- state your hypotheses clearly
  - use a 5% level of significance









3. Ben is studying the Daily Total Rainfall,  $x$  mm, in Leeming for 1987

He used all the data from the large data set and summarised the information in the following table.

$x$	0	0.1–0.5	0.6–1.0	1.1–1.9	2.0–4.0	4.1–6.9	7.0–12.0	12.1–20.9	21.0–32.0	tr
Frequency	55	18	18	21	17	9	9	6	2	29

(a) Explain how the data will need to be cleaned before Ben can start to calculate statistics such as the mean and standard deviation. (2)

Using all 184 of these values, Ben estimates  $\sum x = 390$  and  $\sum x^2 = 4336$

(b) Calculate estimates for

- (i) the mean Daily Total Rainfall,
- (ii) the standard deviation of the Daily Total Rainfall. (3)

Ben suggests using the statistic calculated in part (b)(i) to estimate the annual mean Daily Total Rainfall in Leeming for 1987

(c) Using your knowledge of the large data set,

- (i) give a reason why these data would not be suitable,
- (ii) state, giving a reason, how you would expect the estimate in part (b)(i) to differ from the actual annual mean Daily Total Rainfall in Leeming for 1987 (2)

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---







4. A study was made of adult men from region  $A$  of a country. It was found that their heights were normally distributed with a mean of 175.4 cm and standard deviation 6.8 cm.

(a) Find the proportion of these men that are taller than 180 cm.

(1)

A student claimed that the mean height of adult men from region  $B$  of this country was different from the mean height of adult men from region  $A$ .

A random sample of 52 adult men from region  $B$  had a mean height of 177.2 cm

The student assumed that the standard deviation of heights of adult men was 6.8 cm both for region  $A$  and region  $B$ .

(b) Use a suitable test to assess the student's claim.

You should

- state your hypotheses clearly
- use a 5% level of significance

(4)

(c) Find the  $p$ -value for the test in part (b)

(1)

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



Question 4 continued

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

Ruled area for writing answers, containing 28 horizontal lines.

(Total for Question 4 is 6 marks)



5. Tisam is playing a game.  
She uses a ball, a cup and a spinner.

The random variable  $X$  represents the number the spinner lands on when it is spun.  
The probability distribution of  $X$  is given in the following table

$x$	20	50	80	100
$P(X = x)$	$a$	$b$	$c$	$d$

where  $a$ ,  $b$ ,  $c$  and  $d$  are probabilities.

To play the game

- the spinner is spun to obtain a value of  $x$
- Tisam then stands  $x$  cm from the cup and tries to throw the ball into the cup

The event  $S$  represents the event that Tisam successfully throws the ball into the cup.

To model this game Tisam assumes that

- $P(S | \{X = x\}) = \frac{k}{x}$  where  $k$  is a constant
- $P(S \cap \{X = x\})$  should be the same whatever value of  $x$  is obtained from the spinner

Using Tisam's model,

(a) show that  $c = \frac{8}{5}b$  (2)

(b) find the probability distribution of  $X$  (5)

Nav tries, a large number of times, to throw the ball into the cup from a distance of 100 cm.

He successfully gets the ball in the cup 30% of the time.

- (c) State, giving a reason, why Tisam's model of this game is not suitable to describe Nav playing the game for all values of  $X$  (1)

---

---

---

---

---

---

---

---

---

---



**Question 5 continued**

Lined writing area for the answer to Question 5.

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



P 7 2 8 1 9 A 0 1 3 2 0



**Question 5 continued**

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

**(Total for Question 5 is 8 marks)**

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



6. A medical researcher is studying the number of hours,  $T$ , a patient stays in hospital following a particular operation.

The histogram on the page opposite summarises the results for a random sample of 90 patients.

- (a) Use the histogram to estimate  $P(10 < T < 30)$  (2)

For these 90 patients the time spent in hospital following the operation had

- a mean of 14.9 hours
- a standard deviation of 9.3 hours

Tomas suggests that  $T$  can be modelled by  $N(14.9, 9.3^2)$

- (b) With reference to the histogram, state, giving a reason, whether or not Tomas' model could be suitable. (1)

Xiang suggests that the frequency polygon based on this histogram could be modelled by a curve with equation

$$y = kxe^{-x} \quad 0 \leq x \leq 4$$

where

- $x$  is measured in **tens of hours**
- $k$  is a constant

- (c) Use algebraic integration to show that

$$\int_0^n xe^{-x} dx = 1 - (n + 1)e^{-n} \quad (4)$$

- (d) Show that, for Xiang's model,  $k = 99$  to the nearest integer. (3)

- (e) Estimate  $P(10 < T < 30)$  using  
 (i) Tomas' model of  $T \sim N(14.9, 9.3^2)$  (1)

- (ii) Xiang's curve with equation  $y = 99xe^{-x}$  and the answer to part (c) (2)


The researcher decides to use Xiang's curve to model  $P(a < T < b)$

- (f) State one limitation of Xiang's model. (1)





Question 6 continued




---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



P 7 2 8 1 9 A 0 1 7 2 0



Question 6 continued

Lined writing area for the answer to Question 6 continued.

DO NOT WRITE IN THIS AREA





Please check the examination details below before entering your candidate information

Candidate surname

Other names

Centre Number

Candidate Number

--	--	--	--	--

--	--	--	--	--

**Pearson Edexcel Level 3 GCE**

**Tuesday 20 June 2023**

Afternoon

Paper  
reference

**9MA0/32**

**Mathematics**

**Advanced**

**PAPER 32: Mechanics**

**You must have:**

Mathematical Formulae and Statistical Tables (Green), calculator

Total Marks

**Candidates may use any calculator allowed by Pearson regulations. Calculators must not have the facility for symbolic algebra manipulation, differentiation and integration, or have retrievable mathematical formulae stored in them.**

### Instructions

- Use **black** ink or ball-point pen.
- If pencil is used for diagrams/sketches/graphs it must be dark (HB or B).
- **Fill in the boxes** at the top of this page with your name, centre number and candidate number.
- Answer **all** questions and ensure that your answers to parts of questions are clearly labelled.
- Answer the questions in the spaces provided  
– *there may be more space than you need.*
- You should show sufficient working to make your methods clear. Answers without working may not gain full credit.
- Unless otherwise indicated, whenever a value of  $g$  is required, take  $g = 9.8 \text{ m s}^{-2}$  and give your answer to either 2 significant figures or 3 significant figures.

### Information

- A booklet 'Mathematical Formulae and Statistical Tables' is provided.
- The total mark for this part of the examination is 50. There are 6 questions.
- The marks for **each** question are shown in brackets  
– *use this as a guide as to how much time to spend on each question.*

### Advice

- Read each question carefully before you start to answer it.
- Try to answer every question.
- Check your answers if you have time at the end.

Turn over ►

P72820A

©2023 Pearson Education Ltd.  
N:1/1/1/1/1/1/



  
Pearson





2.



Figure 1

A particle  $P$  has mass 5 kg.

The particle is pulled along a rough horizontal plane by a horizontal force of magnitude 28 N.

The only resistance to motion is a frictional force of magnitude  $F$  newtons, as shown in Figure 1.

- (a) Find the magnitude of the normal reaction of the plane on  $P$  (1)

The particle is accelerating along the plane at  $1.4 \text{ m s}^{-2}$

- (b) Find the value of  $F$  (2)

The coefficient of friction between  $P$  and the plane is  $\mu$

- (c) Find the value of  $\mu$ , giving your answer to 2 significant figures. (1)

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA







3. At time  $t$  seconds, where  $t \geq 0$ , a particle  $P$  has velocity  $\mathbf{v}$   $\text{ms}^{-1}$  where

$$\mathbf{v} = (t^2 - 3t + 7)\mathbf{i} + (2t^2 - 3)\mathbf{j}$$

Find

- (a) the speed of  $P$  at time  $t = 0$  (3)
- (b) the value of  $t$  when  $P$  is moving parallel to  $(\mathbf{i} + \mathbf{j})$  (2)
- (c) the acceleration of  $P$  at time  $t$  seconds (2)
- (d) the value of  $t$  when the direction of the acceleration of  $P$  is perpendicular to  $\mathbf{i}$  (2)

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA





4. [In this question,  $\mathbf{i}$  and  $\mathbf{j}$  are horizontal unit vectors and position vectors are given relative to a fixed origin  $O$ ]

A particle  $P$  is moving on a smooth horizontal plane.

The particle has constant acceleration  $(2.4\mathbf{i} + \mathbf{j})\text{ms}^{-2}$

At time  $t = 0$ ,  $P$  passes through the point  $A$ .

At time  $t = 5$  s,  $P$  passes through the point  $B$ .

The velocity of  $P$  as it passes through  $A$  is  $(-16\mathbf{i} - 3\mathbf{j})\text{ms}^{-1}$

- (a) Find the speed of  $P$  as it passes through  $B$ . (4)

The position vector of  $A$  is  $(44\mathbf{i} - 10\mathbf{j})\text{m}$ .

At time  $t = T$  seconds, where  $T > 5$ ,  $P$  passes through the point  $C$ .

The position vector of  $C$  is  $(4\mathbf{i} + c\mathbf{j})\text{m}$ .

- (b) Find the value of  $T$ . (3)

- (c) Find the value of  $c$ . (3)



Question 4 continued

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

Lined writing area for the response to Question 4.



P 7 2 8 2 0 A 0 9 2 0







5.




Figure 2

A small ball is projected with speed  $28 \text{ m s}^{-1}$  from a point  $O$  on horizontal ground.

After moving for  $T$  seconds, the ball passes through the point  $A$ .

The point  $A$  is  $40 \text{ m}$  horizontally and  $20 \text{ m}$  vertically from the point  $O$ , as shown in Figure 2.

The motion of the ball from  $O$  to  $A$  is modelled as that of a particle moving freely under gravity.

Given that the ball is projected at an angle  $\alpha$  to the ground, use the model to

(a) show that  $T = \frac{10}{7 \cos \alpha}$  (2)

(b) show that  $\tan^2 \alpha - 4 \tan \alpha + 3 = 0$  (5)

(c) find the greatest possible height, in metres, of the ball above the ground as the ball moves from  $O$  to  $A$ . (3)

The model does not include air resistance.

(d) State one other limitation of the model. (1)





**Question 5 continued**

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

Lined writing area for the answer.



P 7 2 8 2 0 A 0 1 3 2 0

**Question 5 continued**

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA





6.




Figure 3

A rod  $AB$  has mass  $M$  and length  $2a$ .

The rod has its end  $A$  on rough horizontal ground and its end  $B$  against a smooth vertical wall.

The rod makes an angle  $\theta$  with the ground, as shown in Figure 3.

The rod is at rest in limiting equilibrium.

- (a) State the direction (left or right on Figure 3 above) of the frictional force acting on the rod at  $A$ . **Give a reason for your answer.**

(1)

The magnitude of the normal reaction of the wall on the rod at  $B$  is  $S$ .

In an initial model, the rod is modelled as being **uniform**.

**Use this initial model to answer parts (b), (c) and (d).**

- (b) By taking moments about  $A$ , show that

$$S = \frac{1}{2} Mg \cot \theta \quad (3)$$

The coefficient of friction between the rod and the ground is  $\mu$

Given that  $\tan \theta = \frac{3}{4}$

- (c) find the value of  $\mu$  (5)
- (d) find, in terms of  $M$  and  $g$ , the magnitude of the resultant force acting on the rod at  $A$ . (3)

In a new model, the rod is modelled as being **non-uniform**, with its centre of mass closer to  $B$  than it is to  $A$ .

A new value for  $S$  is calculated using this new model, with  $\tan \theta = \frac{3}{4}$

- (e) State whether this new value for  $S$  is larger, smaller or equal to the value that  $S$  would take using the initial model. **Give a reason for your answer.** (1)



**Question 6 continued**

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

Lined area for writing the answer to Question 6.



**Question 6 continued**

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



**Question 6 continued**

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

Lined area for writing the answer to Question 6.



