

Physics B (Advancing Physics)

Advanced GCE

Unit **G495**: Field and Particle Pictures

Mark Scheme for June 2012

OCR (Oxford Cambridge and RSA) is a leading UK awarding body, providing a wide range of qualifications to meet the needs of candidates of all ages and abilities. OCR qualifications include AS/A Levels, Diplomas, GCSEs, OCR Nationals, Functional Skills, Key Skills, Entry Level qualifications, NVQs and vocational qualifications in areas such as IT, business, languages, teaching/training, administration and secretarial skills.

It is also responsible for developing new specifications to meet national requirements and the needs of students and teachers. OCR is a not-for-profit organisation; any surplus made is invested back into the establishment to help towards the development of qualifications and support, which keep pace with the changing needs of today's society.

This mark scheme is published as an aid to teachers and students, to indicate the requirements of the examination. It shows the basis on which marks were awarded by examiners. It does not indicate the details of the discussions which took place at an examiners' meeting before marking commenced.

All examiners are instructed that alternative correct answers and unexpected approaches in candidates' scripts must be given marks that fairly reflect the relevant knowledge and skills demonstrated.

Mark schemes should be read in conjunction with the published question papers and the report on the examination.

OCR will not enter into any discussion or correspondence in connection with this mark scheme.















© OCR 2012

Any enquiries about publications should be addressed to:

OCR Publications
PO Box 5050
Annesley
NOTTINGHAM
NG15 0DL

Telephone: 0870 770 6622
Facsimile: 01223 552610
E-mail: publications@ocr.org.uk

Annotations available in Scoris

Annotation	Meaning
	Benefit of doubt given
	Contradiction
	Incorrect response
	Error carried forward
	Follow through
	Not answered question
	Benefit of doubt not given
	Power of 10 error
	Omission mark
	Rounding error
	Error in number of significant figures
	Correct response
	Arithmetic error
	Wrong physics or equation

Annotations in Mark Scheme

Annotation	Meaning
/	alternative and acceptable answers for the same marking point
(1)	Separates marking points
reject	Answers which are not worthy of credit
not	Answers which are not worthy of credit
IGNORE	Statements which are irrelevant
ALLOW	Answers that can be accepted
()	Words which are not essential to gain credit
—	Underlined words must be present in answer to score a mark
ecf	Error carried forward
AW	Alternative wording
ORA	Or reverse argument

Subject specific Marking Instructions

The following questions should be annotated with ticks to show where marks have been awarded in the body of the text:

Question		Answer	Marks	Guidance
1	(a)	C	1	
	(b)	$I = 200 \times 1.8 / 400$ $= 0.90 \text{ A (1)}$	1	No ecf from 1(a)
2	(a)	neutron	1	
	(b)	neutron	1	
3		Units of $k = \text{N C}^{-1} \text{m}^2 \text{C}^{-1}$ (1) (which = $\text{N m}^2 \text{C}^{-2}$)	1	Allow unit derived from algebraic rearrangement to $k = Fr^2/Q_1Q_2$ or Fr^2/Q^2
4	(a)	A	1	
	(b)	mass lost $\text{s}^{-1} = \text{power}/c^2 = 4 \times 10^{26} / 9.0 \times 10^{16}$ (1) $= 4 \times 10^9 \text{ kg (1)}$	2	This is the only question for which a sig fig penalty applies. No more than two sig fig for second mark. Bald correct answer to one or two sig fig scores 2. (eg 4×10^9 or 4.4×10^9) Bald correct answer to more than two sig fig scores 1. 4.4 recurring scores 1
5	(a)	$V = 9.0 \times 10^9 \times 1.6 \times 10^{-19} / 5.0 \times 10^{-6}$ (1) $= 2.9 \times 10^{-4} \text{ J C}^{-1}$ (1)	2	Allow $V (= Er) = 58 \times 5 \times 10^{-6}$ (1) $= 2.9 \times 10^{-4} \text{ J C}^{-1}$ (1) Penalise 1 mark for use of $V = Ed$ (uniform field equation) Allow $3 \times 10^{-4} \text{ J C}^{-1}$, $2.88 \times 10^{-4} \text{ J C}^{-1}$ Allow bald answer
	(b)	Zero (1)	1	Ignore unit
	(c)	Twice the value (1)	1	Accept 2 x value from 5(a) and allow ecf. Ignore unit and any spurious justifications.

Question		Answer	Marks	Guidance
6		$r = 2.5 \times 10^{-20} / 0.7 \times 1.6 \times 10^{-19} \text{ (1)}$ $= 0.22 \text{ (m)} \text{ (1)}$	2	Award 1 mark for mv/Bq Need to see own value Award 1 mark only for bald correct answer to two or more sig figs Allow reverse arguments leading to correct comparison of momentum ($2.24 \times 10^{-20} \text{ kg m s}^{-1}$) and field strength (0.781 T) with given values.
7		$26 \text{ yrs} = 0.864 \text{ half lives (1)}$ $(1/2)^{0.864} \text{ (1)}$ $= 0.549 = 55 \% \text{ (1)}$ or $\lambda = 0.693/30.1 \text{ (1)} = 0.023 \text{ (yr}^{-1}\text{)}$ $N/N_0 = e^{-(0.693/30.1)26} \text{ (1)}$ $= 0.549 = 55\% \text{ (1)}$	3	Allow use of 30 yr for half life (gives 54.8 %). Beware: assumption of linear decay gives 56.8% – scores one mark if correct λ or correct number of half lives. Credit λ in years or seconds ($= 7.3 \times 10^{-10}$) Accept activity equation and arbitrary values of N_0 or A_0 in calculation. Allow use of activity A rather than number, N.
8		$F = 0.23 \times 0.15 \times 0.4 \text{ (1)}$ $= 0.014 \text{ N (1)}$	2	One mark only if any power of ten error. Accept 0.0138 N and 0.01 N.
Section A Total			19	

Question		Answer	Marks	Guidance
9	(a)	<p>Any three from:</p> <ul style="list-style-type: none"> • electrons attracted to nucleus (along line joining centre of particles) • Force or acceleration increases as distance from nucleus decreases AW • So curvature of track increases as distance from nucleus decreases AW 	3	<p>Use ticks to indicate marks awarded when marking this question.</p> <p>In cases of contradiction between the written answer and a diagram the written answer overrides the diagram.</p> <p>Accept F varies with $1/r^2$ for second mark</p> <p>QWC mark only available if path and force variation are clearly linked. Do not award if the answer includes inappropriate physics such as incorrect application of centripetal force.</p>
	(b)	(i) <p>Units of E identified as $\text{kg m}^2 \text{s}^{-2}$ or $\text{kg}(\text{ms}^{-1})^2$ (1) Units of E/c expressed as $\text{kg m}^2 \text{s}^{-2}/\text{m s}^{-1}$ or $\text{kg}(\text{ms}^{-1})^2/\text{m s}^{-1}$ (1)</p> <p>OR</p> <p>E/c has units $\text{N m}/\text{m s}^{-1} = \text{N s}$ (1) p units of N s from $f = \text{rate of change of momentum}$ (1)</p>	2	<p>Allow: $p = E/c = mc^2/c = mc$ (1) mc has units kgms^{-1} (1)</p> <p>Argument must be clearly made. (Going from Nm to answer is not sufficient)</p>
		(ii) <p>$\lambda = 3.0 \times 10^8 \times 6.6 \times 10^{-34} / 6.8 \times 10^{-11}$ (1) $= 2.91 \times 10^{-15}$ (1)</p>	2	<p>(For information momentum is $2.26 \times 10^{-19} \text{kgms}^{-1}$) Need own value of final answer Allow reverse argument leading to energy $6.6 \times 10^{-11} \text{J}$ Any evidence of direct use of $E = hf$ scores zero</p>
	(c)	<p>θ found as $49 - 50^\circ$ (1) $b = 1.2 \times 2.91 \times 10^{-15} / \sin 50^\circ$ (1) $= 4.6 \times 10^{-15} \text{m}$ (1)</p>	3	<p>Credit answers in range $4.5 - 4.8 \times 10^{-15} \text{m}$, but also allow ecf on λ values which round to $3 \times 10^{-15} \text{m}$. Allow ecf within part question for incorrect angle for 2 max</p>
Total			10	

Question			Answer	Marks	Guidance
10	(a)	(i)	mc^2 (at rest) OR m_0c^2	1	Accept "the energy equivalence of rest mass"
		(ii)	$(940 + 220)/940 = 1.23$	1	Need own value or clear working
		(iii)	$1.23^2 = 1/(1 - v^2/c^2)$ (1) (= 1.51) $v^2/c^2 = 0.339$ $v/c = 0.58$ (1)	2	Have to give own value of v (or of v/c, or both) Allow using gamma = 1.2 giving v/c = 0.55 Allow reverse argument: $\text{gamma}^2 = 1/(1 - 0.6^2)$ (1) gamma = 1.25 (1)
	(b)		Proton energy = $220 \times 10^6 \times 1.6 \times 10^{-19} = 3.52 \times 10^{-11}$ (1) Dose equivalent per proton = 1.17×10^{-6} Sv (1) Number of protons to deliver dose equivalent = 1.1×10^5 (1) OR energy = $(125 \times 10^{-3} \times 3.0 \times 10^{-4})/10 = 3.75 \times 10^{-6}$ J (1) No. of protons = $3.75 \times 10^{-6} / (220 \times 10^6 \times 1.6 \times 10^{-19})$ (1) = 1.1×10^5 (1)	3	Credit all other valid methods Accept any answer which will round to 1.1×10^5 Allow reverse argument starting from 10^5 protons: E.G. Energy = $10^5 \times 220 \times 10^6 \times 1.6 \times 10^{-19} = 3.52 \times 10^{-6}$ (1) Effective dose equivalent = $3.52 \times 10^{-6} \times 10/3.0 \times 10^{-4}$ (1) = 117 mSv (1)
	(c)	(i)	Answer = 13-14 cm (1) This is (the depth at which) the steepest gradient occurs (1) (At this depth) the energy loss per cm is greatest (1)	3	Accept greatest rate
		(ii)	Calculation in (b) assumes all energy is absorbed by the cells (1) Graph shows energy is transferred over a range of depths (1)	2	Accept alternative wording
Total				12	

Question			Answer	Marks	Guidance
11	(a)	(i)	Five lines, equally spaced by eye (1) Arrows pointing to the left. (1)	2	lines must be straight, touching the plates, perpendicular to the plates and approximately symmetrical about the centre of the plates. Allow edge effects illustrated.
	(a)	(ii)	vertical line in middle of gap (by eye)	1	label not required
	(b)		$E = 48 \times 10^3 / 0.28 = 0.17$ (1) (MV m^{-1})	1	Need own value or clear working. POT error scores zero.
	(c)	(i)	Drag = weight and calculation showing weight is $1.3 \times 10^{-6} \times 9.8 = 1.3 \times 10^{-5}$ N (1)	1	Both parts needed; don't accept force for drag. Need own value or clear working.
		(ii)	$F = 2.2 \times 10^{-13} \times 0.17 \times 10^6 = 3.7 \times 10^{-8}$ N (1)	1	If 0.2 MV m^{-1} is used accept 4.4×10^{-8} N If 0.171 MV m^{-1} is used accept 3.8×10^{-8} N Need own value or clear working. Allow ecf from (b) if (b) value rounds to 0.2 MV m^{-1}
		(iii)	$a = 3.7 \times 10^{-8} / 1.3 \times 10^{-6} = 0.0285$ (1) $t = 1.9 / 0.8 = 2.38$ (1) $s = 0.5 \times 0.0285 \times 2.38^2 = 0.081\text{m}$ (1)	3	If 4.4×10^{-8} N used = 0.096 m If 3.8×10^{-8} N used = 0.083 m If 4.0×10^{-8} N used = 0.087 m Accept minor variations depending on rounding in intermediate working. Allow ecf from cii if value rounds to 4×10^{-8} Accept ecf within part question
	(d)		Deflection halved (1) E-field halved (1) so force or acceleration halved (1)	3	Correct qualitative response including all three points: max 1 mark.
Total				12	

Question		Answer	Marks	Guidance
12	(a)	$4.6 \times 10^{-6} / 150 \times 31 \times 10^{-3} = 9.9 \times 10^{-7}$ (1) Wb A-turns ⁻¹ (1)	2	Accept Wb A ⁻¹ or H as the only alternative units. Accept 1×10^{-6} Beware POT errors e.g 9.9×10^{-10} Accept bald answers
	(b)	permeability = $9.9 \times 10^{-7} \times 0.27 / 2.6 \times 10^{-4}$ (1) = 1.0×10^{-3} (1)	2	Accept ecf. Accept 1×10^{-3} Accept 1.03×10^{-3} Don't accept 1.02×10^{-3} (rounding error). Accept bald answers
	(c)	(i) air has a much lower permeability	1	Accept 'a lower permeability'
		(ii) flux reduces when <u>permeance</u> reduces. (1) as current-turns value is unchanged (1) or explicit analogy between electric circuit and magnetic circuit: flux analogous to current and current- turns to p.d (1); <u>permeance</u> analogous to conductance and reduces (1)	2	
	(d)	Any two from: <ul style="list-style-type: none"> Permeability a property of material Permeability of iron remains constant Equation is not applicable as it does not account for the permeability of the air gap. 	2	Answer must relate to permeability not permeance
Total			9	

Question		Answer	Marks	Guidance
13		Any two from: Stress concentration at crack (1) Rock fails when local stress at crack exceeds fracture stress (1) Crack propagates through material (1) Under tensile load (1) (clean break with) no plastic flow (1)	2	These points may be made by annotated diagrams Accept breaking stress.
14	(a)	$t = 2\pi r / v = 3.72$ hours (or 3 hours 43 minutes) (1)	1	Answer just in seconds is not sufficient Credit any valid comparison.
	(b)	$6000 \text{ ms}^{-1} / 1\text{s}$ and $3000 \text{ ms}^{-1} / 0.5\text{s}$ (1) Both evaluated to 6000 (m), or equated (1)	2	Accept double v and double f (1) gives same λ as $v=f \lambda$ (1) Units, if given, must be correct otherwise maximum one mark
15		Transverse waves produce vibrations at right angles to direction of travel (1) These vibrations can be in two perpendicular directions to each other (1) Longitudinal waves produce vibrations in direction of travel (so in third spatial dimension) (1)	3	Correct, labelled diagrams can gain all three marks.; (-1 if T and L not labelled); (-1 if direction of travel not labelled). Confusing longitudinal waves with transverse waves scores zero.
16		$v_2 = v_1 \times (\sin \theta_2 / \sin \theta_1)$ (1) $= 6.0 \text{ km/s} \times (\sin 21 / \sin 30)$ (1) $= 4300 \text{ m s}^{-1}$ (1)	3	Correct bald answers gain 3 marks. No ecf if angles are reversed.
13 to 16 Total			11	

Question		Answer	Marks	Guidance
17	(a)	(If driving frequency of earthquakes equals that of natural frequency of seismometer then) resonance occurs (1) resulting in production of large amplitude vibrations of seismometer (1)	2	Do not credit constructive interference.
	(b)	frequency of L waves = $0.05 \text{ Hz} = 1/2\pi (g/L)^{1/2}$ (1) Re-arranging and evaluating gives $L = 99.3 \text{ m}$ (1)	2	Allow use of frequency lower than 0.05 Hz if it rounds to 0.05 Hz Do not credit 99.2 (rounding error). Or reverse calculation from 100m $f = (9.8/100)^{1/2} / 2\pi$ (1) $= 0.0498 \text{ Hz}$ (1)
	(c)	Effective length = $d / \sin \alpha$ (1) So, $\alpha = \sin^{-1} (1 / 100) = 0.6^\circ$ (1)	2	Allow ecf from (b) if value used is in range 50m to 150 m Accept 0.57° or 0.58° Allow tangent (small angle approx)
Total			6	

Question		Answer	Marks	Guidance
18	(a)	Ref to $F = ma$ (1) Small F and large m combine to give small a (1)	2	
	(b)	Displacement small so angle small (1) Force equals $mg \sin \alpha$ (1)	2	Allow evaluated acceleration based estimated displacement substituted in SHM equation (1) If force then evaluated using sensible estimated mass (1)

Question		Answer	Marks	Guidance
19	(a)	each difference of 1 is factor of 10 (1) so 100 times (1)	2	Accept $10^6 / 10^4$ (1) = 10^2 (1)
	(b)	$100^{3/2}$ (1) $= 1000$ (1)	2	Allow for ecf Accept bald answer
Question 18 & 19 Total			8	

Question		Answer	Marks	Guidance	
20	(a)	(Movement causes) change of magnetic <u>flux linkage</u> (1) Induced emf related to the rate of change of flux (linkage or in the coil) (1)	2	Accept flux in <u>coil</u> changes Accept flux (line) cutting arguments for both marks e.g. The coil cuts flux lines (1) Emf depends on rate of cutting flux lines (1).	
	(b)	(i)			
		(i)	Flux = $B \times A = 0.15 \times 4 \times 10^{-4}$ (1) $= 6 \times 10^{-5}$ (Wb) (1)	2	1 mark max if POT error
		(ii)	Flux linkage = $200 \times 6 \times 10^{-5}$ $= 0.012$ Wb turn (1)	1	Accept use of 5×10^{-5} Wb for value of flux, leading to flux linkage of 0.01 Wb Ecf from (b)(i)
		(iii)	Change in flux linkage = $0.012 / 2 = 6 \times 10^{-3}$ Wb turn (1) Time taken for change = $3 \times 10^{-3} / 1.8 \times 10^{-3} \text{ m s}^{-1}$ $= 1.7$ s (1) (Magnitude of) induced emf = $\Delta N\phi / \Delta t$ (1) $= 3.5$ mV (1)	4	Ecf from (b)(ii) and within this part of the question. Allow implicit use of equation Ignore sign Accept 4mV.. Accept 3.6 mV given by $t = 1.67$ s. Correct bald answer gains four marks. 3.5 mV, 3.6 mV, 4mV score 4 marks 7 mV, 7.1 mV, 7.2 mV score 3 marks 5.9 mV scores 3 marks
Total			9		

Question		Answer	Marks	Guidance
21		<p>movement between plate and field induces a voltage (in the plate) (1)</p> <p>This results in (eddy) currents (1)</p> <p>Currents produce magnetic fields which act to oppose the motion or change of flux which caused them (1)</p> <p>Loss of energy due to I^2R (1)</p>	4	<p>Use ticks on this question to indicate mark awards. QWC : only award 4 marks for clear linking of arguments.</p> <p>accept flux changing or flux cutting inducing a voltage.</p> <p>Accept magnetic force on induced current opposes the motion.</p> <p>Accept heating effect of current.</p>
		Total	4	
		Section C total	38	

OCR (Oxford Cambridge and RSA Examinations)
1 Hills Road
Cambridge
CB1 2EU

OCR Customer Contact Centre

Education and Learning

Telephone: 01223 553998

Facsimile: 01223 552627

Email: general.qualifications@ocr.org.uk

www.ocr.org.uk

For staff training purposes and as part of our quality assurance programme your call may be recorded or monitored

Oxford Cambridge and RSA Examinations
is a Company Limited by Guarantee
Registered in England
Registered Office; 1 Hills Road, Cambridge, CB1 2EU
Registered Company Number: 3484466
OCR is an exempt Charity

OCR (Oxford Cambridge and RSA Examinations)
Head office
Telephone: 01223 552552
Facsimile: 01223 552553

© OCR 2012

