

4752 (C2) Concepts for Advanced Mathematics

Section A

1	210 c.a.o.	2	1 for π rads = 180° soi	2
2	(i) 5.4×10^{-3} , 0.0054 or $\frac{27}{5000}$ (ii) 6 www	1 2	M1 for $S = 5.4 / (1 - 0.1)$	3
3	stretch, parallel to the y axis, sf 3	2	1 for stretch plus one other element correct	2
4	[f'(x) =] $12 - 3x^2$ their $f'(x) > 0$ or = 0 soi $-2 < x < 2$	B1 M1 A1	condone $-2 \leq x \leq 2$ or "between -2 and 2"	3
5	(i) grad of chord = $(2^{3.1} - 2^3)/0.1$ o.e. = 5.74 c.a.o. (ii) correct use of A and C where for C, $2.9 < x < 3.1$ answer in range (5.36, 5.74)	M1 A1 M1 A1	or chord with ends $x = 3 \pm h$, where $0 < h \leq 0.1$ s.c.1 for consistent use of reciprocal of gradient formula in parts (i) and (ii)	4
6	[y =] $kx^{3/2} [+ c]$ $k = 4$ subst of (9, 105) in their eqn with c or $c = -3$	M1 A1 M1 A1	may appear at any stage must have c; must have attempted integration	4
7	sector area = 28.8 or $\frac{144}{5}$ [cm ²] c.a.o. area of triangle = $\frac{1}{2} \times 6^2 \times \sin 1.6$ o.e. their sector – their triangle s.o.i. 10.8 to 10.81 [cm ²]	2 M1 M1 A1	M1 for $\frac{1}{2} \times 6^2 \times 1.6$ must both be areas leading to a positive answer	5
8	$a + 10d = 1$ or $121 = 5.5(2a + 10d)$ $5(2a + 9d) = 120$ o.e. $a = 21$ s.o.i. www and $d = -2$ s.o.i. www 4th term is 15	M1 M1 A1 A1 A1	or $121 = 5.5(a + 1)$ gets M2 eg $2a + 9d = 24$	5
9	$x \log 5 = \log 235$ or $x = \frac{\log 235}{\log 5}$ 3.39	M1 A2	or $x = \log_5 235$ A1 for 3.4 or versions of 3.392...	3
10	$2(1 - \cos^2 \theta) = \cos \theta + 2$ $-2 \cos^2 \theta = \cos \theta$ s.o.i. valid attempt at solving their quadratic in $\cos \theta$ $\cos \theta = -\frac{1}{2}$ www $\theta = 90, 270, 120, 240$	M1 A1 DM1 A1 A1	for $1 - \cos^2 \theta = \sin^2 \theta$ substituted graphic calc method: allow M3 for intersection of $y = 2 \sin^2 \theta$ and $y = \cos \theta + 2$ and A2 for all four roots. All four answers correct but unsupported scores B2. 120 and 240 only: B1.	5

18

18

Section B

11	i	$(x+5)(x-2)(x+2)$	2	M1 for $a(x+5)(x-2)(x+2)$	2
	ii	$[(x+2)](x^2+3x-10)$	M1	for correct expansion of one pair of their brackets	2
		$x^3+3x^2-10x+2x^2+6x-20$ o.e.	M1	for clear expansion of correct factors – accept given answer from $(x+5)(x^2-4)$ as first step	
	iii	$y' = 3x^2 + 10x - 4$ their $3x^2 + 10x - 4 = 0$ s.o.i. $x = 0.36\dots$ from formula o.e.	M2 M1 A1	M1 if one error or M1 for substitution of 0.4 if trying to obtain 0, and A1 for correct demonstration of sign change	6
$(-3.7, 12.6)$		B1+1			
iv	$(-1.8, 12.6)$	B1+1	accept $(-1.9, 12.6)$ or f.t. ($\frac{1}{2}$ their max x, their max y)	2	
12	i	Area = (-0.136) seen $[m^2]$ www Volume = $0.34 [m^3]$ or ft from their area $\times 2.5$	4 1	M3 for $0.1/2 \times (0.14 + 0.16 + 2[0.22 + 0.31 + 0.36 + 0.32])$ M2 for one slip; M1 for two slips must be positive	5
	ii	$2x^4 - x^3 - 0.25x^2 - 0.15x$ o.e. value at 0.5 [– value at 0] $= -0.1375$ area of cross section (of trough) or area between curve and x-axis 0.34375 r.o.t. to 3 or more sf $[m^3]$ m^3 seen in (i) or (ii)	M2 M1 A1 E1 B1 U1	M1 for 2 terms correct dep on integral attempted must have neg sign	
13	i	$\log P = \log a + b \log t$ www comparison with $y = mx + c$ intercept = $\log_{10} a$	1 1 1	must be with correct equation condone omission of base	3
	ii	$\log t$ 0 0.78 1.15 1.18 1.20 $\log P$ 1.49 1.64 1.75 1.74 1.76 plots f.t. ruled line of best fit	1 1 1 1	accept to 2 or more dp	
		iii	gradient rounding to 0.22 or 0.23 $a = 10^{1.49}$ s.o.i. $P = 31t^m$ allow the form $P = 10^{0.22 \log t + 1.49}$	2 1 1	M1 for y step / x-step accept 1.47 – 1.50 for intercept accept answers that round to 30 – 32, their positive m
	iv	answer rounds in range 60 to 63	1		1