

OXFORD CAMBRIDGE AND RSA EXAMINATIONS

**Advanced Subsidiary General Certificate of Education
Advanced General Certificate of Education**

MATHEMATICS

4733

Probability & Statistics 2

Wednesday **25 JANUARY 2006** Morning 1 hour 30 minutes

Additional materials:
8 page answer booklet
Graph paper
List of Formulae (MF1)

TIME 1 hour 30 minutes

INSTRUCTIONS TO CANDIDATES

- Write your name, centre number and candidate number in the spaces provided on the answer booklet.
- Answer **all** the questions.
- Give non-exact numerical answers correct to 3 significant figures unless a different degree of accuracy is specified in the question or is clearly appropriate.
- You are permitted to use a graphical calculator in this paper.

INFORMATION FOR CANDIDATES

- The number of marks is given in brackets [] at the end of each question or part question.
- The total number of marks for this paper is 72.
- Questions carrying smaller numbers of marks are printed earlier in the paper, and questions carrying larger numbers of marks later in the paper.
- **You are reminded of the need for clear presentation in your answers.**

This question paper consists of 3 printed pages and 1 blank page.

- 1 In a study of urban foxes it is found that on average there are 2 foxes in every 3 acres.
- (i) Use a Poisson distribution to find the probability that, at a given moment,
- (a) in a randomly chosen area of 3 acres there are at least 4 foxes, [2]
- (b) in a randomly chosen area of 1 acre there are exactly 2 foxes. [3]
- (ii) Explain briefly why a Poisson distribution might not be a suitable model. [2]
- 2 The random variable W has the distribution $B(40, \frac{2}{7})$. Use an appropriate approximation to find $P(W > 13)$. [7]
- 3 The manufacturers of a brand of chocolates claim that, on average, 30% of their chocolates have hard centres. In a random sample of 8 chocolates from this manufacturer, 5 had hard centres. Test, at the 5% significance level, whether there is evidence that the population proportion of chocolates with hard centres is not 30%, stating your hypotheses clearly. Show the values of any relevant probabilities. [7]
- 4 DVD players are tested after manufacture. The probability that a randomly chosen DVD player is defective is 0.02. The number of defective players in a random sample of size 80 is denoted by R .
- (i) Use an appropriate approximation to find $P(R \geq 2)$. [4]
- (ii) Find the smallest value of r for which $P(R \geq r) < 0.01$. [3]
- 5 In an investment model the increase, $Y\%$, in the value of an investment in one year is modelled as a continuous random variable with the distribution $N(\mu, \frac{1}{4}\mu^2)$. The value of μ depends on the type of investment chosen.
- (i) Find $P(Y < 0)$, showing that it is independent of the value of μ . [4]
- (ii) Given that $\mu = 6$, find the probability that $Y < 9$ in each of three randomly chosen years. [4]
- (iii) Explain why the calculation in part (ii) might not be valid if applied to three consecutive years. [1]
- 6 Alex obtained the actual waist measurements, w inches, of a random sample of 50 pairs of jeans, each of which was labelled as having a 32-inch waist. The results are summarised by
- $$n = 50, \quad \Sigma w = 1615.0, \quad \Sigma w^2 = 52\,214.50.$$
- Test, at the 0.1% significance level, whether this sample provides evidence that the mean waist measurement of jeans labelled as having 32-inch waists is in fact greater than 32 inches. State your hypotheses clearly. [10]

- 7 The random variable X has the distribution $N(\mu, 8^2)$. The mean of a random sample of 12 observations of X is denoted by \bar{X} . A test is carried out at the 1% significance level of the null hypothesis $H_0: \mu = 80$ against the alternative hypothesis $H_1: \mu < 80$. The test is summarised as follows: 'Reject H_0 if $\bar{X} < c$; otherwise do not reject H_0 '.

(i) Calculate the value of c . [4]

(ii) Assuming that $\mu = 80$, state whether the conclusion of the test is correct, results in a Type I error, or results in a Type II error if:

(a) $\bar{X} = 74.0$, [1]

(b) $\bar{X} = 75.0$. [1]

(iii) Independent repetitions of the above test, using the value of c found in part (i), suggest that in fact the probability of rejecting the null hypothesis is 0.06. Use this information to calculate the value of μ . [4]

- 8 A continuous random variable X has probability density function given by

$$f(x) = \begin{cases} kx^n & 0 \leq x \leq 1, \\ 0 & \text{otherwise,} \end{cases}$$

where n and k are positive constants.

(i) Find k in terms of n . [3]

(ii) Show that $E(X) = \frac{n+1}{n+2}$. [3]

It is given that $n = 3$.

(iii) Find the variance of X . [3]

(iv) One hundred observations of X are taken, and the mean of the observations is denoted by \bar{X} . Write down the approximate distribution of \bar{X} , giving the values of any parameters. [3]

(v) Write down the mean and the variance of the random variable Y with probability density function given by

$$g(y) = \begin{cases} 4\left(y + \frac{4}{5}\right)^3 & -\frac{4}{5} \leq y \leq \frac{1}{5}, \\ 0 & \text{otherwise.} \end{cases} \quad [3]$$

BLANKPAGE

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (OCR) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

OCR is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.