



## **Mathematics (MEI)**

Advanced GCE **4753**

Methods for Advanced Mathematics (C3)

# **Mark Scheme for June 2010**

---

|   |   |  |
|---|---|--|
| <b>1</b> $\begin{aligned} \int_0^{\pi/6} \cos 3x \, dx &= \left[ \frac{1}{3} \sin 3x \right]_0^{\pi/6} \\ &= \frac{1}{3} \sin \frac{\pi}{2} - 0 \\ &= 1/3 \end{aligned}$  | M1<br>$k \sin 3x, k > 0, k \neq 3$<br>B1<br>$k = (\pm)1/3$<br>A1cao [3]<br>0.33 or better | or M1 for $u = 3x \Rightarrow \int \frac{1}{3} \cos u \, du$ condone 90° in limit<br>or M1 for $\left[ \frac{1}{3} \sin u \right]$<br>so: $\sin 3x$ : M1B0, $-\sin 3x$ : M0B0,<br>$\pm 3\sin 3x$ : M0B0, $-1/3 \sin 3x$ : M0B1             |
| <b>2</b> $fg(x) =  x+1 $<br> $gf(x) =  x  + 1$<br> | B1 B1<br>graph of $ x+1 $ only<br>B1 B1<br>graph of $ x +1$<br>[4]                        | soi from correctly-shaped graphs (i.e. without intercepts)<br>but must indicate which is which<br>bod gf if negative $x$ values are missing<br>'V' shape with $(-1, 0)$ and $(0, 1)$ labelled<br>'V' shape with $(0, 1)$ labelled $(0, 1)$ |
| <b>3(i)</b> $y = (1+3x^2)^{1/2}$<br>$\Rightarrow dy/dx = \frac{1}{2}(1+3x^2)^{-1/2} \cdot 6x$<br>$= 3x(1+3x^2)^{-1/2}$  | M1<br>B1<br>A1<br>[3]   | chain rule<br>$\frac{1}{2} u^{-1/2}$<br>o.e., but must be '3'<br>can isw here  |
| <b>(ii)</b> $y = x(1+3x^2)^{1/2}$<br>$\Rightarrow dy/dx = x \cdot \frac{3x}{\sqrt{1+3x^2}} + 1 \cdot (1+3x^2)^{1/2}$<br>$= \frac{3x^2 + 1 + 3x^2}{\sqrt{1+3x^2}}$<br>$= \frac{1+6x^2}{\sqrt{1+3x^2}} *$               | M1<br>A1ft<br>M1<br>E1<br>[4]   | product rule<br>ft their $dy/dx$ from (i)<br>common denominator or factoring<br>$(1+3x^2)^{-1/2}$<br>www<br>must show this step for M1 E1  |

|   |   |  |  |
|---|---|--|--|
| <p><b>4</b></p> $p = 100/x = 100x^{-1}$ $\Rightarrow \frac{dp}{dx} = -100x^{-2} = -100/x^2$ $\frac{dp}{dt} = \frac{dp}{dx} \times \frac{dx}{dt}$ $\frac{dx}{dt} = 10$ <p>When <math>x = 50</math>, <math>\frac{dp}{dx} = (-100/50^2)</math></p> $\Rightarrow \frac{dp}{dt} = 10 \times -0.04 = -0.4$  | M1<br>A1<br>M1<br>B1<br>M1dep<br>A1cao<br>[6] | attempt to differentiate<br>$-100x^{-2}$ o.e.<br>o.e. soi<br>soi<br>substituting $x = 50$ into their $\frac{dp}{dx}$ dep 2 <sup>nd</sup> M1<br>o.e. e.g. decreasing at 0.4   | condone poor notation if chain rule correct<br>or $x = 50 + 10t$ B1<br>$\Rightarrow P = 100/x = 100/(50 + 10t)$<br>$\Rightarrow \frac{dP}{dt} = -100(50 + 10t)^{-2} \times 10 = -1000/(50 + 10t)^{-2}$ M1<br>A1<br>When $t = 0$ , $\frac{dP}{dt} = -1000/50^2 = -0.4$ A1       |
| <p><b>5</b></p> $y^3 = xy - x^2$ $\Rightarrow 3y^2 \frac{dy}{dx} = x \frac{dy}{dx} + y - 2x$ $\Rightarrow 3y^2 \frac{dy}{dx} - x \frac{dy}{dx} = y - 2x$ $\Rightarrow (3y^2 - x) \frac{dy}{dx} = y - 2x$ $\Rightarrow \frac{dy}{dx} = (y - 2x)/(3y^2 - x) *$ <p>TP when <math>\frac{dy}{dx} = 0 \Rightarrow y - 2x = 0</math></p> $\Rightarrow y = 2x$ $\Rightarrow (2x)^3 = x \cdot 2x - x^2$ $\Rightarrow 8x^3 = x^2$ $\Rightarrow x = 1/8 *(\text{or } 0)$ | B1<br>B1<br>M1<br>E1<br>M1<br>M1<br>E1<br>[7] | $3y^2 \frac{dy}{dx}$<br>$x \frac{dy}{dx} + y - 2x$<br>collecting terms in $\frac{dy}{dx}$ only<br>or $x = 1/8$ and $\frac{dy}{dx} = 0 \Rightarrow y = 1/4$<br>or $(1/4)^3 = (1/8)(1/4) - (1/8)^2$<br>or verifying e.g. $1/64 = 1/64$ | must show ' $x \frac{dy}{dx} + y$ ' on one side<br>or $x = 1/8 \Rightarrow y^3 = (1/8)y - 1/64$ M1<br>verifying that $y = 1/4$ is a solution (must show evidence*) M1<br>$\Rightarrow \frac{dy}{dx} = (1/4 - 2(1/8))/(...) = 0$ E1<br>*just stating that $y = 1/4$ is M1 M0 E0 |
| <p><b>6</b></p> $f(x) = 1 + 2 \sin 3x = y \quad x \leftrightarrow y$ $x = 1 + 2 \sin 3y$ $\Rightarrow \sin 3y = (x - 1)/2$ $\Rightarrow 3y = \arcsin [(x - 1)/2]$ $\Rightarrow y = \frac{1}{3} \arcsin \left[ \frac{x-1}{2} \right] \text{ so } f^{-1}(x) = \frac{1}{3} \arcsin \left[ \frac{x-1}{2} \right]$ <p>Range of <math>f</math> is <math>-1</math> to <math>3</math></p> $\Rightarrow -1 \leq x \leq 3$  | M1<br>A1<br>A1<br>A1<br>M1<br>A1<br>[6]       | attempt to invert<br>must be $y = \dots$ or $f^{-1}(x) = \dots$<br>or $-1 \leq (x - 1)/2 \leq 1$<br>must be 'x', not $y$ or $f(x)$   | at least one step attempted, or reasonable attempt at flow chart inversion<br>(or any other variable provided same used on each side)<br>condone ' $<$ 's for M1<br>allow unsupported correct answers; $-1$ to $3$ is M1 A0  |
| <p><b>7</b></p> <p>(A) True , (B) True , (C) False</p> <p>Counterexample, e.g. <math>\sqrt{2} + (-\sqrt{2}) = 0</math></p>  | B2,1,0<br>B1<br>[3]                           |  |  |

|  |  |   |  |
|--|--|---|--|
| 8(i) When $x = 1, y = 3 \ln 1 + 1 - 1^2 = 0$   | E1 [1]   |   |  |
| (ii) $\frac{dy}{dx} = \frac{3}{x} + 1 - 2x$<br>$\Rightarrow$ At R, $\frac{dy}{dx} = 0 = \frac{3}{x} + 1 - 2x$<br>$\Rightarrow 3 + x - 2x^2 = 0$<br>$\Rightarrow (3 - 2x)(1 + x) = 0$<br>$\Rightarrow x = 1.5, (or -1)$<br>$\Rightarrow y = 3 \ln 1.5 + 1.5 - 1.5^2 = 0.466 \text{ (3 s.f.)}$<br>$\frac{d^2y}{dx^2} = -\frac{3}{x^2} - 2$<br>When $x = 1.5, d^2y/dx^2 (= -10/3) < 0 \Rightarrow \text{max}$ | M1<br>A1cao<br><br>M1<br>M1<br>A1<br>M1<br>A1cao<br><br>B1ft<br><br>E1 [9] | d/dx ( $\ln x$ ) = $1/x$<br><br>re-arranging into a quadratic = 0<br>factorising or formula or completing square<br>substituting their $x$<br><br>ft their $dy/dx$ on equivalent work<br><br>www – don't need to calculate $10/3$ | SC1 for $x = 1.5$ unsupported, SC3 if verified<br><br>but condone rounding errors on 0.466 |
| (iii) Let $u = \ln x, du/dx = 1/x$<br>$dv/dx = 1, v = x$<br>$\Rightarrow \int \ln x dx = x \ln x - \int x \cdot \frac{1}{x} dx$<br>$= x \ln x - \int 1 dx$<br>$= x \ln x - x + c$<br><br>$\Rightarrow A = \int_1^{2.05} (3 \ln x + x - x^2) dx$<br>$= \left[ 3x \ln x - 3x + \frac{1}{2}x^2 - \frac{1}{3}x^3 \right]_1^{2.05}$<br>$= -2.5057 + 2.833..$<br>$= 0.33 \text{ (2 s.f.)}$                       | M1<br>A1<br><br>A1<br><br>B1<br><br>B1ft<br><br>M1dep<br>A1 cao [7]        | parts<br><br>condone no $c$<br><br>correct integral and limits (soi)<br><br>$\left[ 3x \ln x - 3x + \frac{1}{2}x^2 - \frac{1}{3}x^3 \right]$<br>substituting correct limits dep 1 <sup>st</sup> B1                                | allow correct result to be quoted (SC3)  |

| Mark Scheme   |   |   |  |
|---|---|---|--|
| 9(i) (0, $\frac{1}{2}$ )  | B1<br>[1]                                     | allow $y = \frac{1}{2}$ , but not $(x =) \frac{1}{2}$ or $(\frac{1}{2}, 0)$ nor P = 1/2   |  |
| (ii) $\begin{aligned} \frac{dy}{dx} &= \frac{(1+e^{2x})2e^{2x}-e^{2x}\cdot 2e^{2x}}{(1+e^{2x})^2} \\ &= \frac{2e^{2x}}{(1+e^{2x})^2} \\ \text{When } x=0, dy/dx &= 2e^0/(1+e^0)^2 = \frac{1}{2} \end{aligned}$  | M1<br>A1<br>A1<br>B1ft<br>[4]                 | Quotient or product rule<br>correct expression – condone missing bracket<br>cao – mark final answer<br>follow through their derivative  | product rule: $\begin{aligned} \frac{dy}{dx} &= e^{2x} \cdot 2e^{2x}(-1)(1+e^{2x})^{-2} + 2e^{2x}(1+e^{2x})^{-1} \\ &\quad - \frac{2e^{2x}}{(1+e^{2x})^2} \text{ from } (udv - vdu)/v^2 \text{ SC1} \end{aligned}$ |
| (iii) $\begin{aligned} A &= \int_0^1 \frac{e^{2x}}{1+e^{2x}} dx \\ &= \left[ \frac{1}{2} \ln(1+e^{2x}) \right]_0^1 \\ \text{or } &\text{ let } u = 1+e^{2x}, du/dx = 2e^{2x} \\ \Rightarrow A &= \int_2^{1+e^2} \frac{1/2}{u} du = \left[ \frac{1}{2} \ln u \right]_2^{1+e^2} \\ &= \frac{1}{2} \ln(1+e^2) - \frac{1}{2} \ln 2 \\ &= \frac{1}{2} \ln \left[ \frac{1+e^2}{2} \right] * \end{aligned}$  | B1<br>M1<br>A1<br>M1<br>A1<br>M1<br>E1<br>[5] | correct integral and limits (soi)<br>$k \ln(1+e^{2x})$<br>$k = \frac{1}{2}$<br>or $v = e^{2x}, dv/dx = 2e^{2x}$ o.e.<br>[ $\frac{1}{2} \ln u$ ] or [ $\frac{1}{2} \ln(v+1)$ ]<br>substituting correct limits<br>www | condone no dx<br>allow missing dx's or incompatible limits, but penalise missing brackets  |
| (iv) $\begin{aligned} g(-x) &= \frac{1}{2} \left[ \frac{e^{-x} - e^x}{e^{-x} + e^x} \right] = -\frac{1}{2} \left[ \frac{e^x - e^{-x}}{e^x + e^{-x}} \right] = -g(x) \\ \text{Rotational symmetry of order 2 about O} \end{aligned}$   | M1<br>E1<br>B1<br>[3]                         | substituting $-x$ for $x$ in $g(x)$<br>completion www – taking out -ve must be clear<br>must have ‘rotational’ ‘about O’, ‘order 2’ (oe)  | not $g(-x) \neq g(x)$ . Condone use of f for g.  |
| (v) (A) $\begin{aligned} g(x) + \frac{1}{2} &= \frac{1}{2} \cdot \frac{e^x - e^{-x}}{e^x + e^{-x}} + \frac{1}{2} = \frac{1}{2} \cdot \left( \frac{e^x - e^{-x} + e^x + e^{-x}}{e^x + e^{-x}} \right) \\ &= \frac{1}{2} \cdot \left( \frac{2e^x}{e^x + e^{-x}} \right) \\ &= \frac{e^x \cdot e^x}{e^x(e^x + e^{-x})} = \frac{e^{2x}}{e^{2x} + 1} = f(x) \end{aligned}$<br>(B) Translation $\begin{pmatrix} 0 \\ 1/2 \end{pmatrix}$<br>(C) Rotational symmetry [of order 2] about P | M1<br>A1<br>E1<br>M1<br>A1<br>B1<br>[6]       | combining fractions (correctly)<br>translation in y direction<br>up $\frac{1}{2}$ unit dep ‘translation’ used o.e. condone omission of $180^\circ/\text{order 2}$   | allow ‘shift’, ‘move’ in correct direction for M1.<br>$\begin{pmatrix} 0 \\ 1/2 \end{pmatrix}$ alone is SC1.   |