

GCE

Biology B (Advancing Biology)

Unit H422A/03: Practical skills in biology

Advanced GCE

Mark Scheme for June 2017

OCR (Oxford Cambridge and RSA) is a leading UK awarding body, providing a wide range of qualifications to meet the needs of candidates of all ages and abilities. OCR qualifications include AS/A Levels, Diplomas, GCSEs, Cambridge Nationals, Cambridge Technicals, Functional Skills, Key Skills, Entry Level qualifications, NVQs and vocational qualifications in areas such as IT, business, languages, teaching/training, administration and secretarial skills.

It is also responsible for developing new specifications to meet national requirements and the needs of students and teachers. OCR is a not-for-profit organisation; any surplus made is invested back into the establishment to help towards the development of qualifications and support, which keep pace with the changing needs of today's society.

This mark scheme is published as an aid to teachers and students, to indicate the requirements of the examination. It shows the basis on which marks were awarded by examiners. It does not indicate the details of the discussions which took place at an examiners' meeting before marking commenced.

All examiners are instructed that alternative correct answers and unexpected approaches in candidates' scripts must be given marks that fairly reflect the relevant knowledge and skills demonstrated.

Mark schemes should be read in conjunction with the published question papers and the report on the examination.

OCR will not enter into any discussion or correspondence in connection with this mark scheme.

© OCR 2017

Annotations

Annotation	Meaning
DO NOT ALLOW	Answers which are not worthy of credit
IGNORE	Statements which are irrelevant
ALLOW	Answers that can be accepted
()	Words which are not essential to gain credit
_	Underlined words must be present in answer to score a mark
ECF	Error carried forward
AW	Alternative wording
ORA	Or reverse argument

Subject-specific Marking Instructions

INTRODUCTION

Your first task as an Examiner is to become thoroughly familiar with the material on which the examination depends. This material includes:

- the specification, especially the assessment objectives
- the question paper
- the mark scheme.

You should ensure that you have copies of these materials.

You should ensure also that you are familiar with the administrative procedures related to the marking process. These are set out in the OCR booklet **Instructions for Examiners**. If you are examining for the first time, please read carefully **Appendix 5 Introduction to Script Marking: Notes for New Examiners**.

Please ask for help or guidance whenever you need it. Your first point of contact is your Team Leader.

C)uesti	ion	Answer	Marks	Guidance
1	(a)		2.7 ✓ mm week ⁻¹ ✓	2	 IGNORE working determined from reading a single value at 30 weeks i.e. 81/30. Candidates should use the slope of a tangent to a curve as a measure of a rate of change. ALLOW marks within range of 2.35-2.85 when calculated from a tangent ALLOW 'mm per week' or 'mm / week'
	(b)		growth of the fetus is unexpected if the value is outside the, band / range OR growth of the fetus is expected if the value is inside the, band / range ✓	1	 ALLOW 'abnormal growth' for unexpected ALLOW 'normal growth' for expected DO NOT ALLOW answers using just one set of data, e.g. if the growth rate is in the 5th percentile the fetus is growing (too) slowly ALLOW idea that 'BPD values which vary between the three sets of data indicates unexpected growth' DO NOT ALLOW references to unhealthy growth

(c)		2	 DO NOT ALLOW descriptions of methodology as these are not evaluative comments DO NOT ALLOW incorrect reference to inaccurate IGNORE references to black and white / 2D image IGNORE references to clear image/clarity of image
	One mark for useful statement ✓ One mark for idea of limitation ✓		Examples include non-invasive low risk to/safe for, fetus/mother low cost more <u>precise</u> than external measurements e.g. fundal height can monitor growth of different part of fetus' body idea of mobile equipment Examples include
			idea that produces image which can lack detail depends on correct position of fetus requires interpretation by trained medical staff resolution is low(er) (compared to other scans e.g. MRI, CT) level of detail is low(er) (compared to other scans e.g. MRI, CT) image can be blurred due to baby movement (hence the value is <u>imprecise</u>)
	Total	5	

Mark Scheme

June 2017

	Question	Answer	Marks		Guidance
2	(a)		content of the answer. level, accordi nication Stater e Communica	e answer, first ng to the Con ment has been tion Statemer	n met.

9

Level 3 (7-9 marks) Comprehensive details of apparatus and a method to produce reliable data are provided to include the preparation of dilutions of the stock solution of gibberellic acid. Most variables are identified and the method states how these are controlled. Details of a control and safety are included. *There is a well-developed line of reasoning which is*

clear and logically structured and uses scientific terminology at an appropriate level. The information presented is relevant and substantiated.

Level 2 (4-6 marks)

Some details of apparatus and a method to produce reliable data are provided. There is an outline of the preparation of dilutions of the stock solution of gibberellic acid. Some variables are identified and the method states how some are controlled. Details of a control or safety are included.

There is a line of reasoning presented with some structure and use of appropriate scientific language. The information presented in the most part relevant and supported by some evidence.

Level 1 (1–3 marks)

Apparatus and an outline method are suggested to provide some results but information, such as how to dilute the stock solution of gibberellic acid, may be missing. Some variables and safety details are omitted.

There is an attempt at a logical structure with a line of reasoning. The information is in the most part relevant.

0 marks *No response or no response worthy of credit.*

Indicative scientific points may include Apparatus & method:

- growing medium (e.g. soil, cotton wool, filter paper)
- suitable containers (e.g. pots/Petri dishes etc.) for growing medium
- apparatus for precise / accurate volume

measurement

- distilled/deionised water
- details of quantitative preparation of dilution series (for GA) to include volumes & final concentrations
- method of determining whether or not germination
 has occurred
- control included (distilled water or 0 mg dm⁻³ GA)
 - sufficient number of values of IV (different concentrations of GA across appropriate range)

Variables

- independent variable = dilutions of gibberellic acid
- dependent variable = %/proportion/fraction of seeds germinated
- control variables = exposure to light/kept in dark/constant day length, watering of seeds (volume of gibberellic acid solution is not critical but the seeds must be kept wet/moist with the appropriate solution), temperature, planting density/number of seeds per container, time

Reliability

• repeats (for each concentration)

Risk Assessment

potential chemical hazards & control


2 (b)	concentration of, gibberellin/GA ₁ , (mg dm ⁻³) in first column (ascending or descending order) AND length of radicle (mm) to the right of the IV, with each concentration of GA ₁ recorded in separate row AND mean column to the right of the DV AND informative column headings with correct unit symbols AND all cells surrounded by straight ruled lines with complete outer border ✓	4	 DO NOT ALLOW measurements in cm as this is not the most appropriate level of precision for this apparatus as stated in the Q DO NOT ALLOW if the units are in the cells of the table DO NOT ALLOW if column headed as 'average' DO NOT ALLOW if units are incorrectly formatted
	all radicles measured to nearest whole number or nearest 0.5mm AND within acceptable range ✓ three values for 50 are recorded together AND three values for 100 are recorded together AND		ALLOW error carried forward for recording values in cm if the measurements are correct
	three values for 150 are recorded together ✓ all three mean values calculated correctly from candidates own measurements AND recorded to consistent number of decimal places AND all means recorded to, the same / one more decimal place, than raw data ✓		
	Total	13	

Suggested table formats:

Concentration of GA ₁ (mg dm ⁻³)	Length of <u>radicle</u> (mm)	Mean length of radicle (mm)	Concentration of GA ₁ (mg dm ⁻³)	L	ength c (m	of <u>radi</u> m)	<u>icle</u>
		 	(9)	1	2	3	Mean
50		-	150				
100		-	100				
100		-	50				
150		-					

Acceptable ranges of values for radicles: Use data set according to method/end points used by candidate

Concentration		Length range (mm)	Length range (mm)		
of GA ₁		assuming candidate has	assuming candidate has		
	Radicle	measured to the edge of the	measured to the visible emergent point		
(mg dm ⁻³)		seed			
(ing an)		see picture 1	see picture 2		
	1	29-30	35-36		
150	4	26-28	26-28		
	8	25-26	27-30		
	2	16	19-21		
100	5	21-22	23-25		
	9	19-20	22-24		
	3	9-10	10-12		
50	6	11	17-21		
	7	18-19	22-23		



Questi	ion	Answer	Marks	Guidance
3 (a)	(i)	Mark first answer only. Any one from: idea of continuous readings ✓ idea of measured in situ / no need for taking samples out of experimental vessel ✓ greater degree of precision ✓ produces quantitative data ✓	1 max	 ALLOW apparatus remains permanently set up ALLOW no risk of further oxygen dissolving in the water as sample is processed / analysed DO NOT ALLOW greater degree of accuracy / improved accuracy
		objective / not subjective ✓ smaller % error ✓ results can be stored for later, use/processing ✓ time intervals can be varied / time intervals can be more frequent ✓ initial recording time/ start time, can be delayed ✓		DO NOT ALLOW reduces error unqualified OR doesn't make human error
	(ii)	Mark first two answers only: age (of fish) ✓ sex (of fish) ✓ mass (of fish) ✓ food, mass / quantity / type ✓ volume of / mineral ion content in / other organisms in, water ✓	2 max	DO NOT ALLOW species of fish as this is given in the question ALLOW size of fish

	(iii)	Any one from:	1 max	
		to remove, any respiring / photosynthesising microbes/microbial growth (which would interfere with respirometer readings) ✓ <i>idea of</i> to reset the respirometer to zero ✓		Microbes must be stated as either respiring or photosynthesising to gain credit
(b)	(i)	there is no (significant) difference between the <u>means</u> (metabolic rate at different temperatures/at 10 °C & 16 °C) ✓	1	
	(ii)	68.89 ✓	1	ALLOW 68.9

H422/03	
---------	--

(iii)	12.285 🗸 🗸	3	ALLOW e.c.f. throughout from 3(b)(ii) for 3 marks
			For calculations using 68.89 from Q3b(ii): ALLOW 12.291 or 12.284 (as candidate has rounded before the final the step in either case) OR 12.2849 or 12.29 for 2 marks (wrong number of d.p.) ALLOW one mark for correct substitution $t = \frac{ 86.0 - 131.6 }{\sqrt{\frac{68.89}{10} + \frac{68.89}{10}}} \qquad t = \frac{ 131.6 - 86.0 }{\sqrt{\frac{68.89}{68.89} + \frac{68.89}{10}}}$ ALLOW one mark for correct calculation (d.p. not critical here) $\frac{45.6}{\sqrt{\frac{45.6}{13.78}}} \qquad OR \qquad \frac{45.6}{3.71}$
			For calculations using 68.9 from Q3b(ii): ALLOW 12.284 OR ALLOW 12.291 or 12.284 (as candidate has rounded before the final the step in either case) OR 12.2840 or 12.28 for 2 marks (wrong number of d.p.) ALLOW one mark for correct substitution $t = \frac{ 86.0 - 131.6 }{\sqrt{\frac{68.9}{10} + \frac{68.9}{10}}} \qquad t = \frac{ 131.6 - 86.0 }{\sqrt{\frac{68.9}{10} + \frac{68.9}{10}}}$ ALLOW one mark for correct calculation (d.p. not critical here) $\frac{45.6}{\sqrt{13.78}} \qquad \text{OR} \qquad \frac{45.6}{3.71}$

H422/	03
-------	----

(iv)	3 max	ALLOW e.c.f. througout from 3(b)(iii) for 3 marks
		ALLOW 2 marks maximum if the incorrect degrees of freedom have been used
Any three from: degrees of freedom is 18 ✓		
the $t_{calculated}$ value is greater than the $t_{critical}$ value at 5% (p = 0.05) / AW \checkmark		ALLOW correct reference to using 5% probability level ALLOW 12.291 is greater than 2.101
the $t_{calculated}$ value is (also) greater than the $t_{critical}$ value at 1%/0.1% (p = 0.01/0.0001) / AW \checkmark		ALLOW 12.291 is greater than 2.878 OR 3.922
(so the researcher can) reject the null hypothesis \checkmark		
the difference in (mean) metabolic rate is not due to chance / there is a significant difference in the mean metabolic rate ✓		

(c)	Mark first two answers only.	2 max	ALLOW ORA throughout for marine fish in situ
	Any two from: For laboratory investigation: only one species of fish has been investigated ✓ only two temperatures investigated ✓ small sample size ✓ temperature affects the solubility of oxygen ✓ smaller volume of water, is less thermally stable / may vary in salinity (due to evaporation) ✓ correct reference to acclimatisation period ✓		ALLOW aquatic plants affect oxygen availability
	aquatic plants affect oxygen concentration ✓ (fish) have different food / fed at regular intervals ✓ fish in oceans will be affected by wider fluctuations in temperatures (as lab is controlled) ✓ temperature fluctuations at the ocean surface will be greater than at depths ✓ (fish naturally) have different, feeding/ habitat depths ✓ AVP ✓		e.g. idea that increasing sea surface temperatures may affect growth or populations of other marine organisms e.g. metabolic rates of fish in laboratory conditions may be affected by stress e.g. (as temperature increases) marine fish may migrate
	Total	14	

G	Quest	ion	Answer	Marks	Guidance
4	(a)	(i)	<i>Median</i> (aged 18-25) 262 AND (aged 68-75) 299 ✓	2	IGNORE units
			Range (aged 18-25) 37 AND (aged 68-75) 14 ✓		DO NOT ALLOW 258 to 295 or 297 to 311 as command word in the question states <i>calculate</i>
		(ii)	the mean, uses all the data (in the calculation) \checkmark the median, is less affected by outliers / AW \checkmark	2	IGNORE references to 'central tendency' as both averages provide this
		(iii)	Any one from (if one outlier is widely out compared with the other data) the range doesn't always give a true indication of the spread of data	1max	ALLOW 'it' for 'range'
			OR the range can be distorted by an outlier \checkmark		ALLOW outlier can give a false impression of the spread of data OWTTE
	(b)		<i>idea that</i> the data collected are from the same individual (paired measurements) ✓	2	IGNORE the 2 data sets are linked unqualified
			<i>idea of</i> allows comparison of how each individual has changed (not how the average has changed) ✓		

(c)	Summary of instructions to markers: Read through the whole answer. (Be prepared to recognise and credit unexpected approaches where they show relevance.) Using a 'best-fit' approach based on the science content of the answer, first decide which of the level descriptors, Level 1, Level 2 or Level 3, best describes the overall quality of the answer. Then, award the higher or lower mark within the level, according to the Communication Statement (shown in italics):
	 award the higher mark where the Communication Statement has been met. award the lower mark where aspects of the Communication Statement have been missed. The science content determines the level. The Communication Statement determines the mark within a level.

 Level 3 (5–6 marks) Provides a comprehensive consideration of the ethical arguments for and against the use of humans and of animals in this type of experiment. The account is well balanced There is a well-developed line of reasoning which is clear and logically structured and uses scientific terminology at an appropriate level. The information presented is relevant and substantiated Level 2 (3–4 marks) Describes some of the ethical arguments for and against the use of humans and animals in this type of experiment. The account may not be fully balanced (for & against, animals & humans). 	6	 Ethical considerations may include: For animal testing painful experiments can be carried out on animals with the use of analgesics drugs/medicines not yet approved for human use can be used on animals potentially lethal effects can be investigated in animal studies (e.g. sleep deprivation, drugs, gene deletion/'knock out') lifetime effects can be studied on animals without implications for (human) family / friends For human testing informed consent is possible for humans human volunteers can withdraw (e.g. if experiment is painful)
 use of appropriate scientific language. The information presented in the most part relevant and supported by some evidence. Level 1 (1–2 marks) Describes some of the arguments either for or against the use of humans or animals in this type of experiment. The account lacks detail or is not well balanced. There is an attempt at a logical structure with a line of reasoning. The information is in the most part relevant. O marks No response or no response worthy of credit. Total	13	 idea of conclusions/results of drug effectiveness are more applicable to humans can be used to determine effective dosage <i>Against animal testing</i> informed consent is not possible for animals animals cannot withdraw from experiments painful experiments can be carried out on animals / objections on the basis of animal cruelty idea of conclusions/results of drug effectiveness are not fully applicable to humans <i>Against human testing</i> idea that expenses compensation may affect human volunteer pool lifetime effects can have implications for, family / friends / volunteer

Mark Scheme

	Question	Answer	Marks	Guidance
5	(a)	19.85 ✓ ✓	2	ALLOW maximum of one mark for 19.849 OR 19.8 (incorrect number of decimal places)
				DO NOT ALLOW 19.9 (incorrect rounding)
				If answer is incorrect or missing ALLOW maximum of one mark for correct working
				<u>996</u> x 100 5018

H422/	03
-------	----

(b)	 (i) Analysis of data more cases of respiratory tract infections among smokers than expected OR pneumonia is more common than bronchitis among both smokers and non-smokers OR smoking appears to significantly increase the risk of bronchitis <u>and</u> may also (slightly) increase the risk of getting pneumonia ✓ 	3 ALLOW more cases of pneumonia <u>and</u> bronchitis among smokers <u>than expected</u>	
	supporting data referring to <u>expected number of cases</u> ✓	e.g. expected cases of pneumonia among smokers 297.95 or 298 OR expected cases of bronchitis among smokers 150.07 of 150 OR e.g. expected cases of pneumonia among non-smoker 1203.05 or 1203 OR expected cases of bronchitis among non-smokers 605.93 or 606	
	supporting data referring to <u>% of cases</u> ✓	23.09% or 23.1% or 23% of smokers had bronchitis compared with 13.08 or 13.1% or 13% of non-smokers OR about 33.63% or 33.6% or 34% of smokers had pneumonia compared to 28.99% or 29.0% or 29% of non-smokers	;
	<i>Comment on hypothesis</i> null hypothesis should be rejected ✓	ALLOW hypothesis is not supported IGNORE hypothesis is, false / wrong / contradicted	

Data processing for Q 5bi Calculating % values: % of smokers who have bronchitis: <u>230</u> x 100 = 23.09% (23.1%, 23%) 996 % of non-smokers who have bronchitis: <u>5260</u> x 100 = 13.08% (13.1%, 13%) 4022 % of smokers who have pneumonia: <u>335</u> x 100 = 33.63% (33.6%, 34%) 996 % of non-smokers who have pneumonia: <u>1166</u> x 100 = 28.99% (29.0%, 29%) 4022 Calculating expected values: % of population who are non-smokers = $(5018-4022) \times 100 = 19.85\%$ 5018 % of population who are non-smokers = $4022 \times 100 = 80.15\%$ 5018 Expected number of smokers who have bronchitis: <u>19.85</u> x 756 = 150.07 (150.1, 150) 100 Expected number of non-smokers who have bronchitis: 80.15 x 756 = 605.93 (605.9, 606) 100 Expected number of smokers who have pneumonia: 19.85 x 1501 = 297.95 (298.0, 298) 100 Expected number of non-smokers who have pneumonia: 80.15 x 1501 = 1203.05 (1203.1, 1203) 100

	(ii)	<u>chi-squared</u> √	1	ALLOW χ^2 test
	(iii)	(chi-squared test) tests the significance of the difference between, observed and expected results \checkmark	1	
(c)		cilia ✓	1	ALLOW ciliated epithelium ALLOW goblet cell
(d)		states apparatus, to hold solution AND measure volumes ✓	4	 Apparatus could include suitable vessels for holding solutions, e.g. test tubes volumetric apparatus, e.g. measuring cylinder, syringe, (graduated) pipette, burette
		'F' is, stock / initial, solution used in the (serial) dilution \checkmark		 ALLOW 10.0000AU in place of 'F' ALLOW description (serial) dilution of 'F' e.g. measurement of one aliquot (antibiotic solution) and, adding to / mixing with, water removal of one aliquot of new solution and, adding to / mixing with, water (repeated five times)
		correct volumes ✓		Volumes one tenth of each solution forms the aliquot and is mixed with nine aliquots of water ALLOW 'parts' in place of absolute volumes IGNORE scale of volumes
		correct sequence ✓		Sequence must result in correct dilutions being produced, i.e. start with stock solution and proceed in order

H422/03

(e)	E and F have same sized, zone of inhibition ✓ most appropriate (concentration) is, 1.0000 a.u. / E ✓ reason given for not choosing, 10.0000 a.u. / solution F OR reason given for choosing, 1.0000 a.u. / solution E ✓	3	 DO NOT ALLOW E is optimum concentration ALLOW 10.0000 a.u./ solution F, is not optimal because it is wasteful / expensive may cause (worse) side-effects greater % of antibiotic isn't absorbed in gut / AW greater % is egested OWTTE idea that it is good practice to prescribe the lowest effective dose
	Total	15	

OCR (Oxford Cambridge and RSA Examinations) 1 Hills Road Cambridge CB1 2EU

OCR Customer Contact Centre

Education and Learning

Telephone: 01223 553998 Facsimile: 01223 552627 Email: <u>general.gualifications@ocr.org.uk</u>

www.ocr.org.uk

For staff training purposes and as part of our quality assurance programme your call may be recorded or monitored

Oxford Cambridge and RSA Examinations is a Company Limited by Guarantee Registered in England Registered Office; 1 Hills Road, Cambridge, CB1 2EU Registered Company Number: 3484466 OCR is an exempt Charity

OCR (Oxford Cambridge and RSA Examinations) Head office Telephone: 01223 552552 Facsimile: 01223 552553

© OCR 2017



