

4751 (C1) Introduction to Advanced Mathematics

Section A

1	$[v =][\pm] \sqrt{\frac{2E}{m}}$ www	3	M2 for $v^2 = \frac{2E}{m}$ or for $[v =][\pm] \sqrt{\frac{E}{\frac{1}{2}m}}$ or M1 for a correct constructive first step and M1 for $v = [\pm] \sqrt{k}$ ft their $v^2 = k$; if M0 then SC1 for $\sqrt{E/ \frac{1}{2} m}$ or $\sqrt{2E/m}$ etc	3
2	$\frac{3x-4}{x+1}$ or $3 - \frac{7}{x+1}$ www as final answer	3	M1 for $(3x-4)(x-1)$ and M1 for $(x+1)(x-1)$	3
3	(i) 1 (ii) 1/64 www	1 3	M1 for dealing correctly with each of reciprocal, square root and cubing (allow 3 only for 1/64) eg M2 for 64 or -64 or $1/\sqrt[3]{4096}$ or $\frac{1}{4^3}$ or M1 for $1/16^{3/2}$ or 4^3 or -4^3 or 4^{-3} etc	4
4	$6x + 2(2x - 5) = 7$ $10x = 17$ $x = 1.7$ o.e. isw $y = -1.6$ o.e. isw	M1 M1 A1 A1	for subst or multn of eqns so one pair of coeffs equal (condone one error) simplification (condone one error) or appropriate addn/subtn to eliminate variable allow as separate or coordinates as requested graphical soln: M0	4
5	(i) -4/5 or -0.8 o.e. (ii) (15, 0) or 15 found www	2 3	M1 for 4/5 or 4/-5 or 0.8 or -4.8/6 or correct method using two points on the line (at least one correct) (may be graphical) or for -0.8x o.e. M1 for $y =$ their (i) $x + 12$ o.e. or $4x + 5y = k$ and (0, 12) subst and M1 for using $y = 0$ eg $-12 = -0.8x$ or ft their eqn or M1 for given line goes through (0, 4.8) and (6, 0) and M1 for $6 \times 12/4.8$ graphical soln: allow M1 for correct required line drawn and M1 for answer within 2mm of (15, 0)	5

6	<p>f(2) used</p> $2^3 + 2k + 7 = 3$ $k = -6$	<p>M1 M1 A1</p>	<p>or division by $x - 2$ as far as $x^2 + 2x$ obtained correctly or remainder $3 = 2(4 + k) + 7$ o.e. 2nd M1 dep on first</p>	3
7	<p>(i) 56</p> <p>(ii) -7 or ft from -their (i)/8</p>	<p>2 2</p>	<p>M1 for $\frac{8 \times 7 \times 6}{3 \times 2 \times 1}$ or more simplified M1 for 7 or ft their (i)/8 or for $56 \times (-1/2)^3$ o.e. or ft; condone x^3 in answer or in M1 expression; 0 in qn for just Pascal's triangle seen</p>	4
8	<p>(i) $5\sqrt{3}$</p> <p>(ii) common denominator = $(5 - \sqrt{2})(5 + \sqrt{2}) = 23$ numerator = 10</p>	<p>2 M1 A1 B1</p>	<p>M1 for $\sqrt{48} = 4\sqrt{3}$ allow M1A1 for $\frac{5 - \sqrt{2}}{23} + \frac{5 + \sqrt{2}}{23}$ allow 3 only for 10/23</p>	5
9	<p>(i) $n = 2m$</p> $3n^2 + 6n = 12m^2 + 12m \text{ or } = 12m(m + 1)$ <p>(ii) showing false when n is odd e.g. $3n^2 + 6n = \text{odd} + \text{even} = \text{odd}$</p>	<p>M1 M2 B2</p>	<p>or any attempt at generalising; M0 for just trying numbers or M1 for $3n^2 + 6n = 3n(n + 2) = 3 \times \text{even} \times \text{even}$ <u>and</u> M1 for explaining that 4 is a factor of even \times even or M1 for 12 is a factor of $6n$ when n is even <u>and</u> M1 for 4 is a factor of n^2 so 12 is a factor of $3n^2$ or $3n(n + 2) = 3 \times \text{odd} \times \text{odd} = \text{odd}$ or counterexample showing not always true; M1 for false with partial explanation or incorrect calculation</p>	5

Section B

10	i	correct graph with clear asymptote $x = 2$ (though need not be marked)	G2	G1 for one branch correct; condone $(0, -\frac{1}{2})$ not shown SC1 for both sections of graph shifted two to left	6	11	
	ii	$(0, -\frac{1}{2})$ shown 11/5 or 2.2 o.e. isw	G1	allow seen calculated			3
	iii	$x = \frac{1}{x-2}$ $x(x-2) = 1$ o.e. $x^2 - 2x - 1 [= 0]$; ft their equiv eqn attempt at quadratic formula $1 \pm \sqrt{2}$ cao position of points shown	2 M1 M1 M1 A1 B1	M1 for correct first step or equivs with ys or $(x-1)^2 - 1 = 1$ o.e. or $(x-1) = \pm\sqrt{2}$ (condone one error) on their curve with $y = x$ (line drawn or $y = x$ indicated by both coords); condone intent of diagonal line with gradient approx 1 through origin as $y = x$ if unlabelled			2
11	i	$(x-2.5)^2$ o.e. $-2.5^2 + 8$ $(x-2.5)^2 + 7/4$ o.e.	M1 M1 A1	for clear attempt at -2.5^2 allow M2A0 for $(x-2.5) + 7/4$ o.e. with no $(x-2.5)^2$ seen	4		
		min $y = 7/4$ o.e. [so above x axis] or commenting $(x-2.5)^2 \geq 0$	B1	ft, dep on $(x-a)^2 + b$ with b positive; condone starting again, showing $b^2 - 4ac < 0$ or using calculus			
	ii	correct symmetrical quadratic shape 8 marked as intercept on y axis tp $(5/2, 7/4)$ o.e. or ft from (i)	G1 G1 G1	or $(0, 8)$ seen in table	3		
	iii	$x^2 - 5x - 6$ seen or used -1 and 6 obtained $x < -1$ and $x > 6$ isw or ft their solns	M1 M1 M1	or $(x-2.5)^2$ [$>$ or $=$] 12.25 or ft $14 - b$ also implies first M1 if M0, allow B1 for one of $x < -1$ and $x > 6$	3		
iv	min = $(2.5, -8.25)$ or ft from (i) so yes, crosses	M1 A1	or M1 for other clear comment re translated 10 down and A1 for referring to min in (i) or graph in (ii); or M1 for correct method for solving $x^2 - 5x - 2 = 0$ or using $b^2 - 4ac$ with this and A1 for showing real solns eg $b^2 - 4ac = 33$; allow M1A0 for valid comment but error in -8.25 ft; allow M1 for showing y can be neg eg $(0, -2)$ found and A1 for correct conclusion	2	12		

12	i	$(x - 4)^2 - 16 + (y - 2)^2 - 4 = 9$ o.e. $\text{rad} = \sqrt{29}$	M2	M1 for one completing square or for $(x - 4)^2$ or $(y - 2)^2$ expanded correctly <u>or</u> starting with $(x - 4)^2 + (y - 2)^2 = r^2$: M1 for correct expn of at least one bracket and M1 for $9 + 20 = r^2$ o.e.	3
			B1	<u>or</u> using $x^2 - 2gx + y^2 - 2fy + c = 0$ M1 for using centre is (g, f) [must be quoted] and M1 for $r^2 = g^2 + f^2 - c$	
	ii	$4^2 + 2^2$ o.e. $= 20$ which is less than 29	M1 A1	allow 2 for showing circle crosses x axis at -1 and 9 or equiv for y (or showing one positive; one negative); 0 for graphical solutions (often using A and B from (iii) to draw circle)	2
	iii	showing midpt of AB = (4, 2) and showing AB = $2\sqrt{29}$ or showing AC or BC = $\sqrt{29}$ or that A or B lie on circle <u>or</u> showing both A and B lie on circle (or AC = BC = $\sqrt{29}$), and showing AB = $2\sqrt{29}$ or that C is midpt of AB or that C is on AB or that gradients of AB and AC are the same or equiv. <u>or</u> showing C is on AB and showing both A and B are on circle or AC = BC = $\sqrt{29}$	2 2 2 2	in each method, two things need to be established. Allow M1 for the concept of what should be shown and A1 for correct completion with method shown allow M1A0 for AB just shown as $\sqrt{116}$ not $2\sqrt{29}$ allow M1A0 for stating mid point of AB = (4,2) without working/method shown NB showing AB = $2\sqrt{29}$ and C lies on AB is not sufficient – earns 2 marks only	4
iv	$\text{grad AC or AB or BC} = -5/2$ o.e. $\text{grad tgt} = -1/\text{their grad AC}$ $\text{tgt is } y - 7 = \text{their } m(x - 2)$ o.e. $y = 2/5x + 31/5$ o.e.	M1 M1 M1 A1	may be seen in (iii) but only allow this M1 if they go on to use in this part allow for $m_1m_2 = -1$ used eg $y = \text{their } mx + c$ then (2, 7) subst; M0 if grad AC used condone $y = 2/5x + c$ and $c = 31/5$ o.e.	4	