



Oxford Cambridge and RSA

**GCE**

**Mathematics A**

**H240/01: Pure Mathematics**

Advanced GCE

**Mark Scheme for Autumn 2021**

OCR (Oxford Cambridge and RSA) is a leading UK awarding body, providing a wide range of qualifications to meet the needs of candidates of all ages and abilities. OCR qualifications include AS/A Levels, Diplomas, GCSEs, Cambridge Nationals, Cambridge Technicals, Functional Skills, Key Skills, Entry Level qualifications, NVQs and vocational qualifications in areas such as IT, business, languages, teaching/training, administration and secretarial skills.

It is also responsible for developing new specifications to meet national requirements and the needs of students and teachers. OCR is a not-for-profit organisation; any surplus made is invested back into the establishment to help towards the development of qualifications and support, which keep pace with the changing needs of today's society.

This mark scheme is published as an aid to teachers and students, to indicate the requirements of the examination. It shows the basis on which marks were awarded by examiners. It does not indicate the details of the discussions which took place at an examiners' meeting before marking commenced.

All examiners are instructed that alternative correct answers and unexpected approaches in candidates' scripts must be given marks that fairly reflect the relevant knowledge and skills demonstrated.

Mark schemes should be read in conjunction with the published question papers and the report on the examination.

© OCR 2021

## Text Instructions

## 1. Annotations and abbreviations

Annotation in RM assessor	Meaning
✓and ✖	
BOD	Benefit of doubt
FT	Follow through
ISW	Ignore subsequent working
M0, M1	Method mark awarded 0, 1
A0, A1	Accuracy mark awarded 0, 1
B0, B1	Independent mark awarded 0, 1
SC	Special case
^	Omission sign
MR	Misread
BP	Blank Page
Seen	
Highlighting	
Other abbreviations in mark scheme	Meaning
dep*	Mark dependent on a previous mark, indicated by *. The * may be omitted if only one previous M mark
cao	Correct answer only
oe	Or equivalent
rot	Rounded or truncated
soi	Seen or implied
www	Without wrong working
AG	Answer given
awrt	Anything which rounds to
BC	By Calculator
DR	This question included the instruction: In this question you must show detailed reasoning.

## 2. Subject-specific Marking Instructions for A Level Mathematics A

- a Annotations must be used during your marking. For a response awarded zero (or full) marks a single appropriate annotation (cross, tick, M0 or ^) is sufficient, but not required.

For responses that are not awarded either 0 or full marks, you must make it clear how you have arrived at the mark you have awarded and all responses must have enough annotation for a reviewer to decide if the mark awarded is correct without having to mark it independently.

It is vital that you annotate standardisation scripts fully to show how the marks have been awarded.

Award NR (No Response)

- if there is nothing written at all in the answer space and no attempt elsewhere in the script
- OR if there is a comment which does not in any way relate to the question (e.g. 'can't do', 'don't know')
- OR if there is a mark (e.g. a dash, a question mark, a picture) which isn't an attempt at the question.

Note: Award 0 marks only for an attempt that earns no credit (including copying out the question).

If a candidate uses the answer space for one question to answer another, for example using the space for 8(b) to answer 8(a), then give benefit of doubt unless it is ambiguous for which part it is intended.

- b An element of professional judgement is required in the marking of any written paper. Remember that the mark scheme is designed to assist in marking incorrect solutions. Correct solutions leading to correct answers are awarded full marks but work must not always be judged on the answer alone, and answers that are given in the question, especially, must be validly obtained; key steps in the working must always be looked at and anything unfamiliar must be investigated thoroughly. Correct but unfamiliar or unexpected methods are often signalled by a correct result following an apparently incorrect method. Such work must be carefully assessed. When a candidate adopts a method which does not correspond to the mark scheme, escalate the question to your Team Leader who will decide on a course of action with the Principal Examiner. If you are in any doubt whatsoever you should contact your Team Leader.

c The following types of marks are available.

**M**

A suitable method has been selected and *applied* in a manner which shows that the method is essentially understood. Method marks are not usually lost for numerical errors, algebraic slips or errors in units. However, it is not usually sufficient for a candidate just to indicate an intention of using some method or just to quote a formula; the formula or idea must be applied to the specific problem in hand, e.g. by substituting the relevant quantities into the formula. In some cases the nature of the errors allowed for the award of an M mark may be specified.

A method mark may usually be implied by a correct answer unless the question includes the DR statement, the command words “Determine” or “Show that”, or some other indication that the method must be given explicitly.

**A**

Accuracy mark, awarded for a correct answer or intermediate step correctly obtained. Accuracy marks cannot be given unless the associated Method mark is earned (or implied). Therefore M0 A1 cannot ever be awarded.

**B**

Mark for a correct result or statement independent of Method marks.

Unless otherwise indicated, marks once gained cannot subsequently be lost, e.g. wrong working following a correct form of answer is ignored. Sometimes this is reinforced in the mark scheme by the abbreviation isw. However, this would not apply to a case where a candidate passes through the correct answer as part of a wrong argument.

- d When a part of a question has two or more ‘method’ steps, the M marks are in principle independent unless the scheme specifically says otherwise; and similarly where there are several B marks allocated. (The notation ‘dep\*’ is used to indicate that a particular mark is dependent on an earlier, asterisked, mark in the scheme.) Of course, in practice it may happen that when a candidate has once gone wrong in a part of a question, the work from there on is worthless so that no more marks can sensibly be given. On the other hand, when two or more steps are successfully run together by the candidate, the earlier marks are implied and full credit must be given.
- e The abbreviation FT implies that the A or B mark indicated is allowed for work correctly following on from previously incorrect results. Otherwise, A and B marks are given for correct work only – differences in notation are of course permitted. A (accuracy) marks are not given for answers obtained from incorrect working. When A or B marks are awarded for work at an intermediate stage of a solution, there may be various alternatives that are equally acceptable. In such cases, what is acceptable will be detailed in the mark scheme. If this is not the case please, escalate the question to your Team Leader who will decide on a course of action with the Principal Examiner. Sometimes the answer to one part of a question is used in a later part of the same question. In this case, A marks will often be ‘follow through’. In such cases you must ensure that you refer back to the answer of the previous part question even if this is not shown within the image zone. You may find it easier to mark follow through questions candidate-by-candidate rather than question-by-question.
- f We are usually quite flexible about the accuracy to which the final answer is expressed; over-specification is usually only penalised where the scheme explicitly says so.
- When a value **is given** in the paper only accept an answer correct to at least as many significant figures as the given value.

- When a value **is not given** in the paper accept any answer that agrees with the correct value to **3 s.f.** unless a different level of accuracy has been asked for in the question, or the mark scheme specifies an acceptable range.

NB for Specification B (MEI) the rubric is not specific about the level of accuracy required, so this statement reads “2 s.f”.

Follow through should be used so that only one mark in any question is lost for each distinct accuracy error.

Candidates using a value of 9.80, 9.81 or 10 for  $g$  should usually be penalised for any final accuracy marks which do not agree to the value found with 9.8 which is given in the rubric.

g Rules for replaced work and multiple attempts:

- If one attempt is clearly indicated as the one to mark, or only one is left uncrossed out, then mark that attempt and ignore the others.
- If more than one attempt is left not crossed out, then mark the last attempt unless it only repeats part of the first attempt or is substantially less complete.
- If a candidate crosses out all of their attempts, the assessor should attempt to mark the crossed out answer(s) as above and award marks appropriately.

h For a genuine misreading (of numbers or symbols) which is such that the object and the difficulty of the question remain unaltered, mark according to the scheme but following through from the candidate’s data. A penalty is then applied; 1 mark is generally appropriate, though this may differ for some units. This is achieved by withholding one A or B mark in the question. Marks designated as cao may be awarded as long as there are no other errors. If a candidate corrects the misread in a later part, do not continue to follow through. Note that a miscopy of the candidate’s own working is not a misread but an accuracy error.

i If a calculator is used, some answers may be obtained with little or no working visible. Allow full marks for correct answers, provided that there is nothing in the wording of the question specifying that analytical methods are required such as the bold “In this question you must show detailed reasoning”, or the command words “Show” or “Determine”. Where an answer is wrong but there is some evidence of method, allow appropriate method marks. Wrong answers with no supporting method score zero. If in doubt, consult your Team Leader.

j If in any case the scheme operates with considerable unfairness consult your Team Leader.

Question		Answer	Marks	AO	Guidance	
1		$16 - 4(k + 3)$ $-4k - 12 + 16 > 0$ $4k - 4 < 0$  $k < 1$	<b>M1*</b> <b>A1</b> <b>M1dep*</b> <b>A1</b> <b>[4]</b>	<b>1.1</b> <b>2.3</b> <b>1.1</b> <b>1.1</b>	Attempt discriminant Obtain correct inequality Attempt to solve their inequality or equation for $k$ Obtain $k < 1$	Allow $b^2 + 4ac$ for M1, but nothing else Not necessarily expanded  <b>OR</b> (completing the square or differentiating) <b>M1*</b> – attempt to complete the square, or differentiate, and link minimum point to 0 <b>A1</b> – obtain $(k + 3) - 4 < 0$ <b>M1d*</b> – solve their inequality or equation <b>A1</b> – obtain $k < 1$  <b>OR</b> (using perfect square) <b>M1*</b> – link $k + 3$ to 4 <b>A1</b> – obtain $k + 3 < 4$ <b>M1d*</b> – solve their inequality or equation <b>A1</b> – obtain $k < 1$
2	(a)	$(C =) 4000 + 4m$ $(C =) 6(m - 100)$	<b>B1</b> <b>B1</b> <b>[2]</b>	<b>3.3</b> <b>3.3</b>	Correct equation / expression for $A$ Correct equation / expression for $B$	Or $40 + 0.04m$ Or $0.06(m - 100)$ <b>B1B0</b> if units inconsistent in two equations <b>SC B1</b> for both $44 + 0.04m$ and $0.06m$ (or $4400 + 4m$ and $6m$ ) – from using $m = 0$ at 100 minutes
	(b)	$4000 + 4m = 6(m - 100)$ $2m = 4600$  $m = 2300$	<b>M1</b>  <b>A1</b> <b>[2]</b>	<b>1.1</b>  <b>3.4</b>	Attempt to solve simultaneously, from two linear equations in $m$  Obtain 2300 (minutes)	At least one equation must have constant term Could be implied by final answer of 38hrs 20 mins isw once 2300 seen

Question		Answer	Marks	AO	Guidance	
3	(a)	$x = ky^2\sqrt{z}$ $30 = k \times 4 \times 3$ $k = 2.5$  $x = 2.5y^2\sqrt{z}$	M1	3.1a	Attempt to find value for $k$	From $x = ky^2\sqrt{z}$ or $x = kz^2\sqrt{y}$ only Using sum, not product, is M0 but watch for $+\sqrt{z}$ being used for positive square root Ignore modulus sign if used around $\sqrt{z}$ Allow BOD if initial equation stated explicitly, $k$ found correctly but then final equation not seen or seen as now incorrect
			A1	1.1	Correct equation	
	(b)	$x = 2.5 \times 9 \times 5$  $x = 112.5$	M1	1.1	Attempt to find $x$ , from equation in terms of $y, z$ and numerical $k$	Could be from using direct proportion and not their equation from (a)
			A1	1.1	Obtain 112.5	Or any exact equiv
		[2]	[2]			
4	(a)	<b>DR</b> $f(0.5) = 0.25 - 0.75 - 5.5 + 6 = 0$	B1	2.1	Attempt $f(0.5)$ and show equal to 0 Must be using factor theorem so B0 for alternative methods	B0 for just $f(0.5) = 0$ Condone $2(0.5)^3 - 3(0.5)^2 - 11(0.5) + 6 = 0$
		[1]	[1]			
	(b)	<b>DR</b> $f(x) = (2x - 1)(x^2 - x - 6)$	M1	1.1	Attempt complete division by $(2x - 1)$	<b>DR</b> so need to see quadratic factor Allow equivalent complete methods eg coefficient matching / inspection / grid method Condone slip(s) in otherwise correct method
			A1	1.1	Obtain correct quadratic factor	Seen in division / correct coeffs eg $A = 1$ etc / at top of grid



Question			Answer	Marks	AO	Guidance	
			$f(x) = (2x - 1)(x - 3)(x + 2)$	<b>A1</b> [3]	<b>1.1</b>	Obtain correct fully factorised $f(x)$	Must be seen as a product of all 3 factors <b>SC B1</b> for correct factorisation with no DR
	<b>(c)</b>		<b>DR</b> $x = 2^y$  $2^y = 0.5, y = -1$ $2^y = 3, y = 1.58$ $2^y = -2$ , no solutions as $2^y > 0$ for all $y$ Hence $y = -1, y = \log_2 3$	<b>B1</b>  <b>M1</b>  <b>A1</b>  [3]	<b>3.1a</b>  <b>1.1</b>  <b>2.4</b>	State or imply that $x = 2^y$  Attempt to find at least one value of $y$ Obtain both correct values, and no others  Must give <b>reason</b> for $2^y = -2$ having no solution	Could be implied by equating $2^y$ to at least one of their roots Exact or decimal  $1.58$ or better, or $\frac{\log_n 3}{\log_n 2}$ for $\log_2 3$  eg cannot log a negative number $2^y$ always greater than 0
<b>5</b>	<b>(a)</b>	<b>(i)</b>	(2, 9)	<b>B1</b> [1]	<b>1.1</b>	Correct coordinate	And no others
		<b>(ii)</b>	(1, 12)	<b>B1</b> <b>B1</b> [2]	<b>3.1a</b> <b>1.1</b>	Correct $x$ -coordinate Correct $y$ -coordinate	If more than one solution given then award B1 if either co-ordinate is consistent in all solutions
		<b>(iii)</b>	(6, 2)	<b>B1</b> [1]	<b>1.2</b>	Correct coordinate	And no others
	<b>(b)</b>	<b>(i)</b>	$x = -2, x = 4$	<b>B1</b> [1]	<b>3.1a</b>	Both $x$ -coordinates correct, and no others	Ignore any attempt at $y$ values
		<b>(ii)</b>	$x < -2$	<b>B1</b> [1]	<b>1.2</b>	Correct inequality, and no others	Allow $\leq$ Could be written in set notation
		<b>(iii)</b>	$x = 4$	<b>B1</b>	<b>1.2</b>	Correct $x$ -coordinate, and no others	Ignore any attempt at $y$ values

Question		Answer	Marks	AO	Guidance	
			[1]			
6	(a)	$\left(1 - \frac{3}{8}x\right)^{\frac{1}{3}} = 1 + \left(\frac{1}{3}\right)\left(-\frac{3}{8}x\right) + \frac{\left(\frac{1}{3}\right)\left(-\frac{2}{3}\right)\left(-\frac{3}{8}x\right)^2}{2!}$ $= 1 - \frac{1}{8}x - \frac{1}{64}x^2$ $(8 - 3x)^{\frac{1}{3}} = 8^{\frac{1}{3}}\left(1 - \frac{3}{8}x\right)^{\frac{1}{3}} = 2\left(1 - \frac{3}{8}x\right)^{\frac{1}{3}}$ $(8 - 3x)^{\frac{1}{3}} = 2 - \frac{1}{4}x - \frac{1}{32}x^2$	<b>B1</b>  <b>M1</b>  <b>A1</b>  <b>B1FT</b>  <b>[4]</b>	<b>1.1</b>  <b>1.1</b>  <b>1.1</b>  <b>1.1</b>	Obtain correct first two terms  Attempt third term in expansion of $\left(1 - \frac{3}{8}x\right)^{\frac{1}{3}}$  Correct third term  Correct expansion of $(8 - 3x)^{\frac{1}{3}}$	Allow unsimplified second term, including product of two fractions  Allow BOD if no brackets, even if never recovered Allow BOD if no negative sign Allow unsimplified fraction as coefficient, but must be single term FT as 2 x their expansion (at least two terms) Bracket expanded and fractions simplified
	(b)	$ x  < \frac{8}{3}$	<b>B1</b>  <b>[1]</b>	<b>1.2</b>	Allow any equivalent eg $-\frac{8}{3} < x < \frac{8}{3}$  Must be strict inequality Must be condition for $x$ , so B0 for $ 3x  < 8$	
	(c)	$(1 + 2x)^{-2} = 1 + (-2)(2x) + \frac{(-2)(-3)}{2!}(2x)^2$ $= 1 - 4x + 12x^2$ $(2 \times 12) + \left(-\frac{1}{4} \times -4\right) + \left(-\frac{1}{32} \times 1\right)$	<b>M1</b>  <b>A1</b>  <b>M1</b>	<b>3.1a</b>  <b>1.1</b>  <b>1.1</b>	Attempt first three terms of expansion  Obtain correct first three terms  Attempt all 3 relevant products	Must be expanding $(1 + 2x)^{-2}$ Allow BOD if no brackets on $2x$ , even if never recovered Allow unsimplified fraction for coeff of third term Finding 3 appropriate terms from the product of two 3-term quadratics If part of full expansion then M1 when reqd 3 products and no others are combined

Question		Answer	Marks	AO	Guidance
		$\frac{799}{32}$ or $24\frac{31}{32}$	<b>A1</b>  <b>[4]</b>	<b>1.1</b>	Any exact equivalent, including 24.96875  Condone $x^2$ still present
7	(a)	$2x \ln x + \frac{x^2 - 2}{x}$  $2x \ln x + \frac{x^2 - 2}{x} = 0$  $2x^2 \ln x + x^2 - 2 = 0$ <b>A.G.</b>	<b>M1</b>  <b>A1</b>  <b>[2]</b>	<b>3.1a</b>  <b>1.1</b>	Attempt differentiation using product rule  Equate to 0 and obtain given answer  May expand first to give $2x \ln x + \frac{x^2}{x} - \frac{2}{x}$ (allow middle term as just $x$ ) Must be equated to 0 <b>before</b> clearing the fractions Must be equation ie ... = 0
	(b)	$f'(x) = 4x \ln x + 2x^2 \cdot \frac{1}{x} + 2x$  $x_{n+1} = x_n - \frac{2x_n^2 \ln x_n + x_n^2 - 2}{4x_n \ln x_n + 2x_n^2 \cdot \frac{1}{x_n} + 2x_n}$  $x_{n+1} = \frac{x_n (4x_n \ln x_n + 4x_n) - (2x_n^2 \ln x_n + x_n^2 - 2)}{4x_n \ln x_n + 4x_n}$  $x_{n+1} = \frac{4x_n^2 \ln x_n + 4x_n^2 - 2x_n^2 \ln x_n - x_n^2 + 2}{4x_n \ln x_n + 4x_n}$  $x_{n+1} = \frac{2x_n^2 \ln x_n + 3x_n^2 + 2}{4x_n (\ln x_n + 1)}$ <b>A.G.</b>	<b>B1</b>  <b>M1</b>  <b>M1</b>  <b>A1</b>  <b>[4]</b>	<b>1.1</b>  <b>1.1</b>  <b>1.1</b>  <b>2.1</b>	Correct derivative seen  Use correct Newton-Raphson formula, with numerator correct and their derivative in the denominator  Attempt rearrangement into single fraction with brackets expanded  Obtain given answer, with no errors seen  Allow simplified middle term of $2x$ Allow fractional term without subscripts <b>SC</b> Condone use of N-R on $(x^2 - 2) \ln x$  Allow without subscripts N-R not necessarily correct, but must be recognisable attempt <b>SC</b> Rearrange their N-R on $(x^2 - 2) \ln x$  Subscripts needed on RHS at least one step before AG LHS needs $x_{n+1}$ seen
	(c)	$x_2 = 1.25, x_3 = 1.2075$	<b>B1</b>  <b>[1]</b>	<b>1.1</b>	Condone 1.21, or better, for $x_3$  $x_3 = 1.207515437\dots$

Question		Answer	Marks	AO	Guidance	
	(d)	(1.206, -0.102)	<b>B1</b> <b>B1</b>	<b>2.2a</b> <b>2.2a</b>	Correct $x$ -coordinate Correct $y$ -coordinate	Must be 3dp or better Could be given as single coordinate or $x = 1.206, y = -0.102$ Allow BOD if 1.206 given but not identified as $x$ -value
			[2]			
8	(a)	(i)	<b>B1</b>	<b>2.5</b>	Allow alternative notation, or worded equivalent Allow $y$ , or just $f$ , but not $x$	Accept just $\square$ Allow $(-\infty, \infty)$
		(ii)	<b>B1</b>	<b>2.5</b>	Allow alternative notation, or worded equivalent Allow $y$ , or just $g$ , but not $x$	Or $(-\infty, \infty)$ with $(-1, 1)$ clearly excluded
			[1]			
	(b)	(i)	<b>M1</b>  <b>A1</b> <b>[2]</b>	<b>2.1</b>  <b>2.1</b>	Attempt correct composition of functions  Conclude with 5.33	At least one interim value required  <b>SC B1</b> for stating $2\tan(1 \div \cos 0.6) = 5.33$
		(ii)	<b>B1</b>	<b>2.4</b>	Must refer to inverse of $f$ not existing, with reason	Must be clear that referring to the function $f$

Question		Answer	Marks	AO	Guidance
			[1]		
(c)	<p><b>DR</b></p> $4\tan^2x + 6\secx = 0$ $4(\sec^2x - 1) + 6\secx = 0$ $4\sec^2x + 6\secx - 4 = 0$ $\secx = -2, \secx = \frac{1}{2}$ $x = \frac{2}{3}\pi, \frac{4}{3}\pi$	<p><b>M1</b></p> <p><b>A1</b></p> <p><b>M1</b></p> <p><b>A1</b></p>	<p><b>3.1a</b></p> <p><b>1.1</b></p> <p><b>1.1</b></p> <p><b>1.1</b></p>	<p>Attempt use of identity in their equation to obtain quadratic</p> <p>Obtain correct equation in <math>\secx</math> – possibly still with brackets</p> <p>Solve 3 term quadratic and attempt to find at least one value for <math>x</math></p> <p>Obtain at least one correct value</p>	<p>Allow <math>\tan^2x = \pm \sec^2x \pm 1</math></p> <p>Award M1 when reduced to single trig ratio</p> <p>Or correct quadratic in <math>\cosx</math> – possibly still with brackets but with no fractions (<math>4\cos^2x - 6\cosx - 4 = 0</math>)</p> <p>Could solve quadratic BC</p> <p>Must be using root that would give solution for <math>x</math></p> <p>Allow decimals or in degrees</p> <p>Must be from correct solution method of correct quadratic (condone second root not being seen – but must be correct if seen)</p>

Question		Answer	Marks	AO	Guidance	
		$\sec x = \frac{1}{2}$ has no solutions as $ \sec x  \geq 1$	<b>A1</b>  <b>[5]</b>	<b>2.3</b>	Obtain both correct values, and no others, and explain that $\sec x = 0.5$ has no solutions as outside range	Now exact and in radians Or equiv explanation for $\cos x$

<b>9</b>	<b>(a)</b>	$e^{kt} > 0$ for all $t$ , so $y > 0$ for $t \geq 0$ (or for all $t$ ) hence never crosses $x$ -axis	<b>B1</b>  <b>[1]</b>	<b>2.4</b>	Or show that $2e^{-3t} = 0$ has no solutions	Need to see $y \neq 0$ or $y > 0$ and reason relating to exponential (or logarithmic) function Must clearly be referring to $y$ or $2e^{-3t}$
	<b>(b)</b>	$e^{2t} - 4e^t + 3 = 0$ $(e^t - 1)(e^t - 3) = 0$ $e^t = 1, e^t = 3$  $t = 0, t = \ln 3$	<b>M1</b>  <b>A1</b> <b>[2]</b>	<b>3.1a</b>  <b>1.1</b>	Equate to 0 and attempt to solve disguised quadratic  Obtain both correct values	‘determine’ so some evidence of method needed  A0 for $\ln 1$ and not 0
	<b>(c)</b>	$\frac{dx}{dt} = 2e^{2t} - 4e^t$	<b>B1</b>	<b>1.1</b>	Correct $\frac{dx}{dt}$	Mark derivative and condone no/wrong label

Question			Answer	Marks	AO	Guidance	
			$\frac{dy}{dt} = -6e^{-3t}$ $\frac{dy}{dx} = \frac{-6e^{-3t}}{2e^{2t} - 4e^t}$ $\frac{dy}{dx} = \frac{-3e^{-3t}}{e^{2t} - 2e^t} = \frac{3}{e^{3t}(2e^t - e^{2t})} = \frac{3}{2e^{4t} - e^{5t}}$ <b>A.G.</b>	<b>B1</b> <b>M1</b> <b>A1</b>  <b>[4]</b>	<b>1.1</b> <b>2.4</b> <b>2.1</b>	Correct $\frac{dy}{dt}$ Attempt correct method to combine derivatives Show manipulation to given answer	Mark derivative and condone no/wrong label Combine their derivatives correctly Need to see some evidence of how $e^{-3t}$ is dealt with <b>AG</b> so method must be fully correct
	<b>(d)</b>		$2e^{4t} - e^{5t} = 0$ $e^{4t}(2 - e^t) = 0$ $t = \ln 2$ $(-1, \frac{1}{4})$	<b>M1</b>  <b>A1</b>  <b>A1</b>  <b>[3]</b>	<b>3.1a</b>  <b>1.1</b>  <b>1.1</b>	Equate denominator to 0 Solve for $t$ to obtain $t = \ln 2$ Obtain correct coordinate	Or $\frac{dx}{dy} = 0$ or $\frac{dx}{dt} = 0$ No need to see $e^{4t} = 0$ discounted Or $x = -1, y = \frac{1}{4}$
<b>10</b>	<b>(a)</b>	<b>(i)</b>	$\cos y = \frac{CD}{a}$ hence $CD = a \cos y$	<b>B1</b>  <b>[1]</b>	<b>2.4</b>	Justification for $CD$	Need to see either $\cos y = \frac{CD}{a}$ or $\text{adj} = \text{hyp} \times \cos \theta$ before given answer
		<b>(ii)</b>	$\text{area} = \frac{1}{2} AC \cdot CD \sin x = \frac{1}{2} b(a \cos y) \sin x$ $= \frac{1}{2} ab \sin x \cos y$ <b>A.G.</b>	<b>B1</b>  <b>[1]</b>	<b>2.4</b>	Use area of triangle to show given answer	Could quote general expression for area and then show clear substitution If not, then sides being used need to be clearly identified through statement or diagram Could also use right-angled triangle, with base as $AD$ Condone not being rearranged to given expression
		<b>(iii)</b>	$CD = b \cos x$	<b>B1</b>	<b>2.1</b>	Correct $CD$ in terms of $b$ and $x$	

Question	Answer	Marks	AO	Guidance
	Area $BCD =$ $\frac{1}{2}BC \cdot CD \sin y = \frac{1}{2}a(b \cos x) \sin y$ $= \frac{1}{2}ab \cos x \sin y$ Area $ABC =$ $\frac{1}{2}AC \cdot BC \sin(x + y) = \frac{1}{2}ab \sin(x + y)$ $\frac{1}{2}ab \sin(x + y) = \frac{1}{2}ab \sin x \cos y + \frac{1}{2}ab \cos x \sin y$ $\sin(x + y) = \sin x \cos y + \cos x \sin y$	<b>B1</b>  <b>B1</b>  <b>B1</b>  <b>[4]</b>	<b>2.1</b>  <b>1.1</b>  <b>2.1</b>	Correct area of triangle $BCD$  Correct area of triangle $ABC$  Equate area of $ABC$ to the sum of the areas of the two small triangles and complete proof convincingly  Allow alternative proofs eg using lengths  B0 B1 if correct area stated with no justification
<b>(b)</b>	$\sin 30 \cos \alpha + \cos 30 \sin \alpha =$ $\cos 45 \cos \alpha + \sin 45 \sin \alpha$ $\frac{1}{2} \cos \alpha + \frac{1}{2} \sqrt{3} \sin \alpha = \frac{1}{2} \sqrt{2} \cos \alpha + \frac{1}{2} \sqrt{2} \sin \alpha$ $(\sqrt{3} - \sqrt{2}) \sin \alpha = (\sqrt{2} - 1) \cos \alpha$ $\frac{\sin \alpha}{\cos \alpha} = \tan \alpha = \frac{\sqrt{2} - 1}{\sqrt{3} - \sqrt{2}}$ $\frac{(\sqrt{2} - 1)(\sqrt{3} + \sqrt{2})}{(\sqrt{3} - \sqrt{2})(\sqrt{3} + \sqrt{2})} = \frac{\sqrt{6} + 2 - \sqrt{2} - \sqrt{3}}{3 - 2}$ $\tan \alpha = 2 + \sqrt{6} - \sqrt{3} - \sqrt{2}$ <b>A.G.</b>	<b>B1</b>  <b>M1</b>  <b>M1</b>  <b>M1</b>  <b>A1</b>	<b>1.1</b>  <b>1.1</b>  <b>3.1a</b>  <b>3.1a</b>  <b>2.1</b>	Correct use of compound angle formulae  Use exact trig values  Gather like terms and attempt $\tan \alpha$ May still have fractions in the fraction  Attempt to rationalise their denominator  Obtain given answer  Could be implied if exact values used immediately – allow BOD for RHS May be seen as two separate expressions, not yet equated In either equation or two expressions Must see all 4 values, but expansions may not be fully correct $\tan \alpha$ does not yet need to be the subject, but must only appear once for M1  Clear intention seen to multiply throughout by the conjugate of their denominator  With full detail, including (at least) $3 - 2$ in denominator



Question		Answer	Marks	AO	Guidance
11	(a)		[5]		
		$2udu = 2xdx$	B1	1.1a	Any correct expression linking $du$ and $dx$
		$\int \frac{4u(u^2 - 3)}{\sqrt{u^2}} du$	M1*	2.1	Attempt to rewrite integrand in terms of $u$
		$\int (4u^2 - 12) du$	A1	1.1	Obtain correct integrand
		$\frac{4}{3}u^3 - 12u(+c)$	M1dep*	1.1	Attempt integration
		$\frac{4}{3}u(u^2 - 9) + c = \frac{4}{3}(x^2 - 6)\sqrt{x^2 + 3} + c$ A.G.	A1	2.1	Obtain given answer, with at least one intermediate step seen
			[5]		
	(b)	DR $\frac{4}{3}((-5 \times 2) - (-6 \times \sqrt{3}))$	M1	2.1	Attempt to use limits $x = 0$ and $x = 1$ , or $u = \sqrt{3}$ and $u = 2$ in integral in terms of $u$
		$= \frac{4}{3}(6\sqrt{3} - 10)$ or 0.523	A1	1.1	Obtain correct area under curve
		$\frac{dy}{dx} = \frac{12x^2(x^2 + 3)^{\frac{1}{2}} - 4x^3 \cdot 2x \cdot \frac{1}{2}(x^2 + 3)^{-\frac{1}{2}}}{x^2 + 3}$	M1	3.1a	Attempt derivative using the quotient rule
					Correct order and subtraction Attempt to use both limits in their integral to give two terms DR so just stating decimal area is M0 Either using answer from (a) or their integration attempt with +2 or +3 Accept exact (inc unsimplified) or decimal Using +2 gives $\frac{4}{3}(4\sqrt{2} - 3\sqrt{3})$ or 0.614 Or equiv with product rule Need difference of two terms in numerator, at least one term correct, but allow subtraction in incorrect order Using either +2 or +3 equation

Question			Answer	Marks	AO	Guidance	
			at $x = 1$ , $m = \frac{11}{2}$ hence $m' = -\frac{2}{11}$  $y - 2 = -\frac{2}{11}(x - 1)$ when $y = 0$ , $x = 12$  $\text{area} = 8\sqrt{3} - \frac{40}{3} + 11$ $= 8\sqrt{3} - \frac{7}{3}$	<b>A1</b>  <b>M1</b>  <b>M1</b>  <b>A1</b>	<b>1.1</b>  <b>2.1</b>  <b>1.1</b>  <b>3.1a</b>	Obtain correct, unsimplified, derivative Attempt gradient of normal at $x = 1$  Attempt to find point of intersection of normal with $x$ -axis  Obtain correct area Allow any exact (including unsimplified) or decimal equivalent	With either +2 or +3  Substitute $x = 1$ and use negative reciprocal Using +2 gives $m' = -\frac{3}{32}\sqrt{3}$ Can be with $m$ found BC Attempt equation of normal with their gradient and either $(1, 2)$ or $(1, \frac{4}{3}\sqrt{3})$ , and then use $y = 0$ to find $x$ intersection From combining a correct area under curve and a correct area of triangle (either 11 or $\frac{64}{9}\sqrt{3}$ ), even if inconsistent Can still get A1 following M0 for area under curve BC and/or $m$ found BC
<b>12</b>	<b>(a)</b>	<b>(i)</b>	$\frac{d\theta}{dt} = -k$	<b>B1</b>  <b>[1]</b>	<b>3.3</b>	Allow $\frac{d\theta}{dt} = k$ or $\frac{d\theta}{dt} = -3.5$  Both sides of differential equation required	
		<b>(ii)</b>	$\theta = -3.5t + c$  $\theta = 160 - 3.5t$	<b>M1</b>  <b>A1</b> <b>[2]</b>	<b>3.4</b>  <b>1.1</b>	Obtain equation of the form $\theta = \pm 3.5t + c$ , where $c$ could already be numerical and possibly incorrect Obtain correct equation  <b>Alt method</b> For M1, integrate to get $\theta = kt + c$ , then use $(0, 160)$ and $(10, 125)$ to attempt $c$ and hence $k$	

Question		Answer	Marks	AO	Guidance	
	(iii)	The model would predict that the temperature would fall below room temperature, and eventually below freezing point	<b>B1</b>  <b>[1]</b>	<b>3.5b</b>	Any sensible comment	Cooling rate unlikely to be linear Identify that limit (ie room temperature) will be reached

	(b)	(i)	$\frac{d\theta}{dt} = -k(\theta - 20)$	<b>B1</b>  <b>[1]</b>	<b>3.3</b>	Allow $\frac{d\theta}{dt} = k(\theta - 20)$	Both sides of differential equation required ISW if $k = -3.5$ used once correct equation seen (but B0 if only ever seen with $-3.5$ )
		(ii)	$\int \frac{1}{\theta - 20} d\theta = \int -k dt$	<b>M1</b>	<b>3.1a</b>	Separate variables (or invert each side) and attempt integration	Allow M1 for integration of a differential equation not of this form eg $\frac{d\theta}{dt} = \frac{-k}{(\theta - 20)}$ , as long as $t$ and/or $\theta$ are involved – must be attempt at correct rearrangement of their diff eqn

Question	Answer	Marks	AO	Guidance
	$\ln \theta - 20  = -kt + c$ $\ln 140 = c$ $\ln 105 = -10k + \ln 140$ $k = -0.1 \ln 0.75$ $\ln \theta - 20  = (0.1 \ln 0.75)t + \ln 140$ $\theta - 20 = e^{(0.1 \ln 0.75)t + \ln 140} \quad 140e^{(0.1 \ln 0.75)t}$ $\theta = 20 + 140e^{(0.1 \ln 0.75)t}$	<p><b>A1</b></p> <p><b>M1</b></p> <p><b>M1</b></p> <p><b>M1</b></p> <p><b>A1</b></p> <p><b>[6]</b></p>	<p><b>1.1</b></p> <p><b>3.4</b></p> <p><b>1.1a</b></p> <p><b>1.1</b></p> <p><b>1.1</b></p>	<p>Obtain correct integral</p> <p>Use <math>t = 0, \theta = 160</math> in an equation involving both <math>k</math> and <math>c</math></p> <p>Use <math>t = 10, \theta = 125</math> in an equation involving both <math>k</math> and <math>c</math> (<math>c</math> possibly now numerical)</p> <p>Attempt to make <math>\theta</math> the subject</p> <p>Obtain correct equation Allow <math>-0.0288</math> (or better) for <math>0.1 \ln 0.75</math> and/or <math>4.94</math> (or better) for <math>\ln 140</math></p> <p>Or <math>\ln \theta - 20  = kt + c</math> Condone brackets not modulus Equation must be from integration attempt, but could follow M0 As far as numerical <math>c</math> or <math>k</math> Using both pairs of values as limits in a definite integral is M2 As far as numerical <math>c</math> and <math>k</math></p> <p>As far as correctly removing logs Equation must now be of the correct form ie <math>\ln a\theta + b  = ct + d</math> Could still be in terms of <math>c</math> and <math>k</math> to give eg <math>\theta = Ae^{kt} + 20</math> Allow <math>\theta = 20 + e^{(0.1 \ln 0.75)t + \ln 140}</math> Could see <math>\theta = 140(0.75)^{0.1t} + 20</math></p>
(c)	$25 = 160 - 3.5t \Rightarrow t = 38.6 \text{ mins}$ $\ln 5 = (0.1 \ln 0.75)t + \ln 140 \Rightarrow t = 115.8 \text{ mins}$ <p>77 minutes</p>	<p><b>M1</b></p> <p><b>A1</b></p> <p><b>[2]</b></p>	<p><b>3.4</b></p> <p><b>3.4</b></p>	<p>Use <math>\theta = 25</math> in both of their equations to find values for <math>t</math></p> <p>Obtain 77 minutes</p> <p>As far as two numerical values for <math>t</math></p> <p>Accept any answer rounding to 77, with no errors seen</p>

**OCR (Oxford Cambridge and RSA Examinations)**  
**The Triangle Building**  
**Shaftesbury Road**  
**Cambridge**  
**CB2 8EA**

**OCR Customer Contact Centre**

**Education and Learning**

Telephone: 01223 553998

Facsimile: 01223 552627

Email: [general.qualifications@ocr.org.uk](mailto:general.qualifications@ocr.org.uk)

[www.ocr.org.uk](http://www.ocr.org.uk)

For staff training purposes and as part of our quality assurance programme your call may be recorded or monitored