

**Mark Scheme 4753**  
**January 2007**



## Section A

1 (i) P is (2, 1)	B1	
(ii) $ x  = 1\frac{1}{2}$ $\Rightarrow x = (-1\frac{1}{2}) \text{ or } 1\frac{1}{2}$ $ x - 2  + 1 = 1\frac{1}{2} \Rightarrow  x - 2  = \frac{1}{2}$ $\Rightarrow x = (2\frac{1}{2}) \text{ or } 1\frac{1}{2}$	M1 A1  M1 E1	allow $x = 1\frac{1}{2}$ unsupported  or $ 1\frac{1}{2} - 2  + 1 = \frac{1}{2} + 1 = 1\frac{1}{2}$
or by solving equation directly: $ x - 2  + 1 =  x $ $\Rightarrow 2 - x + 1 = x$ $\Rightarrow x = 1\frac{1}{2}$ $\Rightarrow y =  x  = 1\frac{1}{2}$	M1 M1 A1 E1 [4]	equating from graph or listing possible cases
2 $\int_1^2 x^2 \ln x dx$ $u = \ln x$ $dv/dx = x^2 \Rightarrow v = \frac{1}{3}x^3$ $= \left[ \frac{1}{3}x^3 \ln x \right]_1^2 - \int_1^2 \frac{1}{3}x^3 \cdot \frac{1}{x} dx$ $= \frac{8}{3} \ln 2 - \int_1^2 \frac{1}{3}x^2 dx$ $= \frac{8}{3} \ln 2 - \left[ \frac{1}{9}x^3 \right]_1^2$ $= \frac{8}{3} \ln 2 - \frac{8}{9} + \frac{1}{9}$ $= \frac{8}{3} \ln 2 - \frac{7}{9}$	M1 A1  A1  M1  A1 cao [5]	Parts with $u = \ln x$ $dv/dx = x^2 \Rightarrow v = x^3/3$  $\left[ \frac{1}{9}x^3 \right]$ substituting limits  o.e. – not $\ln 1$
3 (i) When $t = 0$ , $V = 10\ 000$ $\Rightarrow 10\ 000 = Ae^0 = A$  When $t = 3$ , $V = 6000$ $\Rightarrow 6000 = 10\ 000 e^{-3k}$ $\Rightarrow -3k = \ln(0.6) = -0.5108\dots$  $\Rightarrow k = 0.17(02\dots)$	M1 A1  M1 M1  A1 [5]	$10\ 000 = Ae^0$ $A = 10\ 000$  taking lns (correctly) on their exponential equation - not logs unless to base 10 art 0.17 or $-(\ln 0.6)/3$ oe
(ii) $2000 = 10\ 000e^{-kt}$ $\Rightarrow -kt = \ln 0.2$  $\Rightarrow t = -\ln 0.2 / k = 9.45$ (years)	M1 A1 [2]	taking lns on correct equation (consistent with their $k$ ) allow art 9.5, but not 9.

<p><b>4</b> Perfect squares are 0, 1, 4, 9, 16, 25, 36, 49, 64, 81 none of which end in a 2, 3, 7 or 8.  Generalisation: no perfect squares end in a 2, 3, 7 or 8.</p>	M1 E1  B1 [3]	Listing all 1- and 2- digit squares. Condone absence of $0^2$ , and listing squares of 2 digit nos (i.e. $0^2 - 19^2$ )  For extending result to include further square numbers.
<b>5 (i)</b> $y = \frac{x^2}{2x+1}$ $\Rightarrow \frac{dy}{dx} = \frac{(2x+1)2x - x^2 \cdot 2}{(2x+1)^2}$ $= \frac{2x^2 + 2x}{(2x+1)^2} = \frac{2x(x+1)}{(2x+1)^2} *$	M1  A1  A1 E1 [4]	Use of quotient rule (or product rule)  Correct numerator – condone missing bracket provided it is treated as present Correct denominator www – do not condone missing brackets
<b>(ii)</b> $\frac{dy}{dx} = 0$ when $2x(x+1) = 0$ $\Rightarrow x = 0$ or $-1$ $y = 0$ or $-1$	B1 B1 B1 B1 [4]	Must be from correct working: SC –1 if denominator = 0
<b>6(i)</b> QA = $3 - y$ , PA = $6 - (3 - y) = 3 + y$ By Pythagoras, PA <sup>2</sup> = OP <sup>2</sup> + OA <sup>2</sup> $\Rightarrow (3+y)^2 = x^2 + 3^2 = x^2 + 9. *$	B1 B1  E1 [3]	must show some working to indicate Pythagoras (e.g. $x^2 + 3^2$ )
<b>(ii)</b> Differentiating implicitly: $2(y+3)\frac{dy}{dx} = 2x$ $\Rightarrow \frac{dy}{dx} = \frac{x}{y+3} *$	M1  E1	Allow errors in RHS derivative (but not LHS) - notation should be correct  brackets must be used
or $9 + 6y + y^2 = x^2 + 9$ $\Rightarrow 6y + y^2 = x^2$ $\Rightarrow (6+2y)\frac{dy}{dx} = 2x$ $\Rightarrow \frac{dy}{dx} = \frac{x}{y+3}$	M1  E1	Allow errors in RHS derivative (but not LHS) - notation should be correct brackets must be used
or $y = \sqrt{x^2 + 9} - 3 \Rightarrow \frac{dy}{dx} = \frac{1}{2}(x^2 + 9)^{-1/2} \cdot 2x$ $= \frac{x}{\sqrt{x^2 + 9}} = \frac{x}{y+3}$	M1  E1	(cao)
<b>(iii)</b> $\frac{dy}{dt} = \frac{dy}{dx} \cdot \frac{dx}{dt}$ $= \frac{4}{2+3} \times 2$ $= \frac{8}{5}$	M1  A1 A1 [3]	chain rule (soi)

## Section B

7(i) When $x = -1, y = -1\sqrt{0} = 0$ Domain $x \geq -1$	E1 B1 [2]	Not $y \geq -1$
(ii) $\frac{dy}{dx} = x \cdot \frac{1}{2}(1+x)^{-1/2} + (1+x)^{1/2}$ $= \frac{1}{2}(1+x)^{-1/2}[x + 2(1+x)]$ $= \frac{2+3x}{2\sqrt{1+x}} *$	B1 B1 M1 E1	$x \cdot \frac{1}{2}(1+x)^{-1/2}$ $\dots + (1+x)^{1/2}$ taking out common factor or common denominator www
$or u = x + 1 \Rightarrow du/dx = 1$ $\Rightarrow y = (u-1)u^{1/2} = u^{3/2} - u^{1/2}$ $\Rightarrow \frac{dy}{du} = \frac{3}{2}u^{\frac{1}{2}} - \frac{1}{2}u^{-\frac{1}{2}}$ $\Rightarrow \frac{dy}{dx} = \frac{dy}{du} \cdot \frac{du}{dx} = \frac{3}{2}(x+1)^{\frac{1}{2}} - \frac{1}{2}(x+1)^{-\frac{1}{2}}$ $= \frac{1}{2}(x+1)^{-\frac{1}{2}}(3x+3-1)$ $= \frac{2+3x}{2\sqrt{1+x}} *$	M1 A1 M1 E1 [4]	taking out common factor or common denominator
(iii) $dy/dx = 0$ when $3x + 2 = 0$ $\Rightarrow x = -2/3$ $\Rightarrow y = -\frac{2}{3}\sqrt{\frac{1}{3}}$ Range is $y \geq -\frac{2}{3}\sqrt{\frac{1}{3}}$	M1 A1ca o A1 B1 ft [4]	o.e. not $x \geq -\frac{2}{3}\sqrt{\frac{1}{3}}$ (ft their y value, even if approximate)
(iv) $\int_{-1}^0 x\sqrt{1+x} dx$ let $u = 1+x, du/dx = 1 \Rightarrow du = dx$ when $x = -1, u = 0, \text{ when } x = 0, u = 1$ $= \int_0^1 (u-1)\sqrt{u} du$ $= \int_0^1 (u^{3/2} - u^{1/2}) du *$	M1 B1 M1 E1	$du = dx$ or $du/dx = 1$ or $dx/du = 1$ changing limits – allow with no working shown provided limits are present and consistent with $dx$ and $du$ . $(u-1)\sqrt{u}$ www – condone only final brackets missing, otherwise notation must be correct
$= \left[ \frac{2}{5}u^{5/2} - \frac{2}{3}u^{3/2} \right]_0^1$ $= \pm \frac{4}{15}$	B1 B1 M1 A1ca o [8]	$\frac{2}{5}u^{5/2}, -\frac{2}{3}u^{3/2}$ (oe) substituting correct limits (can imply the zero limit) $\pm \frac{4}{15}$ or $\pm 0.27$ or better, not 0.26

<p><b>8 (i)</b> <math>f'(x) = 2(e^x - 1)e^x</math></p> <p>When <math>x = 0, f'(0) = 0</math> When <math>x = \ln 2, f'(\ln 2) = 2(2 - 1)2 = 4</math></p>	M1 A1  B1dep M1 A1cao [5]	or $f(x) = e^{2x} - 2e^x + 1$ M1 (or $(e^x)^2 - 2e^x + 1$ plus correct deriv of $(e^x)^2$ ) $\Rightarrow f'(x) = 2e^{2x} - 2e^x$ A1 derivative must be correct, www $e^{\ln 2} = 2$ soi
<p><b>(ii)</b> <math>y = (e^x - 1)^2 \quad x \leftrightarrow y</math></p> $\begin{aligned} &x = (e^y - 1)^2 \\ \Rightarrow &\sqrt{x} = e^y - 1 \\ \Rightarrow &1 + \sqrt{x} = e^y \\ \Rightarrow &y = \ln(1 + \sqrt{x}) \end{aligned}$	M1  M1 E1	reasonable attempt to invert formula  taking lns similar scheme of inverting $y = \ln(1 + \sqrt{x})$
<p>or <math>gf(x) = g((e^x - 1)^2)</math>  <math>= \ln(1 + e^x - 1)</math>  <math>= x</math></p>	M1 M1 E1	constructing gf or fg $\ln(e^x) = x$ or $e^{\ln(1+\sqrt{x})} = 1 + \sqrt{x}$
 Gradient at $(1, \ln 2) = \frac{1}{4}$	B1  B1ft [5]	reflection in $y = x$ (must have infinite gradient at origin)
<p><b>(iii)</b> <math>\int (e^x - 1)^2 dx = \int (e^{2x} - 2e^x + 1) dx</math></p> $\begin{aligned} &= \frac{1}{2}e^{2x} - 2e^x + x + c * \\ &\int_0^{\ln 2} (e^x - 1)^2 dx = \left[ \frac{1}{2}e^{2x} - 2e^x + x \right]_0^{\ln 2} \\ &= \frac{1}{2}e^{2\ln 2} - 2e^{\ln 2} + \ln 2 - (\frac{1}{2} - 2) \\ &= 2 - 4 + \ln 2 - \frac{1}{2} + 2 \\ &= \ln 2 - \frac{1}{2} \end{aligned}$	M1  E1  M1 M1 A1 [5]	expanding brackets (condone $e^{x^2}$ )  substituting limits $e^{\ln 2} = 2$ used must be exact
<p><b>(iv)</b></p>  $\text{Area} = 1 \times \ln 2 - (\ln 2 - \frac{1}{2}) = \frac{1}{2}$	M1 B1  A1cao [3]	subtracting area in (iii) from rectangle rectangle area = $1 \times \ln 2$  must be supported