

**GCE**

**Chemistry A**

Unit **F325**: Equilibria, Energetics and Elements

Advanced GCE

**Mark Scheme for June 2016**

OCR (Oxford Cambridge and RSA) is a leading UK awarding body, providing a wide range of qualifications to meet the needs of candidates of all ages and abilities. OCR qualifications include AS/A Levels, Diplomas, GCSEs, Cambridge Nationals, Cambridge Technicals, Functional Skills, Key Skills, Entry Level qualifications, NVQs and vocational qualifications in areas such as IT, business, languages, teaching/training, administration and secretarial skills.

It is also responsible for developing new specifications to meet national requirements and the needs of students and teachers. OCR is a not-for-profit organisation; any surplus made is invested back into the establishment to help towards the development of qualifications and support, which keep pace with the changing needs of today's society.

This mark scheme is published as an aid to teachers and students, to indicate the requirements of the examination. It shows the basis on which marks were awarded by examiners. It does not indicate the details of the discussions which took place at an examiners' meeting before marking commenced.













All examiners are instructed that alternative correct answers and unexpected approaches in candidates' scripts must be given marks that fairly reflect the relevant knowledge and skills demonstrated.

Mark schemes should be read in conjunction with the published question papers and the report on the examination.

OCR will not enter into any discussion or correspondence in connection with this mark scheme.

© OCR 2016

1. Annotations available in RM Assessor.

Annotation	Meaning
 BOD	Benefit of doubt given
 CON	Contradiction
 X	Incorrect response
 ECF	Error carried forward
 I	Ignore
 NAQ	Not answered question
 NBOD	Benefit of doubt not given
 POT	Power of 10 error
 ^	Omission mark
 RE	Rounding error
 SF	Error in number of significant figures
	Correct response

2. Abbreviations, annotations and conventions used in the detailed Mark Scheme (to include abbreviations and subject-specific conventions).

<b>Annotation</b>	<b>Meaning</b>
<b>DO NOT ALLOW</b>	Answers which are not worthy of credit
<b>IGNORE</b>	Statements which are irrelevant
<b>ALLOW</b>	Answers that can be accepted
( )	Words which are not essential to gain credit
—	Underlined words must be present in answer to score a mark
<b>ECF</b>	Error carried forward
<b>AW</b>	Alternative wording
<b>ORA</b>	Or reverse argument

3. The following questions should be marked using **ALL** appropriate annotations to show where marks have been awarded in the body of the text:
- 2(a)
  - 4(b)(ii)
  - 4(c)
  - 4(d)
  - 5(c)(i)
  - 5(c)(ii)
  - 5(d)(iv)
  - 6(c)
  - 8(e)

Question		Answer	Marks	Guidance	
1	(a)	<p><b>IGNORE</b> any charges shown within complexes (treat as rough working)</p> <p><b>Formulae</b> <span style="float: right;"><b>2 marks</b></span></p> <p style="text-align: center;"><math>[\text{Cu}(\text{NH}_3)_4(\text{H}_2\text{O})_2]^{2+}</math> ✓</p> <p style="text-align: center;"><math>[\text{CuCl}_4]^{2-}</math> ✓</p> <p><b>Colours</b> <span style="float: right;"><b>1 mark</b></span></p> <p>blue <b>AND</b> yellow ✓</p> <p><i>Mark independently of formulae</i></p>	3	<p>For charges, <b>ALLOW</b> +2 and –2</p> <p>Square brackets <b>required</b>, i.e. <b>DO NOT ALLOW</b> <math>\text{Cu}(\text{NH}_3)_4(\text{H}_2\text{O})_2^{2+}</math></p> <p><b>ALLOW</b> Ligands in any order</p> <p><b>ALLOW</b> <math>\text{CuCl}_4^{2-}</math> i.e. no brackets <b>OR</b> <math>\text{Cu}(\text{Cl})_4^{2-}</math></p> <p>For <math>\text{CuCl}_4^{2-}</math>, <b>ALLOW</b> green–yellow <b>OR</b> yellow–green <b>DO NOT ALLOW</b> green</p> <p>For <math>[\text{Cu}(\text{NH}_3)_4(\text{H}_2\text{O})_2]^{2+}</math> <b>DO NOT ALLOW</b> pale blue, light blue</p> <p><b>DO NOT ALLOW</b> precipitate with blue <b>OR</b> yellow</p>	
1	(b)	(i)	<p><b>Donates two</b> electron <b>pairs</b> to a metal ion/metal/<math>\text{Cu}^{2+}</math> <b>AND</b> forms <b>two</b> coordinate <b>bonds</b> to a metal ion/metal/<math>\text{Cu}^{2+}</math> ✓</p>	1	<p><b>ALLOW</b> lone pairs for electron pairs <b>ALLOW</b> molecule/atom/ion/substance for 'ligand' <b>ALLOW</b> dative (covalent) bonds for coordinate bonds <b>ALLOW</b> transition element for metal</p> <p><b>Two is needed once only</b> e.g. <b>Donates two</b> electron <b>pairs</b> to form coordinate <b>bonds</b> to a metal ion/metal/<math>\text{Cu}^{2+}</math> <b>Donates</b> electron <b>pairs</b> to form <b>two</b> coordinate <b>bonds</b> to a metal ion/metal/<math>\text{Cu}^{2+}</math></p> <p><b>DO NOT ALLOW</b> donates <b>two</b> electron pairs to form <b>one/a</b> coordinate bond</p>

Question			Answer	Marks	Guidance						
1	(b)	(ii)		3	<p><b>FULL ANNOTATIONS MUST BE USED</b></p> <p>-----</p> <p><b>2 marks:</b> one for each correct isomer ✓✓</p> <p><b>TAKE CARE:</b> structures may be in different orientations and in different order</p> <p><b>IF BOTH</b> isomers are 'correct', but O connectivity wrong, <b>AWARD</b> 1 mark for both structures  <b>Check</b> H<sub>2</sub>O ligands carefully for connectivity</p> <p><b>ALLOW</b> H<sub>2</sub>O reversed shown as -O<sub>2</sub>H</p> <p><b>IGNORE</b> charges (anywhere)</p> <p>-----</p> <p><b>NOTE:</b> For each structure, <b>ALL</b> O atoms must be shown <b>AND</b> For (COO<sup>-</sup>)<sub>2</sub>, <b>ALLOW</b> skeletal, structural or displayed formula</p> <p><b>DO NOT ALLOW</b> structures such as those shown below</p> <p>-----</p> <p><b>1 mark:</b> for whole of 2nd row for whole of 'Type' row  i.e. (cis <b>AND</b> optical) <b>AND</b> trans only</p>						
			<table border="1"> <tbody> <tr> <td><i>cis</i></td> <td>✓</td> </tr> <tr> <td><i>trans</i></td> <td></td> </tr> <tr> <td>optical</td> <td>✓</td> </tr> </tbody> </table>	<i>cis</i>	✓	<i>trans</i>		optical	✓		
<i>cis</i>	✓										
<i>trans</i>											
optical	✓										
			<table border="1"> <tbody> <tr> <td><i>cis</i></td> <td></td> </tr> <tr> <td><i>trans</i></td> <td>✓</td> </tr> <tr> <td>optical</td> <td></td> </tr> </tbody> </table>	<i>cis</i>		<i>trans</i>	✓	optical			
<i>cis</i>											
<i>trans</i>	✓										
optical											



Question			Answer	Marks	Guidance
1	(b)	(iii)	$\text{CuC}_4\text{H}_4\text{O}_{10}^{2-}$ Formula ✓ 2- charge ✓ <b>MARK formula and charge INDEPENDENTLY</b>	2	<b>Empirical</b> formula essential, e.g. <b>DO NOT ALLOW</b> $\text{Cu}(\text{COO})_2(\text{H}_2\text{O})_2$ for formula mark <b>ALLOW</b> any order of elements in formula <b>ALLOW</b> -2 for charge
			Total	9	

Question	Answer	Marks	Guidance
2 (a)	<p><b>initial rates data (3 marks)</b>  <b>NOTE:</b> Each comparison <b>MUST</b> relate to the <b>actual</b> change in concentration/rate in the experiments</p> <p style="text-align: right;"><b>EXPTS</b></p> <p><math>\text{H}_2\text{O}_2</math>: <math>[\text{H}_2\text{O}_2] \times 2</math>      rate <math>\times 2</math>      (1 &amp; 2)  <b>AND</b>  1st order ✓</p> <p><math>\text{H}^+</math>: <math>[\text{H}^+] \times 2</math>      rate does not change      (2 &amp; 3)  <b>AND</b>  Zero order ✓</p> <p><math>\text{I}^-</math>: <math>[\text{I}^-] \times 2</math> <b>AND</b> <math>[\text{H}_2\text{O}_2] \times 2</math>      rate <math>\times 4</math>      (2 &amp; 4)  <b>OR</b> <math>[\text{I}^-] \times 2</math> <b>AND</b> <math>[\text{H}_2\text{O}_2] \times 4</math>      rate <math>\times 8</math>      (1 &amp; 4)  <b>OR</b> <math>[\text{I}^-] \times 2</math> <b>AND</b> <math>[\text{H}_2\text{O}_2] \times 2</math>      rate <math>\times 4</math>      (3 &amp; 4)  <b>AND</b>  1st order ✓</p>	3	<p><b>FULL ANNOTATIONS MUST BE USED</b>  -----</p> <p><b>THROUGHOUT,</b></p> <ul style="list-style-type: none"> <li>• Square brackets <b>NOT REQUIRED</b> around <math>\text{H}_2\text{O}_2</math>, <math>\text{H}^+</math> and <math>\text{I}^-</math></li> <li>• <b>ALLOW</b> 'doubles' for <math>\times 2</math>; quadruples for <math>\times 4</math></li> </ul> <p><b>ALLOW</b> direct comparison of concentrations and rate, e.g.  <math>[\text{H}_2\text{O}_2]</math> changes by <math>\frac{0.0020}{0.0010} = 2</math>, rate changes by <math>\frac{1.14 \times 10^{-5}}{5.70 \times 10^{-6}} = 2</math>  <b>AND</b> 1st order (Expts 1 &amp; 2)</p> <p><b>DO NOT ALLOW</b> <math>\text{I}_2</math> for <math>\text{I}^-</math></p> <p><b>IGNORE</b> <math>[\text{H}^+]</math> for Expts 3 &amp; 4</p>
	<p><b>Calculation of rate constant (3 marks),</b>  <b>EITHER</b>  <math>k = \frac{5.70 \times 10^{-6}}{0.0010 \times 0.20}</math> <b>OR</b> <math>2.85 \times 10^{-2}</math> <b>OR</b> <math>0.0285</math> <b>OR</b> <math>0.029</math> ✓</p> <p><math>k = 2.9 \times 10^{-2}</math> ✓ (2 SF in standard form)  <i>Subsumes previous mark if no working shown</i></p> <p style="text-align: center;"><math>\text{dm}^3 \text{mol}^{-1} \text{s}^{-1}</math> ✓</p>	3	<p><b>IGNORE</b> working</p> <p><b>DO NOT ALLOW</b> 0.03</p> <p><b>ALLOW ECF</b> from error in powers of 10 <b>ONLY</b>  e.g. <math>2.9 \times 10^{-3}</math> by use of 0.010 instead of 0.0010  <b>DO NOT ALLOW</b> <math>2.90 \times 10^{-2}</math> (3 SF)  <b>OR</b> <math>29 \times 10^{-3}</math> (Not standard form)</p> <p><b>ALLOW</b> <math>\text{mol}^{-1}</math>, <math>\text{dm}^3</math> and <math>\text{s}^{-1}</math> in any order, e.g. <math>\text{mol}^{-1} \text{dm}^3 \text{s}^{-1}</math></p>





Question		Answer	Marks	Guidance
3	(a)	(enthalpy change for) 1 mole of gaseous ions <b>OR</b> 1 mole of hydrated ions/aqueous ions ✓  gaseous ions forming aqueous/hydrated ions ✓	2	<p><b>one mole</b> can be stated just once <b>EITHER</b> with gaseous ions <b>OR</b> with aqueous ions, e.g.</p> <ul style="list-style-type: none"> <li>• 1 mole of gaseous ions <b>forms</b> hydrated ions/aqueous ions</li> <li>• Gaseous ions <b>form 1 mole of</b> hydrated ions/aqueous ions</li> </ul> <p><b>ALLOW</b> 1 mol for 1 mole</p> <p><b>IGNORE</b> 'energy released' <b>OR</b> 'energy required'</p> <p>For 2nd mark  <b>IGNORE</b> gaseous ions are hydrated  <b>IGNORE</b> gaseous ions dissolve in water  <i>Particles formed not stated</i></p> <p><b>ALLOW</b> 1 mark for:  1 mole of gaseous <b><u>IONS</u></b> forms aqueous/hydrated <b>atoms/ particles/ molecules</b></p>

Question	Answer	Marks	Guidance
3 (b) (i)	<p><b>4 marks</b> for species <b>AND</b> state symbols on all 4 energy levels (including added energy level)</p>  <p><b>1 mark</b> for <b>B, C AND D</b> labels <b>OR</b> enthalpy values <b>AND</b> arrow directions correct ✓</p> <p><b>ALLOW</b> <math>K_2SO_4(aq)</math> for <math>2K^+(aq) + SO_4^{2-}(aq)</math></p> <p><b>ALLOW</b> arrows not touching lines.</p> <p><b>Direction</b> is important:</p> <ul style="list-style-type: none"> <li>• <b>FROM</b> <math>2K^+(g) + SO_4^{2-}(g)</math> line</li> <li>• <b>FROM</b> <math>K_2SO_4(s)</math> line</li> </ul> <p><b>See APPENDIX</b></p> <p>'2 x' is <b>NOT</b> required – <i>part of calculation mark</i></p>	5	<p><b>IF</b> extra energy level is <b>above</b> top line <b>OR below</b> bottom line, <b>DO NOT ALLOW</b> mark for species on this line. <b>See APPENDIX</b></p> <p><b>ALLOW C</b> and <b>D</b> with associated labels, the other way round:</p>  <p>State symbols are <b>essential</b></p> <p><b>IF</b> no extra energy level is shown with <b>C</b> and <b>D</b> combined forming <math>2K^+(aq) + SO_4^{2-}(aq)</math>,</p> <ul style="list-style-type: none"> <li>• No mark for the extra energy level with species</li> <li>• No mark for labels as <b>C</b> and <b>D</b> are combined</li> </ul> <p>Therefore 3 max for species on energy levels provided</p>
3 (b) (ii)	$\Delta H(\text{hydration}) SO_4^{2-} = -1099 \text{ (kJ mol}^{-1}\text{)} \checkmark$	1	<b>ONLY</b> correct answer

Question			Answer	Marks	Guidance
3	(c)	(i)	<p><b>Aqueous</b> particles are more disordered than <b>solid</b> (particles)  <b>OR</b>  <b>Solid</b> particles are more ordered than <b>aqueous</b> (particles) ✓</p>	1	<p>For particles, <b>ALLOW</b> ions  <b>DO NOT ALLOW</b> molecules/atoms</p> <p><b>ALLOW</b> 'When the state changes from solid to aqueous, disorder increases'</p> <p>For more disordered, <b>ALLOW</b> less ordered/ more freedom/ more ways of arranging energy/ more random</p> <p>For aqueous particles, <b>ALLOW particles in solution</b></p> <p><b>IGNORE</b> dissolved</p>
3	(c)	(ii)	<p><b>Calculation (2 marks)</b>  <math>\Delta G = 24 - (298 \times 0.225)</math> <b>OR</b> <math>24 - 67.05</math> (in kJ)  <b>OR</b> <math>24000 - (298 \times 225)</math> <b>OR</b> <math>24000 - 67050</math> (in J) ✓</p> <p><b>Calculation of <math>\Delta G</math> (IGNORE UNITS)</b>  <math>\Delta G = -43</math> (kJ mol<sup>-1</sup>) <b>OR</b> <math>-43000</math> (J mol<sup>-1</sup>) ✓  <i>Subsumes 1st calculation mark</i></p> <p><b>Reason for solubility</b>            Calculated value of <math>\Delta G</math> that is <b>negative</b>  <b>AND</b>            Statement that:  <math>\Delta G</math> is negative <b>OR</b> <math>\Delta G &lt; 0</math> <b>OR</b> <math>-43 &lt; 0</math>  <b>OR</b> <math>\Delta H - T\Delta S &lt; 0</math> <b>OR</b> <math>T\Delta S &gt; \Delta H</math> ✓</p>	3	<p><b>Contact TL if solely entropy approach rather than <math>\Delta G</math></b></p> <p><b>ALLOW</b> <math>-43.1</math> <b>OR</b> <math>-43.05</math> (<i>calculator value</i>)</p> <p><b>ALLOW 1 calculation mark (IGNORE units) for</b>  <math>-67.(026)</math> <b>OR</b> <math>-67026</math> <b>ECF</b> from 225 instead of 0.225  <math>18.(375)</math> <b>OR</b> <math>+18.375</math> <b>ECF</b> from 25 instead of 298</p> <p><b>ALLOW other ECF</b> from <b>ONE</b> error in 1st step of calc, e.g. incorrect value for <math>\Delta H</math> such as <math>-1099</math> from 3bii <math>\rightarrow -1166.05</math>  <b>TAKE CARE</b> that same units used for <math>\Delta H</math> and <math>\Delta S</math></p> <p><b>NO reason mark from a +ve value of <math>\Delta G</math></b></p>
<b>Total</b>				<b>12</b>	

Question	Answer	Marks	Guidance
4 (a)	Iodine is non-polar <b>OR</b> Iodine does not form H bonds <b>with water</b> ✓	1	<b>IGNORE</b> iodine is slightly polar <b>IGNORE</b> 'cannot bond to water' (too vague) <b>IGNORE</b> 'Lack of a lone pair' <b>IGNORE</b> 'inability to induce a dipole'
4 (b)	<p><b>FIRST, CHECK THE ANSWER ON ANSWER LINE</b>  <b>IF <math>K_c = 104 \text{ dm}^3 \text{ mol}^{-1}</math> award 4 marks:</b>  <b>3 for calculation of 104 from data, 1 for units</b></p> <p>-----</p> <p><b>Equilibrium concentrations (mol × 5) (1 mark)</b>  <math>I_2 = 4.00 \times 10^{-5} \times 5 = 2.00 \times 10^{-4} \text{ (mol dm}^{-3}\text{)}</math>  <b>AND <math>I^- = 9.404 \times 10^{-2} \times 5 = 0.4702 \text{ (mol dm}^{-3}\text{)}</math> ✓</b>  <b>AND <math>I_3^- = 1.96 \times 10^{-3} \times 5 = 9.80 \times 10^{-3} \text{ (mol dm}^{-3}\text{)}</math></b></p> <p><b>Calculation of <math>K_c</math> and units (3 marks)</b></p> $K_c = \frac{[I_3^-(aq)]}{[I_2(aq)] \times [I^-(aq)]} \text{ OR } \frac{9.80 \times 10^{-3}}{2.00 \times 10^{-4} \times 0.4702} \checkmark$ <p>= 104 ✓ <b>Must be 3 SF</b></p> <p><math>\text{dm}^3 \text{ mol}^{-1}</math> <b>OR</b> <math>\text{mol}^{-1} \text{ dm}^3</math> ✓</p>	4	<p><b>FULL ANNOTATIONS MUST BE USED</b></p> <p>-----</p> <p>Throughout, at least <b>3SF</b> but <b>ALLOW</b> absence of trailing zeroes  e.g. for <math>9.80 \times 10^{-3}</math> <b>ALLOW</b> <math>9.8 \times 10^{-3}</math>  <b>FOR <math>I^-</math> 0.4702, ALLOW 0.47(0) (mol dm<sup>-3</sup>)</b> still →104 for calc</p> <p>State symbols <b>not</b> required in <math>K_c</math> expression  <b>ALLOW ECF</b> from incorrect concentrations</p> <p>Any <b>ECF</b> value <b>MUST</b> be to <b>3 SF</b> for <math>K_c</math> value</p> <p>-----</p> <p><b>COMMON ERRORS</b></p> <p><b>104.2 → 104.2109741 (calc) &gt; 3 SF 2 marks + units</b></p> <p><b>521 no × 5 for concs 2 marks + units</b></p> <p><b>521.1 → 521.0548703 as above and &gt; 3SF 1 mark + units</b></p> <p><b>2610 ÷ 5 instead of × 5 for concs 2 marks + units</b></p> <p><b><math>9.60 \times 10^{-3}</math> <math>K_c</math> upside down, correct concs 2 marks + units</b></p> <p><b><math>1.92 \times 10^{-3}</math> <math>K_c</math> upside down, no × 5 for concs 1 mark + units</b></p> <p><b>NOTE:</b> With <math>K_c</math> upside down, units become <math>\text{mol dm}^{-3}</math> by ECF</p>

Question		Answer	Marks	Guidance
4	(c)	<p>Ag<sup>+</sup>/silver nitrate reacts with I<sup>-</sup> to form AgI/silver iodide  <b>OR</b> Ag<sup>+</sup> + I<sup>-</sup> → AgI ✓</p> <p>yellow precipitate/solid forms ✓</p> <p><b>Equilibrium 2</b> shifts to the left ✓</p> <p><b>Equilibrium 1</b> shifts to left  <b>AND</b>  I<sub>2</sub> comes out of solution/less I<sub>2</sub> dissolves/  I<sub>2</sub> precipitates/black solid /grey solid /violet solid ✓</p>	4	<p><b>FULL ANNOTATIONS MUST BE USED</b></p> <p>-----</p> <p><b>DO NOT ALLOW</b> cream <b>OR</b> cream–yellow  <b>ALLOW</b> just ‘yellow’ if supported by AgI(s) somewhere</p>
4	(d)	<p><b>in all equations ALLOW equilibrium signs</b>  <b>IGNORE</b> state symbols</p> <p>-----</p> <p><b>Reaction 1: 1 mark</b>  2I<sub>2</sub> + 5O<sub>2</sub> → 2I<sub>2</sub>O<sub>5</sub> ✓</p> <p>-----</p> <p><b>Reaction 2: 2 marks</b>  <b>1st mark: ALL CORRECT species</b></p> <p>e.g.: I<sub>2</sub> + OH<sup>-</sup> → I<sup>-</sup> + IO<sub>3</sub><sup>-</sup> + H<sub>2</sub>O</p> <p><b>2nd mark for CORRECT balanced equation</b>  3I<sub>2</sub> + 6OH<sup>-</sup> → 5I<sup>-</sup> + IO<sub>3</sub><sup>-</sup> + 3H<sub>2</sub>O  ✓✓</p>	3	<p><b>FULL ANNOTATIONS MUST BE USED</b></p> <p>-----</p> <p><b>ALLOW</b> correct multiples throughout, e.g. I<sub>2</sub> + 2½O<sub>2</sub> → I<sub>2</sub>O<sub>5</sub></p> <p>-----</p> <p><b>For 1st mark, IGNORE</b> e<sup>-</sup> present</p> <p><b>ALLOW</b> species/equation with NaOH or KOH,  e.g. 3I<sub>2</sub> + 6NaOH → 5I<sup>-</sup> + IO<sub>3</sub><sup>-</sup> + 3H<sub>2</sub>O + 6Na<sup>+</sup>  3I<sub>2</sub> + 6NaOH → 5NaI + NaIO<sub>3</sub> + 3H<sub>2</sub>O</p> <p><b>ALLOW</b>  Species: I<sub>2</sub> + OH<sup>-</sup> → I<sup>-</sup> + IO<sub>2</sub><sup>+</sup> + H<sub>2</sub>O ✓  <b>OR</b> Equation: 3I<sub>2</sub> + 4OH<sup>-</sup> → 5I<sup>-</sup> + IO<sub>2</sub><sup>+</sup> + 2H<sub>2</sub>O ✓✓</p> <p>Species: I<sub>2</sub> + OH<sup>-</sup> → I<sup>-</sup> + IO<sub>3</sub><sup>+</sup> + H<sub>2</sub>O ✓  <b>OR</b> Equation: 3I<sub>2</sub> + 2OH<sup>-</sup> → 5I<sup>-</sup> + IO<sub>3</sub><sup>+</sup> + H<sub>2</sub>O ✓✓</p>
		<b>Total</b>	<b>12</b>	

Question		Answer	Marks	Guidance
5	(a)	$(K_a =) \frac{[H^+][NO_2^-]}{[HNO_2]} \checkmark$ <p><b>IGNORE</b> state symbols</p>	1	<p><b>IGNORE</b> <math>\frac{[H^+]^2}{[HNO_2]}</math> <b>OR</b> <math>\frac{[H^+][A^-]}{[A]}</math></p> <p><b>ALLOW</b> <math>H_3O^+</math> for <math>H^+</math></p> <p>Square brackets <b>required</b></p>
5	(b)	<p><b>FIRST, CHECK THE ANSWER ON ANSWER LINE</b>  <b>IF answer = 2.12 award 2 marks</b></p> <p>-----</p> $[H^+] = \sqrt{K_a [HNO_2]} = 7.502 \times 10^{-3} \text{ (mol dm}^{-3}\text{)} \checkmark$ $\text{pH} = -\log 7.502 \times 10^{-3} = 2.12 \checkmark \quad \text{pH to 2 DP}$	2	<p>-----</p> <p><b>ALLOW</b> intermediate value from 3 SF (7.50 up to calculator value of <math>7.501999733 \times 10^{-3}</math>)</p> <p><b>ALLOW</b> 1 mark for 2.1 <b>OR</b> answer &gt; 2 DP (i.e. not 2 DP)</p> <p><b>ONLY ALLOW</b> pH mark by <b>ECF</b> if <math>K_a</math> <b>AND</b> 0.120 used and <b>AND</b> pH &lt; 7</p> <p>-----</p> <p><b>COMMON ERRORS (MUST be to 2 DP)</b></p> <p><b>pH = 4.25 No square root: 1 mark</b>  <math>[H^+] = (4.69 \times 10^{-4} \times 0.120) = 5.628 \times 10^{-5} \text{ (mol dm}^{-3}\text{)}</math>  <math>\text{pH} = -\log 5.628 \times 10^{-5} = 4.25 \checkmark</math></p> <p><b>pH = 0.92 no <math>K_a</math> used: zero marks</b>  <math>\text{pH} = -\log 0.120 = 0.92</math></p> <p><b>pH = 13.08 <math>K_w</math>/pOH used: zero marks</b>  <math>\text{pH} = -\log \frac{1.00 \times 10^{-14}}{0.120}</math> <b>OR</b> <math>14 - \log 0.120 = 13.08</math></p>


Question	Answer	Marks	Guidance
5 (c) (i)	<p><b>FIRST, CHECK THE ANSWER ON ANSWER LINE</b>  <b>IF answer = 3.43, AWARD 4 marks</b></p> <p>-----</p> <p>Expression: <math>K_a \times \text{acid/base ratio}</math></p> <p>Use of <math>K_a \times \frac{[\text{HNO}_2]}{[\text{NO}_2^-]}</math> <b>OR</b> <math>4.69 \times 10^{-4} \times \frac{[\text{HNO}_2]}{[\text{NO}_2^-]}</math> ✓</p> <p>Using correct concs/mol in expression</p> <p><math>[\text{H}^+] = 4.69 \times 10^{-4} \times \frac{0.0400}{0.0500}</math> ✓ Subsumes previous mark</p> <p>Calculation of <math>[\text{H}^+]</math></p> <p><math>[\text{H}^+] = 3.752 \times 10^{-4} \text{ (mol dm}^{-3}\text{)}</math> ✓</p> <p>pH to 2 DP (From 3.42573717)</p> <p><math>\text{pH} = -\log 3.752 \times 10^{-4} = 3.43</math> ✓</p> <p><b>NO marks</b> are available using</p> <p><math>K_a</math> square root approach (weak acid pH)</p> <p><math>K_w / 10^{-14}</math> approach (strong base pH)</p> <p>-----</p> <p><b>ALLOW</b> alternative approach based on Henderson–Hasselbalch equation (<b>ALLOW</b> <math>-\log K_a</math> for <math>\text{p}K_a</math>)</p> <p><math>\text{pH} = \text{p}K_a + \log \frac{[\text{NO}_2^-]}{[\text{HNO}_2]}</math> <b>OR</b> <math>\text{p}K_a - \log \frac{[\text{HNO}_2]}{[\text{NO}_2^-]}</math> ✓</p> <p><math>\text{pH} = \text{p}K_a + \log \frac{0.0500}{0.0400}</math> <b>OR</b> <math>\text{p}K_a - \log \frac{0.0400}{0.0500}</math> ✓</p> <p><math>\text{pH} = \text{p}K_a + 0.097</math> ✓</p> <p><math>\text{pH} = 3.329 + 0.097 = 3.43</math> ✓</p>	4	<p><b>FULL ANNOTATIONS MUST BE USED</b></p> <p>-----</p> <p><b>ALLOW</b> just <math>K_a \times \frac{\text{acid}}{\text{salt}}</math> expression</p> <p><b>Mark by ECF</b> from <math>4.69 \times 10^{-4} \times \frac{[\text{NO}_2^-]}{[\text{HNO}_2]}</math> inverted expression</p> <p><b>Mark by ECF</b> from incorrect <math>[\text{HNO}_2]</math> and <math>[\text{NO}_2^-]</math>  <b>ONLY</b> award <b>marks</b> for a pH calculation via <math>K_a</math> <b>AND</b> using concentrations/mol derived from the question</p> <p><b>DO NOT ALLOW</b> final pH mark by <b>ECF</b> if <math>\text{pH} &gt; 7</math></p> <p>-----</p> <p><b>COMMON ERRORS BUT CHECK WORKING</b></p> <p><b>pH = 2.82      3 marks</b>  initial concs: 0.200 and 0.0625</p> <p><b>pH = 3.23      3 marks</b>  0.0400 and 0.0500 acid/base ratio inverted</p> <p><b>pH = 3.83      2 marks</b>  initial concs: 0.200 and 0.0625 and ratio inverted</p> <p><b>pH = 2.73      3 marks</b>  Incorrect <math>[\text{NO}_2^-] = 0.01</math> and correct <math>[\text{HNO}_2] = 0.04</math></p> <p><b>pH = 4.03      3 marks</b>  correct <math>[\text{NO}_2^-] = 0.05</math> and incorrect <math>[\text{HNO}_2] = 0.01</math></p>



Question			Answer	Marks	Guidance
5	(c)	(ii)	<p><b>Equilibrium: 1 mark</b>  <math>\text{HNO}_2 \rightleftharpoons \text{H}^+ + \text{NO}_2^-</math> ✓            (ignore state symbols)</p> <p><b>Control of pH: 2 marks (QWC)</b>  <b>Added HCl</b>  <math>\text{NO}_2^-</math> reacts with added acid/HCl/<math>\text{H}^+</math>  <b>OR</b> <math>\text{NO}_2^- + \text{H}^+ \rightarrow</math>  <b>OR</b> more <math>\text{HNO}_2</math> forms ✓</p> <p><b>Added NaOH</b>  <math>\text{HNO}_2</math> reacts with added alkali/NaOH/<math>\text{OH}^-</math>  <b>OR</b> <math>\text{HNO}_2 + \text{OH}^- \rightarrow</math>  <b>OR</b> more <math>\text{NO}_2^-</math> forms  <b>OR</b> <math>\text{H}^+</math> reacts with added alkali/NaOH  <b>OR</b> <math>\text{H}^+ + \text{OH}^- \rightarrow</math> ✓</p> <p><b>Equilibrium shift:</b>  <b>1 mark for shifts in</b> <math>\text{HNO}_2 \rightleftharpoons \text{H}^+ + \text{NO}_2^-</math> (See 1st mark)            Equilibrium for added <b>acid</b> → <b>left</b>  <b>AND</b> Equilibrium for added <b>alkali</b> → <b>right</b> ✓ (QWC)</p>	4	<p><b>FULL ANNOTATIONS MUST BE USED</b>            -----</p> <p><b>IGNORE</b> <math>\text{HA} \rightleftharpoons \text{H}^+ + \text{A}^-</math></p> <p>Equilibrium sign <b>essential</b>  <b>BUT ALLOW</b> small slips in its appearance if it is obviously an attempt to show an equilibrium sign rather than an arrow</p> <p><b>QWC: Quality of written communication</b></p> <p><b>DO NOT ALLOW</b> HA and <math>\text{A}^-</math> for <math>\text{HNO}_2</math> and <math>\text{NO}_2^-</math></p> <p><b>IGNORE just</b> acid reacts with added alkali</p> <p><b>IGNORE just</b> conjugate base/salt/base reacts with added acid  <b>DO NOT ALLOW</b> salt/base reacts with added acid</p> <p><b>AWARD</b> 'shift mark' <b>ONLY</b> if correct equilibrium equation has been given  <b>IGNORE</b> any other equilibria in response</p>

Question			Answer	Marks	Guidance
5	(d)	(i)	Endothermic <b>AND</b> $K_w$ increases with temperature <b>OR</b> Endothermic <b>AND</b> dissociation increases <b>with temperature</b> <b>OR</b> Endothermic <b>AND</b> (dissociation) involves breaking bonds ✓	1	Endothermic <b>and</b> reason required for the mark  <b>ALLOW</b> Endothermic <b>AND</b> increasing temperature shifts equilibrium/reaction to the right/favours forward reaction  <b>DO NOT ALLOW</b> breaking hydrogen bonds <b>OR</b> intermolecular bonds/forces
5	(d)	(ii)	<b><i>OH<sup>-</sup> concentration</i></b> $[\text{OH}^-] = \frac{9.311 \times 10^{-14}}{1.00 \times 10^{-7}} = 9.311 \times 10^{-7} \text{ (mol dm}^{-3}\text{)} \checkmark$ <b><i>Explanation (dependent on 1st mark)</i></b> $9.311 \times 10^{-7} > 1.00 \times 10^{-7}$ <b>OR</b> $[\text{OH}^-] > [\text{H}^+]$ <b>OR</b> $\text{OH}^-$ in excess <b>AND</b> Alkaline ✓	2	<b><i>H<sup>+</sup> OR OH<sup>-</sup> concentration (neutral pH)</i></b> $[\text{H}^+] = [\text{OH}^-] = \sqrt{(9.311 \times 10^{-14})} = 3.05 \times 10^{-7} \text{ (mol dm}^{-3}\text{)} \checkmark$  <b><i>Explanation (dependent on 1st mark)</i></b> $\text{pH} = -\log(3.05 \times 10^{-7}) = 6.5 \rightarrow 6.515501837$ (calc) <b>AND</b> Alkaline ✓
5	(d)	(iii)	$\text{p}K_w = 13.03 \checkmark$	1	<b>ONLY</b> correct answer

Question			Answer	Marks	Guidance
5	(d)	(iv)	<p><b>FIRST, CHECK THE ANSWER ON ANSWER LINE</b>  <b>IF answer = 10.76, award 3 marks</b></p> <p>-----</p> <p><b>Dilution 1 mark</b>  <math>[\text{OH}^-(\text{aq})] = [\text{NaOH}(\text{aq})] = \frac{0.0270}{5} = 0.00540 \text{ (mol dm}^{-3}\text{)} \checkmark</math></p> <p><b>[H<sup>+</sup>] 1 mark</b>  <math>[\text{H}^+(\text{aq})] = \frac{9.311 \times 10^{-14}}{0.00540} = 1.72 \times 10^{-11} \text{ (mol dm}^{-3}\text{)} \checkmark</math>  Calculator: <math>1.724259259 \times 10^{-11}</math></p> <p><b>pH 1 mark</b>  <math>\text{pH} = -\log 1.72 \times 10^{-11} = \mathbf{10.76} \checkmark</math></p> <p>-----</p> <p><b>ALLOW</b> pOH method for 2nd and 3rd mark:</p> <p><math>\text{pOH} = -\log 0.00540 = 2.27 \checkmark</math> (calculator 2.26760624)  <math>\text{pH} = 13.03 - 2.27 = 10.76 \checkmark</math></p>	3	<p><b>FULL ANNOTATIONS MUST BE USED</b></p> <p>-----</p> <p><b>ALLOW</b> dilution <b>AFTER</b> calculation of <math>[\text{H}^+(\text{aq})]</math>  i.e. original <math>[\text{H}^+] = \frac{9.311 \times 10^{-14}}{0.0270} = 3.45 \times 10^{-12} \text{ (mol dm}^{-3}\text{)} \checkmark</math>  After dilution, <math>[\text{H}^+] = 3.45 \times 10^{-12} \times 5 = 1.72 \times 10^{-11} \text{ (mol dm}^{-3}\text{)} \checkmark</math>  <math>\text{pH} = -\log 1.72 \times 10^{-11} = \mathbf{10.76} \checkmark</math>  <b>ALLOW ECF</b> from incorrect <math>[\text{H}^+(\text{aq})]</math> provided that <math>\text{pH} &gt; 7</math></p> <p>-----</p> <p><b>COMMON ERRORS (MUST be to 2 DP)</b></p> <p><b>pH = 11.73</b> At 25°C (<math>1.00 \times 10^{-14}</math>): <b>2 marks</b>  <math>\text{pH} = -\log 1.85 \times 10^{-12} = \mathbf{11.73}</math></p> <p><b>pH = 11.46</b> No dilution at 60°C (<math>9.311 \times 10^{-14}</math>) <b>2 marks</b>  <math>\text{pH} = -\log(3.45 \times 10^{-12}) = \mathbf{11.46}</math></p> <p><b>pH = 12.43</b> No dilution <b>AND</b> 25°C (<math>1.00 \times 10^{-14}</math>) <b>1 mark</b>  <math>\text{pH} = -\log(3.70 \times 10^{-13}) = \mathbf{12.43}</math></p> <p><b>pH = 12.16</b> <math>\times 5</math> instead of <math>\div 5</math> at 60°C (<math>9.311 \times 10^{-14}</math>) <b>2 marks</b>  <math>\text{pH} = -\log(6.879 \times 10^{-13}) = \mathbf{12.16}</math></p> <p><b>pH = 13.13</b> <math>\times 5</math> instead of <math>\div 5</math> at 25°C (<math>1.00 \times 10^{-14}</math>) <b>1 mark</b>  <math>\text{pH} = -\log(7.407 \times 10^{-14}) = \mathbf{13.13}</math></p> <p><b>NOTE:</b> Attempts at dilution <math>\rightarrow 0.0270</math> with error in powers of 10  <math>\rightarrow 12.46</math> from <math>0.00270</math>, etc may give 2 marks by <b>ECF</b></p>
			<b>Total</b>	<b>18</b>	

Question		Answer	Marks	Guidance
6	(a)	<p><b>Definition</b> The e.m.f. (of a half-cell) compared with/connected to a (standard) hydrogen half-cell/(standard) hydrogen electrode ✓</p> <p><b>Standard conditions</b> <i>Units essential</i> Temperature of 298 K / 25°C <b>AND</b> (solution) concentrations of 1 mol dm<sup>-3</sup> <b>AND</b> pressure of 100 kPa <b>OR</b> 10<sup>5</sup> Pa <b>OR</b> 1 bar ✓</p>	2	<p>For e.m.f., <b>ALLOW</b> voltage <b>OR</b> potential difference/p.d. <b>OR</b> electrode/reduction/redox potential <b>ALLOW</b> e.m.f. of a cell .....</p> <p><b>ALLOW</b> /(standard) hydrogen cell <b>IGNORE</b> S.H.E. (as abbreviation for standard hydrogen electrode) <b>DO NOT ALLOW</b> hydrogen fuel cell</p> <p><b>ALLOW</b> 1M <b>OR</b> 1 mol/dm<sup>3</sup> <b>DO NOT ALLOW</b> 1 mol <b>OR</b> 1 mole <b>ALLOW</b> 1 atmosphere/1 atm <b>OR</b> 101 kPa <b>OR</b> 101325 Pa</p>
6	(b)	<p>(i)</p> <p><b>Complete</b> circuit with voltmeter <b>AND</b> labelled salt bridge linking two half-cells ✓</p>  <p>Cu electrode in Cu<sup>2+</sup> ✓</p> <p>Pt electrode in V<sup>2+</sup> <b>AND</b> V<sup>3+</sup> ✓</p> <p>Cu shown as + <b>AND</b> Pt shown as - ✓</p> <p>electrons in wire <b>AND</b> ions in salt bridge ✓ <i>On diagram or stated</i></p>	5	<p>Half cells can be drawn in either order Half cells must show electrodes dipping into solutions <b>ALLOW</b> small gaps in circuit <b>DO NOT ALLOW</b> half-cell with H<sub>2</sub> added</p> <p><b>IGNORE</b> any stated concentrations <b>IGNORE</b> 'anode' and 'cathode'</p> <p>In salt bridge, <b>ALLOW</b> any stated ion that may be present, e.g. K<sup>+</sup>, NH<sub>4</sub><sup>+</sup>, NO<sub>3</sub><sup>-</sup>, Cu<sup>2+</sup>, V<sup>2+</sup>, V<sup>3+</sup></p> <p><b>IGNORE</b> direction of travel of ions and electrons.</p> <p><b>ALLOW</b> Cu half cell as + <b>AND</b> V half cell as -</p>

Question			Answer	Marks	Guidance
6	(b)	(ii)	0.60 <b>OR</b> 0.6 (V) ✓	1	<b>IGNORE</b> any sign
6	(c)		<p><b>Definitions: 1 mark</b> Oxidising agent removes/accepts/gains electrons <b>OR</b> increases oxidation number (of another species) <b>AND</b> Reducing agent adds/donates/loses electrons <b>OR</b> decreases oxidation number (of another species) ✓</p> <p><b>Oxidising agent: 2 marks</b> Cr<sup>3+</sup> oxidises Al <b>OR</b> Cr<sup>3+</sup> acts as oxidising agent <b>AND</b> 3Cr<sup>3+</sup> + Al → 3Cr<sup>2+</sup> + Al<sup>3+</sup> ✓ <b>Explanation (dependent on Cr<sup>3+</sup> oxidising Al above)</b> E of redox system 2 (Cr<sup>3+</sup>/Cr<sup>2+</sup>) is <b>more</b> positive /<b>less</b> negative (than E of system 1 (Al<sup>3+</sup>/Al)) <b>ORA</b>, i.e. in terms of 1 being more negative (than 2) ✓</p> <p><b>Reducing agent: 3 marks</b> Cr<sup>3+</sup> reduces FeO<sub>4</sub><sup>2-</sup>/(H<sup>+</sup>) ✓ 2Cr<sup>3+</sup> + 2FeO<sub>4</sub><sup>2-</sup> + 2H<sup>+</sup> → Cr<sub>2</sub>O<sub>7</sub><sup>2-</sup> + 2Fe<sup>3+</sup> + H<sub>2</sub>O ✓ <b>Explanation (dependent on Cr<sup>3+</sup> reducing FeO<sub>4</sub><sup>2-</sup> above)</b> E of redox system 5 (Cr<sub>2</sub>O<sub>7</sub><sup>2-</sup>/Cr<sup>3+</sup>) is <b>less</b> positive/ <b>more</b> negative (than E of system 6 (FeO<sub>4</sub><sup>2-</sup>/Fe<sup>3+</sup>)) <b>ORA</b>, i.e. in terms of 6 being more positive (than 5) ✓</p>	6	<p><b>FULL ANNOTATIONS MUST BE USED</b></p> <p>-----</p> <p><b>ALLOW</b> oxidising agent decreases its oxidation number <b>AND</b> reducing agent increases its oxidation number</p> <p><b>IGNORE</b> oxidising agent oxidises/is reduced <b>OR</b> reducing agent reduces/is oxidised</p> <p><b>In equations,</b></p> <ul style="list-style-type: none"> <li><b>IGNORE</b> state symbols (even if incorrect)</li> <li><b>ALLOW</b> ⇌ in equation</li> </ul> <p><b>IF</b> more than one equation shown for <b>Cr<sup>3+</sup> as oxidising agent</b>, <b>CON</b> and <b>zero marks</b> for 2 oxidising agent marks <b>IGNORE</b> equations with Cr<sup>2+</sup> as reactant</p> <p>Explanations <b>MUST</b> be in terms of positive/negative: <b>IGNORE</b> 'higher' E <b>OR</b> 'greater'</p> <p><b>ALLOW</b> E<sub>cell</sub> = +1.25 V (+ sign required)</p> <p><b>IF</b> more than one equation shown for <b>Cr<sup>3+</sup> as a reducing agent</b>, <b>CON</b> and <b>zero marks</b> for 3 reducing agent marks <b>IGNORE</b> equations with Cr<sup>2+</sup> as reactant</p> <p>Explanations <b>MUST</b> be in terms of positive/negative: <b>IGNORE</b> 'higher' E <b>OR</b> 'greater'</p> <p><b>ALLOW</b> E<sub>cell</sub> = +0.87 V (+ sign required)</p>
			<b>Total</b>	<b>14</b>	

Question			Answer	Marks	Guidance
7	(a)	(i)	<p><b>IGNORE</b> any charges shown within complexes (treat as rough working)</p> <p><b>Complex ion C:</b> <math>[\text{Ni}(\text{H}_2\text{O})_6]^{2+}</math> ✓</p> <p><b>Solid D:</b> <math>\text{Ni}(\text{OH})_2</math> ✓</p> <p><b>Complex ion E:</b> <math>[\text{Ni}(\text{CN})_4]^{2-}</math> ✓</p>	3	<p><b>ALLOW</b> +2 and –2 for charges</p> <p>Square brackets required</p> <p><b>ALLOW</b> <math>\text{Ni}(\text{H}_2\text{O})_4(\text{OH})_2</math>  <math>(\text{H}_2\text{O})_4</math> and <math>(\text{OH})_2</math> in any order</p> <p><b>IGNORE</b> any square brackets</p> <p>Square brackets required</p> <p><b>TAKE CARE</b> for round brackets within complex ion, i.e. <math>(\text{H}_2\text{O})</math>, <math>(\text{OH})</math> and <math>(\text{CN})</math></p>

Question			Answer	Marks	Guidance
7	(a)	(ii)	<p><b>Mark independently of 7(a)(i)</b>  <b>ALLOW</b> +2 and –2 for charges  <b>IGNORE</b> any charges shown within complexes (treat as rough working)</p> $\text{Ni}^{2+} + 2\text{OH}^- \rightarrow \text{Ni}(\text{OH})_2 \checkmark$ <p>Type of reaction: precipitation ✓  <b>INDEPENDENT of equation</b></p> $[\text{Ni}(\text{H}_2\text{O})_6]^{2+} + 4\text{CN}^- \rightarrow [\text{Ni}(\text{CN})_4]^{2-} + 6\text{H}_2\text{O}(\text{l}) \checkmark$ <p>Type of reaction: ligand substitution ✓  <b>INDEPENDENT of equation</b></p>	4	<p><b>For equations: IGNORE</b> state symbol (even if wrong)  Square brackets <b>not</b> required for <math>\text{Ni}(\text{OH})_2</math></p> <p><b>ALLOW</b> <math>[\text{Ni}(\text{H}_2\text{O})_6]^{2+} + 2\text{OH}^- \rightarrow [\text{Ni}(\text{H}_2\text{O})_4(\text{OH})_2] + 2\text{H}_2\text{O}</math>  <b>ALLOW</b> <math>[\text{Ni}(\text{H}_2\text{O})_6]^{2+} + 2\text{OH}^- \rightarrow \text{Ni}(\text{OH})_2 + 6\text{H}_2\text{O}</math>  <b>ALLOW</b> <math>\text{NiSO}_4(\text{aq}) + 2\text{OH}^-(\text{aq}) \rightarrow \text{Ni}(\text{OH})_2(\text{s}) + \text{SO}_4^{2-}(\text{aq})</math>  <b>ALLOW</b> <math>\text{NiSO}_4(\text{aq}) + 2\text{KOH}(\text{aq}) \rightarrow \text{Ni}(\text{OH})_2(\text{s}) + \text{K}_2\text{SO}_4(\text{aq})</math></p> <p><b>ALLOW</b> acid/base <b>OR</b> neutralisation <b>OR</b> deprotonation  <b>ONLY IF</b> <math>[\text{Ni}(\text{H}_2\text{O})_6]^{2+}</math> <b>AND</b> <math>[\text{Ni}(\text{H}_2\text{O})_4(\text{OH})_2]</math> used</p> <p><b>ALLOW</b> precipitate</p> <p><b>ALLOW</b> <math>[\text{Ni}(\text{H}_2\text{O})_6]^{2+} + 4\text{KCN} \rightarrow [\text{Ni}(\text{CN})_4]^{2-} + 6\text{H}_2\text{O} + 4\text{K}^+</math></p> <p><b>LOOK</b> at formulae for <b>E</b> from <b>7(a)(i)</b> (copied at bottom)  <b>ALLOW ECF</b> in <b>7a(ii)</b> Equation for no round brackets around CN, i.e. <math>[\text{NiCN}_4]^{2-}</math> in <b>7a(i)</b>  This is the only <b>ECF</b> allowed from <b>7a(i)</b> structures.</p> <p><b>ALLOW</b> ligand exchange</p>
7	(b)	(i)	linear ✓	1	<b>IGNORE</b> planar

Question			Answer	Marks	Guidance
7	(b)	(ii)	Au/Gold has been <b>oxidised</b> from 0 to +1 ✓  O/Oxygen/O <sub>2</sub> has been <b>reduced</b> from 0 to -2 ✓	2	<b>IF</b> Ag referred to, rather than Au, treat as a slip and apply <b>BOD ALLOW</b> 0 to 1 (i.e. no + sign for +1)  <b>ALLOW</b> 1 mark for <b>ALL</b> oxidation numbers correct with no oxidised or reduced <b>OR</b> oxidation and reduction wrong way round, e.g. Au goes from 0 to +1 and O goes from 0 to -2 ✓ Au is reduced from 0 to +1 and O is oxidised from 0 to -2 ✓
7	(b)	(iii)	<b>IGNORE</b> any charges shown within complexes (treat as rough working) $4\text{Au} + 8\text{CN}^- + 2\text{H}_2\text{O} + \text{O}_2 \rightarrow 4[\text{Au}(\text{CN})_2]^- + 4\text{OH}^-$ ✓✓  First mark for all 6 species  Second mark for balancing	2	<b>IF</b> Ag referred to, rather than Au, treat as a slip and apply <b>BOD</b>  <b>IGNORE</b> state symbols <b>CARE:</b> In $[\text{Au}(\text{CN})_2]^-$ , - sign is <b>OUTSIDE</b> square brackets  <b>For 1st mark, IGNORE</b> e <sup>-</sup> present  <b>ALLOW</b> 1 mark for balanced equation with CN <sup>-</sup> missing, i.e. $4\text{Au} + 2\text{H}_2\text{O} + \text{O}_2 \rightarrow 4\text{Au}^+ + 4\text{OH}^-$  <b>ALLOW</b> 1 mark rogue e <sup>-</sup> on either side  <b>ALLOW</b> multiples, e.g. $2\text{Au} + 4\text{CN}^- + \text{H}_2\text{O} + \frac{1}{2}\text{O}_2 \rightarrow 2[\text{Au}(\text{CN})_2]^- + 2\text{OH}^-$ $\text{Au} + 2\text{CN}^- + \frac{1}{2}\text{H}_2\text{O} + \frac{1}{4}\text{O}_2 \rightarrow [\text{Au}(\text{CN})_2]^- + \text{OH}^-$
7	(b)	(iv)	$\text{ClO}^- + 2\text{H}^+ + 2\text{e}^- \rightarrow \text{Cl}^- + \text{H}_2\text{O}$ ✓	1	<b>IGNORE</b> state symbols  <b>ALLOW</b> e for electron  <b>ALLOW</b> multiples
<b>Total</b>				<b>13</b>	




Question		Answer	Marks	Guidance
8	(a)	$\text{Cu}^{2+}: (1s^2) 2s^2 2p^6 3s^2 3p^6 3d^9 \checkmark$ $\text{Cu}^+: (1s^2) 2s^2 2p^6 3s^2 3p^6 3d^{10} \checkmark$	2	<p><b>IGNORE</b> repeated <math>1s^2</math> after <math>1s^2</math> prompt on answer line  <b>ALLOW</b> <math>4s^0</math>, either before or after <math>3d</math></p> <p><b>ALLOW</b> upper case D, etc and subscripts,  e.g. ....<math>3S_23P^6</math>  <b>DO NOT ALLOW</b> [Ar] as shorthand for <math>1s^2 2s^2 2p^6 3s^2 3p^6</math></p>
8	(b)	<p><b>IGNORE</b> any charges shown within formulae  (treat as rough working)</p> $\text{CuCO}_3 + 2\text{HCOOH} \rightarrow \text{Cu}(\text{HCOO})_2 + \text{H}_2\text{O} + \text{CO}_2$ <p><b>OR</b> <math>\text{CuO} + 2\text{HCOOH} \rightarrow \text{Cu}(\text{HCOO})_2 + \text{H}_2\text{O}</math></p> <p><b>OR</b> <math>\text{Cu}(\text{OH})_2 + 2\text{HCOOH} \rightarrow \text{Cu}(\text{HCOO})_2 + 2\text{H}_2\text{O} \checkmark</math></p>	1	<p><b>IGNORE</b> state symbols  In formula of <math>\text{HCOOH}/\text{HCOO}</math>, <b>ALLOW</b> H, C and O in <b>ANY</b> order  <b>ALLOW</b> <math>\text{H}_2\text{CO}_3</math> for <math>\text{H}_2\text{O}</math> and <math>\text{CO}_2</math> in carbonate equation</p> <p><b>ALLOW</b> <math>(\text{HCOO})_2\text{Cu}</math> for <math>\text{Cu}(\text{HCOO})_2</math></p> <p><b>DO NOT ALLOW</b> equation with <math>\text{CuSO}_4</math></p>
8	(c)	$2\text{Cu}^{2+} + 4\text{I}^- \rightarrow 2\text{CuI}(\text{s}) + \text{I}_2 \checkmark$ State symbol for $\text{CuI}(\text{s})$ <b>ONLY</b> required	1	<p><b>ALLOW</b> multiples, e.g. <math>\text{Cu}^{2+} + 2\text{I}^- \rightarrow \text{CuI}(\text{s}) + \frac{1}{2}\text{I}_2</math></p> <p><b>IGNORE</b> other state symbols, even if incorrect</p>
8	(d)	Starch $\checkmark$ Blue/black to colourless/white $\checkmark$ <b>MARK INDEPENDENTLY</b>	2	<p><b>IGNORE</b> 'brown' in composite colour with blue or black, i.e.  <b>ALLOW</b> blue/brown to colourless  <b>ALLOW</b> black/brown to colourless</p> <p><b>DO NOT ALLOW</b> just 'it turns colourless/is decoloured'  <i>Initial colour required</i></p> <p><b>IGNORE</b> clear for colourless</p>

Question	Answer	Marks	Guidance
8 (e)	<p><b>WORKING REQUIRED</b>            Correct answer: <math>x = 4</math> required evidence of working            -----  <math>n(\text{S}_2\text{O}_3^{2-})</math> OR <math>n(\text{Cu}^{2+}) = \frac{0.0420 \times 23.5}{1000} = 9.87 \times 10^{-4}</math> (mol) ✓              In 250.0 cm<sup>3</sup> solution, <math>n(\text{Cu}^{2+}) = 9.87 \times 10^{-3}</math> (mol) ✓    <math>M(\text{Cu}(\text{HCOO})_2 \cdot 4\text{H}_2\text{O}) = \frac{2.226}{9.87 \times 10^{-3}} = 225.5</math> (g mol<sup>-1</sup>) ✓    <math>x(\text{H}_2\text{O})</math> has mass of <math>225.5 - M(\text{Cu}(\text{HCOO})_2)</math>  <math>= 225.5 - 153.5</math>  <math>= 72(.0)</math> ✓    <math>x = \frac{72(.0)}{18(.0)} = 4</math>                      <b>WHOLE NUMBER</b> needed  <b>AND</b>            evidence of working ✓</p>	5	<p><b>FULL ANNOTATIONS MUST BE USED</b>            -----            At least <b>3 SF</b> required throughout    <i>Alternative approach for final 3 marks based on mass:</i>            mass <math>\text{Cu}(\text{HCOO})_2 = 9.87 \times 10^{-3} \times 153.5 = 1.515</math> g ✓  <math>n(\text{H}_2\text{O}) = \frac{2.226 - 1.515}{18(.0)} = \frac{0.711}{18(.0)} = 0.0395</math> (mol) ✓    <math>x = \frac{0.0395}{9.87 \times 10^{-3}} = 4</math> ✓    <b>ALLOW</b> <math>\text{Cu}(\text{HCOO})_2 \cdot 4\text{H}_2\text{O}</math>            -----  <b>COMMON ERRORS for 4 marks</b>  <math>x = 117</math> (calc 116.78)  <i>Use of <math>9.87 \times 10^{-4}</math> (no scaling <math>\times 10</math>) <math>\rightarrow M = 2255.319</math></i>    <math>x = 17</math> (calc 16.53)                      4 marks  <i>Use of <math>4.935 \times 10^{-4}</math> (Use of <math>0.5 \times 9.87 \times 10^{-3}</math>)</i>              Check <math>n(\text{Cu}^{2+})</math> for other <b>ECFs</b>            Check for <b>ECFs</b> from incorrect <math>M(\text{anhydr salt})</math>    Actual = 153.5</p>
	<b>Total</b>	<b>11</b>	

## APPENDIX Q3(b)


Extra energy line placed **ABOVE** top line  
3 out of 4 marks awarded for energy lines and species.

Top arrow is shown **FROM**  $2\text{K}^+(\text{g}) + \text{SO}_4^{2-}(\text{g})$  and arrow directions correct. Letter labels correct so last mark is awarded. **4/5 marks**




Same as left-hand response

**BUT** top arrow shown **TO**  $2\text{K}^+(\text{g}) + \text{SO}_4^{2-}(\text{g})$  so last mark not awarded  
**3/5 marks**




Extra energy line placed **BELOW** bottom line  
3 out of 4 marks awarded for energy lines and species.

Top arrow is shown **FROM**  $\text{K}_2\text{SO}_4(\text{s})$  and arrow directions correct. Letter labels correct so last mark is awarded. **4/5 marks**



Same as left-hand response

**BUT** bottom arrow shown **TO**  $\text{K}_2\text{SO}_4(\text{s})$  so last mark not awarded  
**3/5 marks**



**OCR (Oxford Cambridge and RSA Examinations)**  
**1 Hills Road**  
**Cambridge**  
**CB1 2EU**

**OCR Customer Contact Centre**

**Education and Learning**

Telephone: 01223 553998

Facsimile: 01223 552627

Email: [general.qualifications@ocr.org.uk](mailto:general.qualifications@ocr.org.uk)

[www.ocr.org.uk](http://www.ocr.org.uk)

For staff training purposes and as part of our quality assurance programme your call may be recorded or monitored

**Oxford Cambridge and RSA Examinations**  
**is a Company Limited by Guarantee**  
**Registered in England**  
**Registered Office; 1 Hills Road, Cambridge, CB1 2EU**  
**Registered Company Number: 3484466**  
**OCR is an exempt Charity**

**OCR (Oxford Cambridge and RSA Examinations)**  
**Head office**  
**Telephone: 01223 552552**  
**Facsimile: 01223 552553**

© OCR 2016

