

**Mark Scheme 4724
June 2006**

1	$\frac{d}{dx}(xy) = x \frac{dy}{dx} + y$ $\frac{d}{dx}(y^2) = 2y \frac{dy}{dx}$ <p>Substitute (1,2) into their differentiated equation and attempt to solve for $\frac{dy}{dx}$. [Allow subst of (2,1)]</p> $\frac{dy}{dx} = -2$	<p>B1</p> <p>B1</p> <p>M1 dep at least 1 x B1</p> <p>A1</p>	<p>s.o.i. e.g. $2x \frac{dy}{dx} + y$</p> <p>Or attempt to solve their diff equation for $\frac{dy}{dx}$ and then substitute (1,2)</p> <p>4</p>
<hr/>			
2	<p>(i) $1 + (-2)(-3x) + \frac{(-2)(-3)}{1.2}(-3x)^2$ (+ ... ignore)</p> <p>= 1 + 6x</p> <p>... + 27x²</p> <p>(ii) $(1 + 2x)^2(1 - 3x)^{-2}$</p> <p>Attempt to expand $(1 + 2x)^2$ & select (at least) 2 relevant products and add</p> <p>55 (Accept 55x²)</p> <p><u>SR 1</u> For expansion of $(1 + 2x)^2$ with 1 error, A1√</p> <p><u>SR 2</u> For expansion of $(1 + 2x)^2$ & > 1 error, A0</p> <p>Alternative Method</p> <p>For correct method idea of long division</p> <p>1 +10x +55x²</p>	<p>M1</p> <p>B1</p> <p>A1</p> <p>M1</p> <p>A2√</p> <p>M1</p> <p>A1,A1,A1(4)</p>	<p>State or imply; accept $-3x^2$ & $-9x^2$</p> <p>Correct first 2 terms</p> <p>Correct third term</p> <p>For changing into suitable form, seen/implied</p> <p>Selection may be after multiplying out</p> <p>4 If (i) is $a + bx + cx^2$, f.t. $4(a + b) + c$</p>
<hr/>			
3	<p>(i) $\frac{A}{x} + \frac{B}{3-x}$ & c-u rule or $A(3-x) + Bx \equiv 3 - 2x$</p> <p>$\frac{1}{x}$</p> <p>$-\frac{1}{3-x}$</p> <p>(ii) $\int \frac{1}{x}(dx) = \ln x$ or $\ln x$</p> <p>$\int \frac{1}{3-x}(dx) = -\ln(3-x)$ or $-\ln 3-x$</p> <p>Correct method idea of substitution of limits</p> <p>$\ln 2 (+ \ln 1 - \ln 1) - \ln 2 = 0$</p> <p>Alternative Method</p> <p>If ignoring PFs, $\ln x(3-x)$ immediately</p> <p>As before</p> <p>(iii) Suitable statement or clear implication e.g. Equal amounts (of area) above and below (axis) or graph crosses axis or there's a root (Be lenient)</p>	<p>M1</p> <p>A1</p> <p>A1</p> <p>B1</p> <p>B1</p> <p>M1</p> <p>A1</p> <p>B2</p> <p>M1,A1 (4)</p> <p>B1</p>	<p>Correct format + suitable method</p> <p>seen in (i) or (ii)</p> <p>3 ditto; $\frac{1}{x} - \frac{1}{3-x}$ scores 3 immediately</p> <p>Check sign carefully; do not allow $\ln(x-3)$</p> <p>Dep on an attempt at integrating</p> <p>Clearly seen; WWW AG</p> <p>$\ln x(x-3) \rightarrow 0$</p> <p>1</p>

4	(i) Working out $\mathbf{b} - \mathbf{a}$ or $\mathbf{a} - \mathbf{b}$ or $\mathbf{c} - \mathbf{a}$ or $\mathbf{a} - \mathbf{c}$	M1) Irrespective of label
	= $\pm(-3\mathbf{i} - \mathbf{j} - \mathbf{k})$ or $\pm(-2\mathbf{i} + \mathbf{j} - 2\mathbf{k})$	A1) If not scored, these 1 st 3 marks can be
	Method for finding magnitude of <u>any</u> vector	M1) awarded in part (ii)
	Method for finding scalar product of <u>any</u> 2 vectors	M1	
	Using $\cos \theta = \frac{a \cdot b}{ a b }$ AEF for <u>any</u> 2 vectors	M1	
	[Alternative cosine rule method] $ \vec{BC} = \sqrt{6}$	B1	
	Cosine rule used	M1	'Recognisable' form
	$45.3^\circ, 0.79(0), \frac{\pi}{3.97}$ (45.289378, 0.7904487)	A1	6 Do not accept supplement (134.7 etc)

(ii) Use of $\frac{1}{2} \vec{AB} \vec{AC} \sin \theta$	M1	Accept $\left \frac{1}{2} \vec{AB} \times \vec{AC} \right $
3.54 (3.5355) or $\frac{5\sqrt{2}}{2}$	A1	2 Accept from correct supp (134.7 etc)

5	(i) $\frac{dA}{dt}$ or kA^2 seen	M1	
	$\frac{dA}{dt} = kA^2$	A1	2

(ii) Separate variables + attempt to integrate	*M1	Accept if based on $\frac{dA}{dt} = kA^2$ or A^2
$-\frac{1}{A} = kt + c$ or $-\frac{1}{kA} = t + c$ or $-\frac{1}{A} = t + c$	A1	
Subst one of (0,0), (1,1000) or (2,2000) into eqn.	dep*M1	Equation must contain k and/or c
Subst another of (0,0), (1,1000) or (2,2000) into eqn	dep*M1	This equation must contain k <u>and</u> c
Substitute $A = 3000$ into eqn with k and c subst	dep*M1	
$t = \frac{7}{3}$ ISW	A1	6 Accept 2.33, 2h 20 m

6	(i) Attempt to connect du and dx e.g. $\frac{du}{dx} = e^x$	M1	But not $du = dx$
	Use of $e^{2x} = (e^x)^2$ or $(u-1)^2$ s.o.i.	A1	
	Simplification to $\int \frac{u-1}{u} (du)$ WWW	A1	3 AG

(ii) Change $\frac{u-1}{u}$ to $1 - \frac{1}{u}$ or use parts	M1	If parts, may be twice if $\int \ln x dx$ is involved
$\int \frac{1}{u} du = \ln u$	A1	Seen anywhere in this part
<u>Either</u> attempt to change limits <u>or</u> resubstitute	M1 (indep)	Expect new limits $e+1$ & 2
Show as $e+1 - \ln(e+1) - \{2 \text{ or } (1+1)\} + \ln 2$	A1	
WWW show final result as $e-1 - \ln\left(\frac{e+1}{2}\right)$	A1	5 AG

7	<p>(i) Produce at least 2 of the 3 relevant eqns in λ and μ M1 e.g. $1 + 3\lambda = -8 + \mu$, $-2 + \lambda = 2 - 2\mu$</p> <p>Solve the 2 eqns in λ & μ as far as $\lambda = \dots$ or $\mu = \dots$ M1</p> <p>1st solution: $\lambda = -2$ or $\mu = 3$ A1</p> <p>2nd solution: $\mu = 3$ or $\lambda = -2$ f.t. A1✓</p> <p>Substitute their λ and μ into 3rd eqn and find 'a' M1</p> <p>Obtain $a = 2$ & clearly state that a cannot be 2 A1 6</p>
	<p>(ii) Subst their λ or μ (& poss a) into either line eqn M1</p> <p>Point of intersection is $-5\mathbf{i} - 4\mathbf{j}$ A1 2 Accept any format <u>No f.t. here</u></p> <p>N.B. In this question, award marks irrespective of labelling of parts</p>
8	<p>(i) <u>Integration method</u></p> <p>Attempt to change $\cos^2 6x$ into $f(\cos 12x)$ M1</p> <p>$\cos^2 6x = \frac{1}{2}(1 + \cos 12x)$ A1 with $\cos^2 6x$ as the subject of the formula</p> <p>$\int = \frac{1}{2}x + \frac{1}{24}\sin 12x + c$ A1 AG Accept $\frac{1}{2}\left(x + \frac{1}{12}\sin 12x\right)$</p> <p><u>Differentiation method</u></p> <p>Differentiate RHS producing $\frac{1}{2} + \frac{1}{2}\cos 12x$ ---(E) B1</p> <p>Attempt to change $\cos 12x$ into $f(\cos 6x)$ M1 Accept $+/- 2\cos^2 6x + /- 1$</p> <p>Simplify (E) WWW to $\cos^2 6x$ + satis finish A1 3</p> <hr/> <p>(ii) Parts with $u = x$, $dv = \cos^2 6x$ *M1</p> <p>$x\left(\frac{1}{2}x + \frac{1}{24}\sin 12x\right) - \int\left(\frac{1}{2}x + \frac{1}{24}\sin 12x\right)dx$ A1 Correct expression only</p> <p>$\int \sin 12x dx = -\frac{1}{12}\cos 12x$ B1 Clear indication somewhere in this part</p> <p>Correct use of limits to <u>whole</u> integral dep*M1 Accept () (-0)</p> <p>$\frac{\pi^2}{288} - \frac{\pi^2}{576} - \frac{1}{288} - \frac{1}{288}$ A1 AE unsimp exp. Accept $12x24, \sin \pi$ here</p> <p>$\frac{\pi^2}{576} - \frac{1}{144}$ +A1 6 Tolerate e.g. $\frac{2}{288}$ here</p> <p>S.R. If final marks are A0 + A0, allow SR A1 for 0.01/0.010/0.0101/0.0102/0.0101902</p>

9	<p>(i) $\frac{dy}{dx} = \frac{\frac{dy}{dt}}{\frac{dx}{dt}}$</p> <p>$\frac{dx}{dt} = -4 \sin t$ or $\frac{dy}{dt} = 3 \cos t$</p> <p>$\frac{dy}{dx} = -\frac{3 \cos t}{4 \sin t}$ or $\frac{3 \cos t}{-4 \sin t}$ ISW</p> <p>SR: M1 for Cartesian eqn attempt + B1 for $\frac{d}{dx}(y^2) = 2y \frac{dy}{dx}$ + A1 as before (must be in terms of t)</p>	<p>M1</p> <p>*B1</p> <p>dep*A1</p>	<p>Used, not just quoted</p> <p>Also $\frac{-3 \cos t}{4 \sin t}$ provided B0 not awarded</p>
<hr/>			
(ii)	<p>$y - 3 \sin p = \left(\text{their } \frac{dy}{dx} \right) (x - 4 \cos p)$</p> <p>or $y = \left(\text{their } \frac{dy}{dx} \right) x + c$ & subst cords to find c</p> <p>$4y \sin p - 12 \sin^2 p = -3x \cos p + 12 \cos^2 p$</p> <p>or $c = \frac{12 \sin^2 p + 12 \cos^2 p}{4 \sin p}$</p> <p>$3x \cos p + 4y \sin p = 12$ WWW</p>	<p>M1</p> <p>A1</p> <p>A1</p> <p>A1</p>	<p>Accept p or t here</p> <p>Ditto</p> <p>Correct equation cleared of fractions</p> <p>3 AG Only p here. Mixture earlier \rightarrow A0</p>
<hr/>			
(iii)	<p>Subst $x = 0$ and $y = 0$ separately in tangent eqn</p> <p>Produce $\frac{3}{\sin p}$ and $\frac{4}{\cos p}$</p> <p>Use $\Delta = \frac{1}{2} \left(\frac{3}{\sin p} \cdot \frac{4}{\cos p} \right) = \frac{12}{\sin 2p}$ WWW</p>	<p>M1</p> <p>A1</p> <p>A1</p>	<p>to find R & S</p> <p>Accept $\frac{12}{4 \sin p}$ and/or $\frac{12}{3 \cos p}$</p> <p>3 AG</p>
<hr/>			
(iv)	<p>Least area = 12</p> <p>$p = \frac{1}{4} \pi$ as final or only answer</p> <p>S.R. $45^\circ \rightarrow$ B1 ;</p>	<p>B1</p> <p>B2</p>	<p>3 These B marks are independent.</p> <p>S.R. [-12 and e.g. $-\pi/4 \rightarrow$ B1]</p>

