

Please write clearly in	า block capitals.	
Centre number	Candidate number	
Surname		
Forename(s)		
Candidate signature	I declare this is my own work.	ノ

GCSE CHEMISTRY

F

Foundation Tier Paper 1

Time allowed: 1 hour 45 minutes

Materials

For this paper you must have:

- a ruler
- · a scientific calculator
- the periodic table (enclosed).

Instructions


- Use black ink or black ball-point pen.
- · Pencil should only be used for drawing.
- Fill in the boxes at the top of this page.
- Answer all questions in the spaces provided. Do not write outside the box around each page or on blank pages.
- If you need extra space for your answer(s), use the lined pages at the end of this book. Write the question number against your answer(s).
- Do all rough work in this book. Cross through any work you do not want to be marked.
- In all calculations, show clearly how you work out your answer.

Information

- The maximum mark for this paper is 100.
- The marks for questions are shown in brackets.
- You are expected to use a calculator where appropriate.
- You are reminded of the need for good English and clear presentation in your answers.

For Examiner's Use					
Question	Mark				
1					
2					
3					
4					
5					
6					
7					
8					
9					
10					
TOTAL	li de la companya de				







0 1	This question is about elements, compounds and mixtures.	
0 1.1	Substance A contains only one type of atom.	
	Substance A does not conduct electricity.	
	Which type of substance is A ?	[4 mouls]
	Tick (✓) one box.	[1 mark]
	Compound	
	Metallic element	
	Mixture	
	Non-metallic element	
0 1.2	Substance B contains two types of atoms. The atoms are chemically combined together in fixed proportions. Which type of substance is B ? Tick (✓) one box. Compound Metallic element Mixture Non-metallic element	[1 mark]



Do not write outside the box


0 1.3	What is the name of the elements in Group 0 of the periodic table?			
	Tick (✓) one box.	[1 mark]		
	Alkali metals			
	Halogens			
	Noble gases			
	Transition metals			
0 1.4	Which statement about the elements in Group 0 is correct?	[4 mork]		
	Tick (✓) one box.	[1 mark]		
	All elements in the group are very reactive.			
	All elements in the group form negative ions.			
	The boiling points increase down the group.			
	The relative atomic masses (<i>A</i> _r) decrease down the group.			



Neon is in Group 0.		
What type of particles a	re in a sample of neon?	[4 magula]
Tick (✓) one box.		[1 mark]
Atoms		
lons		
Molecules		
	What type of particles a Tick (✓) one box. Atoms lons	What type of particles are in a sample of neon? Tick (✓) one box. Atoms

0 1. 6 Figure 1 represents part of the structure of an oxide of a metal.

Figure 1



Determine the empirical formula of this oxide.

[1 mark]


Empirical formula = XO_____



A nanoparticle of a metallic element is a cube.

Figure 2 shows a diagram of the nanoparticle.

Figure 2



0	1		7	The surface are	a of a	cube is	given b	y the ed	quation
---	---	--	---	-----------------	--------	---------	---------	----------	---------

surface area = $(length of side)^2 \times 6$

Calculate the surface area of the cube in Figure 2.

Give your answer to 2 significant figures.

[3	ma	rks]
L		

-			
-			

Surface area (2 significant figures) = nm²

0 1 . 8	Fine and coarse particles of the metallic element are also cubes.	l
	The length of a fine particle cube is 10 times smaller than the length of a coarse particle cube.	
	How does the surface area to volume ratio of the fine particle cube compare with that of the coarse particle cube? [1 mark]	
	Tick (✓) one box.	
	Both surface area to volume ratios are the same.	
	The surface area to volume ratio of the fine particle is 10 times greater.	_
	The surface area to volume ratio of the fine particle is 10 times smaller.	1

Turn over for the next question



		8					
2	This question is about chem	This question is about chemical cells and batteries.					
2 . 1	. 1 Three different types of battery can be used to power a TV remote control.						
	Table 1 gives information al	bout these batter	ies.				
		Table 1					
		Zinc-carbon battery	Alkaline battery	Nickel- metal hydride			
				battery			
Cos	st of battery in £ (pounds)	0.17	0.50	1.50			
	st of battery in £ (pounds)	0.17 No	0.50 No	•			

Give one advantage	of each type	of battery.
--------------------	--------------	-------------

[3 marks]

Zinc-carbon			
Alkaline			
Nickel-metal hydride			

0 2 . Figure 3 shows a symbol printed on batteries.



This symbol shows that batteries should not be put in household waste.

Suggest why batteries should **not** be put in household waste.

[1 mark]







Figure 4



0 2. 3 The order of reactivity of three metals is shown below.

Iron (Most reactive)

Tin

Copper (Least reactive)

Which combination of metal electrodes would give the highest voltage in the chemical cell in **Figure 4**?

[1 mark]

Copper and iron	

Tick (✓) one box.

Iron and tin

Tin and copper

The voltage produced by the cell in **Figure 4** depends on the type of electrodes and the type of electrolyte.

Suggest **one** other factor that could affect the voltage produced.

[1 mark]



0 2 . 5	Water is produced in a hydroge	n fuel cell.		Do not write outside the box	
	Complete the word equation to show the reaction that produces water in a hydrogen fuel cell.				
	ruer cen.			[2 marks]	
			water	8	
	+ _		→ water		



0 3	This question is	about Group 1 elements.		
0 3.1	Complete Table	2 to show the electronic s	tructure of a potassium a	tom. [1 mark]
		Table 2		
	Atom	Number of electrons	Electronic structure	
	Sodium	11	2,8,1	
	Potassium	n 19		
0 3.2	Tick (✓) one box They have the sa	elements have similar che ame number of electron shame number of outer shell lectrons in the first shell.	nells.	[1 mark]
0 3.3	What is the type Tick (✓) one box Covalent Ionic Metallic	of bonding in sodium?		[1 mark]



Table 3 shows observations made when lithium, potassium and rubidium react with water.

Table 3

Element	Observations
Lithium	Bubbles slowly Floats Moves slowly
Sodium	1
Potassium	Bubbles very quickly Melts into a ball Floats Moves very quickly Flame
Rubidium	Sinks Melts into a ball Explodes with a flame

0	3	. 4	Give two observations you could make when sodium reacts with water
---	---	-----	--

Write your answers in **Table 3**.

[2 marks]



Do not write outside the box

0 3.5	How does the reactivity of the elements change going down Group 1? [1 mark]
0 3.6	Give two ways in which the observations in Table 3 show the change in reactivity going down Group 1. [2 marks]
	2
0 3.7	Which gas is produced when Group 1 elements react with water? [1 mark] Tick (✓) one box.
	Carbon dioxide Hydrogen
	Nitrogen
	Oxygen



0 3 . 8

Sodium fluoride is an ionic compound.

Figure 5 shows dot and cross diagrams for a sodium atom and a fluorine atom.

Complete **Figure 5** to show what happens when a sodium atom and a fluorine atom react to produce sodium fluoride.


You should:

Sodium ion

- complete the electronic structures of the sodium ion and the fluoride ion
- give the charges on the sodium ion and the fluoride ion.

[3 marks]

Figure 5







Fluoride ion

Do not write outside the Turn over for the next question DO NOT WRITE ON THIS PAGE ANSWER IN THE SPACES PROVIDED



0 4

A student investigated the reactivity of metals with hydrochloric acid.

This is the method used.

- 1. Measure 50 cm³ of hydrochloric acid into a polystyrene cup.
- 2. Measure the temperature of the hydrochloric acid.
- 3. Add one spatula of metal powder to the hydrochloric acid and stir.
- 4. Measure the highest temperature the mixture reaches.
- 5. Calculate the temperature increase for the reaction.
- 6. Repeat steps 1 to 5 three more times.
- 7. Repeat steps 1 to 6 with different metals.

Table 4 shows the student's results.

Table 4

Metal	Temp	perature	increase	in °C	Mean temperature
IVIELAI	Trial 1	Trial 2	Trial 3	Trial 4	increase in °C
Cobalt	6	7	5	9	7
Magnesium	54	50	37	55	x
Zinc	18	16	18	20	18

0 4.1	Calculate the mean temperature increase X for magnesium in Table 4 .	
	Do not include the anomalous result in your calculation.	[2 marks]
	X =	°C




0 4.2	Determine the order of reactivity for the metals cobalt, magnesium and zinc.	
	Use Table 4.	·4 1.9
	Most reactive	1 mark]
	Least reactive	
0 4.3	The range of measurements either side of the mean shows the uncertainty in t mean temperature increase.	:he
	Complete the sentence.	
	Use Table 4 .	1 mark]
	The mean temperature increase for zinc is 18 ±°C	,
0 4.4	What type of variable is the volume of hydrochloric acid in this investigation? Tick (✓) one box.	[1 mark]
	Control Dependent	
	Independent	
0 4.5	Suggest one way of improving step 3 in the method to give results which are repeatable.	more [1 mark]



0 4 . 6

Figure 6 shows a reaction profile for the reaction of magnesium with hydrochloric acid.





What do labels A, B and C represent on Figure 6?

Choose answers from the box.

[3 marks]

9

activation energy	energy	overall energy change
products	progress of reaction	reactants

Α			

В_____

C



0 5	This question is about acids and alkalis.
0 5 . 1	Which ion do acids produce in aqueous solution?
	Tick (✓) one box.
	H ⁺ OH ⁻ O ²⁻
0 5.2	Acids react with alkalis.
	What is the name of this type of reaction?
	Tick (✓) one box.
	Decomposition
	Electrolysis
	Neutralisation
	Redox
0 5.3	Balance the equation for the reaction between sulfuric acid and potassium hydroxide. [1 mark]
	$H_2SO_4 + \underline{\hspace{1cm}} KOH \rightarrow K_2SO_4 + \underline{\hspace{1cm}} H_2O$
0 5.4	Universal indicator turns purple in potassium hydroxide solution.
	What is the pH of the solution?
	Tick (✓) one box.
	1 7 14






Do not write outside the

A student does a titration to find the volume of sulfuric acid that reacts with 25 cm³ of potassium hydroxide solution.

Figure 7 shows the equipment used.

Figure 7



0	5	5	The 25 cm ³ of potassium hydroxide solution is measured with the measuring cylinder.

Which piece of equipment could the student use to measure the 25 cm³ of potassium hydroxide solution more accurately?

[1 mark]

Tick (✓) one box.	
Beaker	
Evaporating basin	
Pipette	
Test tube	



0 5 . 6	Describe how the student would use the equipment in Figure 7 to		outsid bo
	complete the titration.	[5 marks]	
			10

Turn over for the next question




0 6

This question is about metal carbonates.

A student investigated the reaction of copper carbonate with an acid.

Figure 8 shows the apparatus.

Figure 8




This is the method used.

- 1. Pour 25 cm³ of the acid into a conical flask.
- 2. Weigh 0.10 g of copper carbonate.
- 3. Remove the stopper and add the copper carbonate to the flask.
- 4. Quickly replace the stopper.
- 5. Record the maximum volume of gas collected in the gas syringe.
- 6. Repeat steps 1 to 5 with different masses of copper carbonate.

0 6 . 1 Figure 9 shows the gas syringe during the experiment.

Figure 9



What is the reading on the gas syringe?

[1 mark]


cm³



0 6.2 The student plotted the results on a graph.

Figure 10 shows the student's graph.

Figure 10



Determine the gradient of the line of best fit.

You should:

- calculate the values of the change in y and the change in x
- calculate the gradient of the line of best fit.

[4 marks]

Change in y =	cm ³

O t			
Gradient			



			_
0 6.3	Copper chloride was produced in the reaction.		
	Which acid reacts with copper carbonate to produce copper chloride?	[4 mork	
	Tick (✓) one box.	[1 mark]	l
	Hydrochloric acid		
	Nitric acid		
	Sulfuric acid		
0 6.4	The reaction between copper carbonate and the acid produced a gas.		
	What was the gas?	[1 mark]	
	Tick (✓) one box.	[i iliai k	1
	Carbon dioxide		
	Chlorine		
	Hydrogen		
	Oxygen		




Do not write outside the box

Do not write outside the box

A different student produced a pure, dry sample of copper chloride using same reaction.	
This is the method used.	
1. Add excess copper carbonate to the acid.	
2. Filter the mixture.	
3. Heat the solution gently until crystals start to form.	
4. Leave for 24 hours.	
5. Remove the crystals.	
6. Rinse with water and dry the crystals.	
0 6. 5 Why was the solution heated gently in step 3?	54 m and 3
Tick (✓) one box.	[1 mark]
To evaporate acid	
To evaporate copper carbonate	
To evaporate water	
0 6.6 How should the solution be heated gently in step 3?	[1 mark]







0 7


This question is about electrolysis.

Some students investigated the electrolysis of silver nitrate solution.

This electrolysis produces silver at the negative electrode.

Figure 11 shows the apparatus.

Figure 11



This is the method used.

- 1. Weigh the negative electrode.
- 2. Set up the apparatus shown in Figure 11.
- 3. Switch on the power supply.
- 4. Switch off the power supply after five minutes.
- 5. Rinse the negative electrode with water and allow to dry.
- 6. Reweigh the negative electrode.
- 7. Repeat steps 1 to 6 for different times.



Do not write outside the box

0 7.1	Some silver did not stick to the negative electrode but fell to the bottom of the beaker.		
	The students needed to weigh this silve	er.	
	How could the students separate the si	lver from the silver nitrate solution?	[1 mark]
	Tick (✓) one box.		[1 mark]
	By chromatography		
	By crystallisation		
	By distillation		
	By filtration		

Table 5 shows the students' results.

Table 5

Time in minutes	Mass of silver in g
0	0.00
5	0.06
10	0.12
15	0.18
20	0.24
25	0.30




0 7. 2 Draw a graph on Figure 12.

You should:

- use a suitable scale for the x-axis
- plot the data from Table 5
- · draw a line of best fit.

[4 marks]

Figure 12



Time in minutes

0 7 . 3 Determine the mass of silver that would be produced after 12 minutes.

Use Figure 12.

[1 mark]

Mass of silver =

Question 7 continues on the next page



A student investigated the electrolysis of two aqueous salt solutions.

Hydrogen is produced at the negative electrode when the metal in the salt solution is more reactive than hydrogen.

Complete **Table 6** to show what the student would **observe** at the negative electrode for each salt solution.

[2 marks]

Table 6

Salt solution	Observation at negative electrode
Copper sulfate	
Sodium chloride	

0 7. 5 A teacher demonstrates the electrolysis of molten lead bromide

The products at the electrodes are lead and bromine.

Why should the teacher do the demonstration in a fume cupboard?

[1 mark]

0 7 . 6 Two other molten compounds are electrolysed.

Complete **Table 7** to show the molten compounds and the products.

[3 marks]

Table 7

Molten compound electrolysed	Product at the negative electrode	Product at the positive electrode
Zinc chloride		
	Potassium	lodine

12



Do not write outside the Turn over for the next question DO NOT WRITE ON THIS PAGE ANSWER IN THE SPACES PROVIDED



0 8 This question is about carbon and its compounds.

Fullerenes are molecules of carbon atoms.

The first fullerene to be discovered was Buckminsterfullerene (C₆₀).

0 8. 1 What shape is a Buckminsterfullerene molecule?

[1 mark]


0 8. 2 Give **one** use of a fullerene.

[1 mark]

Propanone is a compound of carbon, hydrogen and oxygen.

Figure 13 shows the dot and cross diagram for a propanone molecule.

Figure 13






0 8.3	Complete Figure 14 to show a propanone molecule.
	Use a line to represent each single bond.
	Use Figure 13.
	[1 mark]
	Figure 14
0 8 . 4	— C — U O O Determine the molecular formula of propanone.
	Use Figure 13 .
	[1 mark]
	Molecular formula =
0 8 . 5	Propanone is a liquid with a low boiling point.
	Why does propanone have a low boiling point? [1 mark]
	Tick (✓) one box.
	The covalent bonds are strong.
	The covalent bonds are weak.
	The intermolecular forces are strong.
	The intermolecular forces are weak.





0 8.6 Figure 15 represents the structure of graphite.

Figure 15



Explain why graphite is:

- a good electrical conductor
- soft and slippery.

You should answer in terms of structure and bonding.

[6 marks]

11



Do not write outside the Turn over for the next question DO NOT WRITE ON THIS PAGE ANSWER IN THE SPACES PROVIDED



0 9			structure and the periodic table.		
0 9.1		eaning of 'isotopes	s'. of subatomic particles.		[2 marks]
0 9.2	Table 8 sho isotopes of		nbers and percentage abundance Table 8	es of the	
			Tuble 0		
		Mass number	Percentage abundance (%)		
		69	60		
		71	40		
		ne relative atomic	mass ($A_{ m r}$) of gallium.		
	·		•		[2 marks]
	_				
		Relative atomi	c mass (1 decimal place) =		



	Gallium (Ga) is in Group 3 of the modern periodic table.	outside box
0 9.3	Give the numbers of electrons and neutrons in an atom of the isotope $^{69}_{31}$ Ga [2 marks]	
	Number of electrons	
	Number of neutrons	
0 9 . 4	What is the most likely formula of a gallium ion? [1 mark] Tick (✓) one box.	
	Ga⁺	
	Ga ⁻	
	Ga ³⁺	
	Ga ³⁻	
0 9 . 5	Gallium was discovered six years after Mendeleev published his periodic table.	
	Give two reasons why the discovery of gallium helped Mendeleev's periodic table to become accepted. [2 marks]	
	1	
	2	9





1 0	This question is about the extraction of metals.
	Element $\bf R$ is extracted from its oxide by reduction with hydrogen. The equation for the reaction is: $ 3H_2 + {\bf R}O_3 \to {\bf R}+ 3H_2O $
10.1	The sum of the relative formula masses (M_r) of the reactants (3 H ₂ + RO ₃) is 150 Calculate the relative atomic mass (A_r) of R. Relative atomic masses (A_r): H = 1 O = 16 [2 marks]
	Relative atomic mass (<i>A</i> _r) of R =
1 0.2	Identify element R. You should use: • your answer to question 10.1 • the periodic table. [1 mark]
	Identity of R =

1 0 . 3	Carbon is used to extract tin (Sn) from tin oxide (SnO ₂).	
	The equation for the reaction is:	
	$SnO_2 + C \rightarrow Sn + CO_2$	
	Calculate the percentage atom economy for extracting tin in this reaction.	
	Relative atomic masses (A_r): $C = 12$ $O = 16$ $Sn = 119$	[3 marks]
	Percentage atom economy =	%

Question 10 continues on the next page



1 0.4 Tungsten (W) is a metal.

Tungsten is extracted from tungsten oxide (WO₃).

All other solid products from the extraction method must be separated from the tungsten.

Table 9 shows information about three possible methods to extract tungsten from tungsten oxide.

Table 9


Method	Reactant	Relative cost of reactant	Products
			Tungsten solid
1	Carbon	Low	Carbon dioxide gas
			Tungsten carbide solid
2	2 Hydrogon High		Tungsten solid
	Hydrogen	High	Water vapour
3	Iron	Low	Tungsten solid
3			Iron oxide solid

Evaluate the three possible methods for extracting tungsten from tungsten oxide. [4 marks	s]
	_
	_
	_

END OF QUESTIONS



10





Question number	Additional page, if required. Write the question numbers in the left-hand margin.



Question number	Additional page, if required. Write the question numbers in the left-hand margin.



Question number	Additional page, if required. Write the question numbers in the left-hand margin.
	Copyright information
	For confidentiality purposes, all acknowledgements of third-party copyright material are published in a separate booklet. This booklet is published after each live examination series and is available for free download from www.aqa.org.uk.
	Permission to reproduce all copyright material has been applied for. In some cases, efforts to contact copyright-holders may have been unsuccessful and AQA will be happy to rectify any omissions of acknowledgements. If you have any queries please contact the Copyright Team.
	Copyright © 2021 AQA and its licensors. All rights reserved.



