

GCE

Mathematics

Unit **4722**: Core Mathematics 2

Advanced Subsidiary GCE

Mark Scheme for June 2017

OCR (Oxford Cambridge and RSA) is a leading UK awarding body, providing a wide range of qualifications to meet the needs of candidates of all ages and abilities. OCR qualifications include AS/A Levels, Diplomas, GCSEs, Cambridge Nationals, Cambridge Technicals, Functional Skills, Key Skills, Entry Level qualifications, NVQs and vocational qualifications in areas such as IT, business, languages, teaching/training, administration and secretarial skills.

It is also responsible for developing new specifications to meet national requirements and the needs of students and teachers. OCR is a not-for-profit organisation; any surplus made is invested back into the establishment to help towards the development of qualifications and support, which keep pace with the changing needs of today's society.

This mark scheme is published as an aid to teachers and students, to indicate the requirements of the examination. It shows the basis on which marks were awarded by examiners. It does not indicate the details of the discussions which took place at an examiners' meeting before marking commenced.

All examiners are instructed that alternative correct answers and unexpected approaches in candidates' scripts must be given marks that fairly reflect the relevant knowledge and skills demonstrated.

Mark schemes should be read in conjunction with the published question papers and the report on the examination.

OCR will not enter into any discussion or correspondence in connection with this mark scheme.

© OCR 2017

Annotations and abbreviations

Annotation in scoris	Meaning
✓ and ✖	
BOD	Benefit of doubt
FT	Follow through
ISW	Ignore subsequent working
M0, M1	Method mark awarded 0, 1
A0, A1	Accuracy mark awarded 0, 1
B0, B1	Independent mark awarded 0, 1
^	Omission sign
MR	Misread
NGE	Not good enough
BP	Blank page
Highlighting	
Other abbreviations in mark scheme	Meaning
E1	Mark for explaining
U1	Mark for correct units
G1	Mark for a correct feature on a graph
M1 dep*	Method mark dependent on a previous mark, indicated by *
cao	Correct answer only
oe	Or equivalent
rot	Rounded or truncated
soi	Seen or implied
www	Without wrong working
cwo	Correct working only

Subject-specific Marking Instructions for GCE Mathematics Pure strand

- a Annotations should be used whenever appropriate during your marking.

The A, M and B annotations must be used on your standardisation scripts for responses that are not awarded either 0 or full marks. It is vital that you annotate standardisation scripts fully to show how the marks have been awarded.

For subsequent marking you must make it clear how you have arrived at the mark you have awarded

- b An element of professional judgement is required in the marking of any written paper. Remember that the mark scheme is designed to assist in marking incorrect solutions. Correct *solutions* leading to correct answers are awarded full marks but work must not be judged on the answer alone, and answers that are given in the question, especially, must be validly obtained; key steps in the working must always be looked at and anything unfamiliar must be investigated thoroughly.

Correct but unfamiliar or unexpected methods are often signalled by a correct result following an *apparently* incorrect method. Such work must be carefully assessed. When a candidate adopts a method which does not correspond to the mark scheme, award marks according to the spirit of the basic scheme; if you are in any doubt whatsoever (especially if several marks or candidates are involved) you should contact your Team Leader.

- c The following types of marks are available.

M

A suitable method has been selected and *applied* in a manner which shows that the method is essentially understood. Method marks are not usually lost for numerical errors, algebraic slips or errors in units. However, it is not usually sufficient for a candidate just to indicate an intention of using some method or just to quote a formula; the formula or idea must be applied to the specific problem in hand, eg by substituting the relevant quantities into the formula. In some cases the nature of the errors allowed for the award of an M mark may be specified.

A

Accuracy mark, awarded for a correct answer or intermediate step correctly obtained. Accuracy marks cannot be given unless the associated Method mark is earned (or implied). Therefore M0 A1 cannot ever be awarded.

B

Mark for a correct result or statement independent of Method marks.

E

A given result is to be established or a result has to be explained. This usually requires more working or explanation than the establishment of an unknown result.

Unless otherwise indicated, marks once gained cannot subsequently be lost, eg wrong working following a correct form of answer is ignored. Sometimes this is reinforced in the mark scheme by the abbreviation isw. However, this would not apply to a case where a candidate passes through the correct answer as part of a wrong argument.

- d When a part of a question has two or more 'method' steps, the M marks are in principle independent unless the scheme specifically says otherwise; and similarly where there are several B marks allocated. (The notation 'dep *' is used to indicate that a particular mark is dependent on an earlier, asterisked, mark in the scheme.) Of course, in practice it may happen that when a candidate has once gone wrong in a part of a question, the work from there on is worthless so that no more marks can sensibly be given. On the other hand, when two or more steps are successfully run together by the candidate, the earlier marks are implied and full credit must be given.
- e The abbreviation ft implies that the A or B mark indicated is allowed for work correctly following on from previously incorrect results. Otherwise, A and B marks are given for correct work only — differences in notation are of course permitted. A (accuracy) marks are not given for answers obtained from incorrect working. When A or B marks are awarded for work at an intermediate stage of a solution, there may be various alternatives that are equally acceptable. In such cases, exactly what is acceptable will be detailed in the mark scheme rationale. If this is not the case please consult your Team Leader.

Sometimes the answer to one part of a question is used in a later part of the same question. In this case, A marks will often be 'follow through'. In such cases you must ensure that you refer back to the answer of the previous part question even if this is not shown within the image zone. You may find it easier to mark follow through questions candidate-by-candidate rather than question-by-question.

- f Wrong or missing units in an answer should not lead to the loss of a mark unless the scheme specifically indicates otherwise. Candidates are expected to give numerical answers to an appropriate degree of accuracy, with 3 significant figures often being the norm. Small variations in the degree of accuracy to which an answer is given (e.g. 2 or 4 significant figures where 3 is expected) should not normally be penalised, while answers which are grossly over- or under-specified should normally result in the loss of a mark. The situation regarding any particular cases where the accuracy of the answer may be a marking issue should be detailed in the mark scheme rationale. If in doubt, contact your Team Leader.
- g Rules for replaced work

If a candidate attempts a question more than once, and indicates which attempt he/she wishes to be marked, then examiners should do as the candidate requests.

If there are two or more attempts at a question which have not been crossed out, examiners should mark what appears to be the last (complete) attempt and ignore the others.

NB Follow these maths-specific instructions rather than those in the assessor handbook.

- h For a *genuine* misreading (of numbers or symbols) which is such that the object and the difficulty of the question remain unaltered, mark according to the scheme but following through from the candidate's data. A penalty is then applied; 1 mark is generally appropriate, though this may differ for some units. This is achieved by withholding one A or B mark in the question.

Note that a miscopy of the candidate's own working is not a misread but an accuracy error.

Question		Answer	Marks	Guidance	
1	(i)	$(2\sqrt{7})^2 = x^2 + (x + 2)^2 - 2x(x + 2)\cos 60$ $x^2 + 2x - 24 = 0$ $(x + 6)(x - 4) = 0$ $x = 4$	M1	Attempt use of correct cosine rule	Must be attempt to use correct rule but allow BOD on lack of brackets eg $2\sqrt{7}^2$ not $(2\sqrt{7})^2$, even if subsequently 14, and the same for the terms involving x Allow omission of a square sign when substituting as long as correct formula has been seen No need to evaluate $\cos 60$ for M1 Evaluating in radian mode (-0.952) still can get M1 as long as $\cos 60$ seen first
			A1	Obtain correct 3 term quadratic	Must be simplified to three terms but not necessarily all on one side of the equation
			M1	Attempt to solve 3 term quadratic equation	See additional guidance for valid methods
			A1	Obtain $x = 4$ only	Must be from a correct solution of a correct quadratic, though only the positive root may ever be seen Could draw attention to required root by giving both answers and then eg underlining $x = 4$ A0 if $x = -6$ still present If the other root is stated, before being discarded, it must have been $x = -6$
			[4]		
	(ii)	$\frac{1}{2} \times 4 \times 6 \times \sin 60$ $= 6\sqrt{3}$	M1	Attempt area of the triangle, using their x	Must be using correct formula, including $\frac{1}{2}$ Allow equiv methods, such as $\frac{1}{2}bh$ as long as valid attempt at b and h Must be using a positive, numerical, value of x from (i)
			A1	Obtain $6\sqrt{3}$	Must be given as simplified surd No ISW if then given as decimal, unless the exact value is indicated as the final answer (underlined etc)
			[2]		

Question		Answer	Marks	Guidance
2	(i)	$0.5 \times 0.2 \{ \cos 0 + \cos 0.8 + 2(\cos 0.2 + \cos 0.4 + \cos 0.6) \}$ $= 0.715$	<p>B1</p> <p>M1*</p> <p>M1d*</p> <p>A1</p> <p>[4]</p>	<p>State the 5 correct y-values, and no others</p> <p>Attempt to find area between $x = 0$ and $x = 0.8$, using $k\{y_0 + y_n + 2(y_1 + \dots + y_{n-1})\}$</p> <p>Use $k = 0.5 \times 0.2$ so i</p> <p>Obtain 0.715, or better</p> <p>Working in degrees: B1 if exact values seen (ie $\cos 0.2$ etc), but B0 if straight into decimals M1 M1 is then possible as long as it is clear where each value is being placed</p> <p>$0.5 \times 0.2 \{1.00 + 1.00 + 2(1.00 + 1.00 + 1.00)\} = 0.800$ will be 0/4 unless more detail shown</p>

Question	Answer	Marks	Guidance
(ii)	<p>Graph of $y = \cos x$, with 4 trapezia drawn</p> <p>Tops of the trapezia are below the curve</p>	<p>B1</p> <p>B1</p> <p>[2]</p>	<p>Correct $y = \cos x$ graph, with exactly 4 trapezia of roughly equal width</p> <p>Any valid explanation</p> <p>Trapezia must be plausibly $[0, 0.8]$, allow BOD as long as final trapezium ends before $\frac{\pi}{2}$ Curve may be shown beyond $x = 0.8$, but B0 if clearly of the incorrect shape beyond $x = 0.8$ No need for scale on either axis Exactly four trapezia must be shown, of roughly equal widths, with top vertices on the curve.</p> <p>Not dependent on previous B1 Must refer to the tops of the trapezia so B0 for ‘trapezia are below curve’ (ie ‘top’ not used) Allow ‘trapezium’ rather than ‘trapezia’ Concave / convex is B0 B0 if comparing to exact area B1 for decreasing gradient (but B0 for decreasing curve) Candidates could also use their diagram as part of their explanation – as long as there was an intention to draw trapezia then they are eligible for the second B1 even if B0 for the diagram. This could include a single trapezium (even if labelled $0 - 0.8$), several trapezia whose tops are collinear, an incorrect $y = \cos x$ graph (including $y = \sin x$) and similar. Use of rectangles to support their explanation however is B0. They could shade gaps on their diagram but some text also required B0 for ‘some area not calculated’ unless clear which area - could be described or shaded ISW any irrelevant comments, but B0 if contradictory comments</p>

Question		Answer	Marks	Guidance
3	(i)	$1 + 4x + 7x^2 + 7x^3$	<p>B1</p> <p>M1</p> <p>A1</p> <p>A1</p> <p>[4]</p>	<p>Obtain $1 + 4x$</p> <p>Attempt at least one more term - product of correct binomial coeff and power of $\frac{1}{2}x$</p> <p>Obtain $7x^2$</p> <p>Obtain $7x^3$</p> <p>Must be 1, not 1^8 Must be $4x$ not unsimplified equiv Allow separate terms not linked by '+' eg 1, $4x$</p> <p>Powers of $\frac{1}{2}x$ must be consistent with the binomial coeff being used Binomial coeff must be numerical, so 8C_2 is not yet sufficient Allow M1 if powers only applied to x and not $\frac{1}{2}$</p> <p>Coeff must be simplified As part of sum, or part of list</p> <p>Coeff must be simplified As part of sum, or part of list</p> <p>ISW any attempt to 'simplify'</p> <p>If expanding brackets: Mark as above, but must consider all 8 brackets for the M mark (allow irrelevant terms to be discarded)</p>
	(ii)	$4(y + y^2) + 7(y + y^2)^2$ Hence coefficient of y^2 is 11	<p>M1</p> <p>A1FT</p> <p>[2]</p>	<p>Attempt to use $x = y + y^2$ in their expansion from (i)</p> <p>Obtain coefficient of 11 (or $11y^2$), FT their (i)</p> <p>Replace x with $y + y^2$ in both relevant terms and attempt expansion, including relevant numerical coeffs from (i) Allow M1 if using their attempt at a 'simplified' expansion Could instead attempt a new expansion - must use correct binomial coeffs and powers of $\frac{1}{2}(y + y^2)$</p> <p>Ignore terms involving powers other than y^2 The FT is on their $1 + 4x + 7x^2 + 7x^3$, not a multiple of this as a result of any attempt to 'simplify'</p>

Question	Answer	Marks	Guidance
4	$\frac{dy}{dx} = 5x^{\frac{3}{2}} - 10x$ $y = 2x^{\frac{5}{2}} - 5x^2 + c$ $11 = 64 - 80 + c \Rightarrow c = 27$ $y = 2x^{\frac{5}{2}} - 5x^2 + 27$	<p>B1</p> <p>M1</p> <p>A1</p> <p>A1</p> <p>M1</p> <p>A1</p> <p>[6]</p>	<p>Expand bracket to obtain correct expression</p> <p>Attempt integration</p> <p>Obtain at least one correct term</p> <p>Obtain fully correct expression (allow no +c)</p> <p>Attempt to find c, using (4, 11)</p> <p>Obtain fully correct equation, including $y = \dots$</p> <p>Each term must be of form kx^n, so $5x\sqrt{x}$ is not sufficient</p> <p>M0 if bracket not expanded first M1 can still be gained for integrating an incorrect expansion as long as there are two terms For an 'integration attempt' there must be an increase in power by 1 for both terms</p> <p>Allow unsimplified coeffs</p> <p>Allow unsimplified coeffs</p> <p>There must have been a clear attempt at integration, but can follow M0 eg if the bracket was not expanded first Need to get as far as actually attempting c M1 could be implied by eg $11 = 64 - 80$ followed by an attempt to include a constant to balance the equation M0 if no $+ c$ seen or implied M0 for $x = 11, y = 4$ Allow a slip when substituting, as long as it is clear that use of $x = 4, y = 11$ is intended</p> <p>Coefficients now need to be simplified Must be an equation ie $y = \dots$, so A0 for 'f(x) = ...' or 'equation = ...' ISW an incorrect attempt to further simplify the equation</p> <p>NB this question can also be done using integration by parts – 4/4 for correct integral (but no partial credit) and then M1A1 as per MS</p>

Question	Answer	Marks	Guidance	
5	$r\theta = 6$ $\frac{1}{2}r^2\theta = 24$ $\frac{1}{2}r \times 6 = 24$ $r = 8, \theta = 0.75$ segment area = $24 - \frac{1}{2} \times 8^2 \times \sin 0.75$ $= 2.19$	B1* B1* M1d* A1 M1 A1 [6]	State $r\theta = 6$ State $\frac{1}{2}r^2\theta = 24$ Attempt to solve simultaneously to find r or θ Obtain $r = 8, \theta = 0.75$ (aef) Attempt area of segment Obtain 2.19, or better [6]	Or exact equiv from using a fraction of the circle Or exact equiv from using a fraction of the circle Allow B1 for $\frac{1}{2} r \times \text{arc} = 24$ Stating both $\frac{1}{2}r^2\theta = 24$ and $\frac{1}{2}r^2\sin\theta = 24$ is B0 unless only the correct equation is subsequently used B1 B1 can be implied by a correct equation in a single variable As far as attempting r or θ , using a valid method (but allow slips) Must be using the two correct equations in r and θ Both values required 24 – area of triangle, using $\frac{1}{2}r^2\sin\theta$ or equiv Allow if evaluated in degree mode (gives 23.58) Allow M1 for attempting $\frac{1}{2}r^2(\theta - \sin\theta)$ with their r and θ , even if this does not give area of sector as 24 Allow final answer in range [2.187, 2.188] if > 3sf Could use variables other than r and θ Alt method for working in degrees B1 - state $\frac{\theta}{360} \times 2\pi r = 6$ B1 - state $\frac{\theta}{360} \times \pi r^2 = 24$ M1 - attempt to solve simultaneously A1 - obtain $r = 8, \theta = 43.0^\circ$ or better (42.97...) M1 - attempt area of segment NB using $\frac{1}{2}r^2(\theta - \sin\theta)$ with θ in degrees is M0 as incorrect attempt at area of sector A1 - obtain 2.19 or better

Question	Answer	Marks	Guidance	
6	$\int (11 - x - 2x^2) dx = 11x - \frac{1}{2}x^2 - \frac{2}{3}x^3$ $\int 8x^{-3} dx = -4x^{-2}$ $(22 - 2 - \frac{16}{3}) - (11 - \frac{1}{2} - \frac{2}{3}) = \frac{29}{6}$ $(-1) - (-4) = 3$ $\frac{29}{6} - 3 = \frac{11}{6}$	<p>M1</p> <p>A1</p> <p>M1</p> <p>A1</p> <p>M1</p> <p>M1</p> <p>A1</p> <p>[7]</p>	<p>Attempt integration of $11 - x - 2x^2$</p> <p>Obtain $11x - \frac{1}{2}x^2 - \frac{2}{3}x^3$</p> <p>Attempt integration of $8x^{-3}$</p> <p>Obtain $-4x^{-2}$</p> <p>Use limits of $x = 1, 2$</p> <p>Attempt correct method to find shaded area (at any point)</p> <p>Obtain $\frac{11}{6}$, or exact equiv</p>	<p>Increase in power by 1 for at least 2 terms</p> <p>Obtain correct integral</p> <p>Integrate to kx^{-2}</p> <p>Allow unsimplified coeff</p> <p>In both integrals Must follow clear attempt at integration Must be $F(2) - F(1)$ ie correct order and subtraction</p> <p>M0 if incorrect order of subtraction, even if $\frac{11}{6}$ subsequently appears as final answer M1 can follow M0 for use of limits</p> <p>A0 for decimal answer unless clearly a recurring decimal (but not eg 1.833...) ISW if $\frac{11}{6}$ seen but then followed by eg 1.83</p> <p>Answer only is 0/7 - need to see evidence of integration, but use of limits does not need to be explicit</p> <p>Alternative MS for subtracting first: M1 - attempt subtraction in correct order M1 - attempt integration of $\pm(11 - x - 2x^2)$ A1 - obtain $\pm(11x - \frac{1}{2}x^2 - \frac{2}{3}x^3)$, signs must be consistent with their subtraction M1 - attempt integration of $\pm 8x^{-3}$ A1 - obtain $\mp 4x^{-2}$, sign must be consistent with their subtraction M1 - correct use of limits in entire integral A1 - obtain $\frac{11}{6}$</p>

Question			Answer	Marks	Guidance
					<p>Ignore sight of $11 - x - 2x^2 = 8x^{-3}$ prior to subtraction occurring</p> <p>Adding functions prior to integration will get max of 5 marks - M0M1A1M1A1M1A0 (Alt MS) – to give same credit as integrating separately, using limits and then adding</p> <p>Multiplying through by x^3 prior to integration can get M1 for use of limits, and possibly M1 if subtraction happens before multiplying through</p>
7	(a)	$(x + 1)\log_3 = 500\log_2$ $x + 1 = 315.46$ $x = 314$	<p>M1</p> <p>A1</p> <p>M1</p> <p>A1</p> <p>[4]</p>	<p>Introduce logs and drop power(s)</p> <p>Obtain correct linear equation</p> <p>Attempt to find x</p> <p>Obtain 314, or better</p>	<p>Any base (or no explicit base) as long as consistent If using logs to any base other than 2 or 3, then both powers must be dropped Allow BOD if $x + 1$ is not in brackets</p> <p>aef eg $x + 1 = 500\log_3 2$ Brackets must now be explicit, or implied by a later correct equation</p> <p>Correct order of operations and correct operations, so M0 for eg $x = 500\log_3 2 + 1$</p> <p>Allow answer in range [314.4, 314.5] if > 3sf ISW subsequent incorrect rounding once more accurate answer has been seen</p> <p>Answer only, or T&I, is 0/4 Writing 2 as $3^{0.6309}$ with no evidence of use of logs to find the index is 0/4</p>

Question		Answer	Marks	Guidance	
	(b)	(i)			
		(ii)			

Question		Answer	Marks	Guidance	
8	(a)	$a + 6d = 2(a + 4d)$ $a = -2d$ $\frac{7}{2}(2a + 6d) = 84$ $\frac{7}{2}(2a - 3a) = 84$ $a = -24$	M1	Attempt $u_7 = 2u_5$	Using correct $u_n = a + (n - 1)d$
			A1	Obtain $a = -2d$ or equiv	Obtain correct simplified equation, with like terms combined
			B1	State $\frac{7}{2}(2a + 6d) = 84$	Or equiv, including unsimplified
			M1	Attempt to solve simultaneously	As far as attempting a or d Must be solving two equations in a and d , from attempt at $u_7 = 2u_5$ and attempt at S_7 (but could be from incorrect formulae eg $u_n = a + nd$)
			A1	Obtain $a = -24$	If d is also given then it must be correct ($d = 12$)
			[5]		Could use variables other than a and d

Question	Answer	Marks	Guidance	
(b)	$r^2 = 2 \text{ hence } r = \sqrt{2}$ $\frac{a(1-\sqrt{2}^7)}{1-\sqrt{2}} = 254$ $a = \frac{254(1-\sqrt{2})}{1-8\sqrt{2}}$ $a = \frac{254(1-\sqrt{2})(1+8\sqrt{2})}{(1-8\sqrt{2})(1+8\sqrt{2})}$ $a = \frac{254(-15+7\sqrt{2})}{-127}$ $a = 30 - 14\sqrt{2}$	<p>B1</p> <p>M1</p> <p>A1</p> <p>B1</p> <p>M1</p> <p>A1</p> <p>[6]</p>	<p>State $r = \sqrt{2}$ www</p> <p>Attempt $S_7 = 254$</p> <p>Rearrange to obtain correct numerical expression for a aef</p> <p>Use $(\sqrt{2})^7 = 8\sqrt{2}$ soi</p> <p>Attempt to rationalise denominator</p> <p>Obtain correct value in surd form</p>	<p>B0 if from $ar^7 = 2ar^5$ (but then allow all of the remaining marks) Allow decimal value (1.41) Allow B1 for $r = \pm \sqrt{2}$</p> <p>Must be correct formula, using their numerical r, which could be exact or a decimal value Must also equate to 254</p> <p>Must be in an exact form, but could involve $(\sqrt{2})^7$ or $\sqrt{128}$ rather than $8\sqrt{2}$ Ignore second value for a from using $r = -\sqrt{2}$</p> <p>Equation may no longer be fully correct</p> <p>Must be using $r = \sqrt{2}$ only Must be explicit evidence of rationalising Could use $(1 + (\sqrt{2})^7)$ or $(1 + \sqrt{128})$ Allow M1 if denominator now incorrect, as long as of form $\pm (1 - k\sqrt{2})$ or equiv M0 if rationalising $1 - \sqrt{2}$ only (ie before making a the subject)</p> <p>Allow any exact answer in form $p + q\sqrt{r}$ A0 if additional answer from using $r = -\sqrt{2}$ A0 if final answer results from subsequent attempt to simplify eg $a = 15 - 7\sqrt{2}$ (ie no ISW)</p> <p>Could use variables other than a and r</p> <p>If $a = 30 - 14\sqrt{2}$ obtained, but no evidence of dealing with $(\sqrt{2})^7$ or rationalising denominator then maximum of B1 M1 A1 ie 3 marks (as the given form has not been 'shown')</p>

Question		Answer	Marks	Guidance	
9	(i)	$f(1/2) = 1/2 + 9/2 - 5 = 0$ $f(x) = (2x - 1)(2x^2 + x + 5)$	B1	Confirm $f(1/2) = 0$, with detail shown $4(1/2)^3 + 9(1/2) - 5 = 0$ is sufficient B0 for just $f(1/2) = 0$ No conclusion needed If using division to justify then must draw attention to the zero remainder	
			M1	Attempt complete division or equiv Must be dividing by $(2x - 1)$ Must be complete method - ie all 3 terms attempted Long division - must subtract lower line (allow one slip) Inspection - expansion must give at least three correct terms of the cubic Coefficient matching - must be valid attempt at all coeffs of quadratic, considering all relevant terms each time Synthetic division - must be using 0.5 (not -0.5) and adding within each column (allow one slip); expect to see $ \begin{array}{r rrrr} 0.5 & 4 & 0 & 9 & -5 \\ & & 2 & 1 & \\ \hline & 4 & 2 & 10 & \end{array} $	
			A1	Obtain correct quotient	Allow $4x^2 + 2x + 10$ from dividing by $x - 1/2$
			A1	Obtain $(2x - 1)(2x^2 + x + 5)$	Must be written as a product Allow $(x - 1/2)(4x^2 + 2x + 10)$ ISW any attempt to write as 3 linear factors, or to find roots
			[4]		

Question		Answer	Marks	Guidance
(ii)	(a)	$4 \sin 2\theta \cos 2\theta + \frac{5}{\cos 2\theta} = \frac{13 \sin 2\theta}{\cos 2\theta}$ $4 \sin 2\theta \cos^2 2\theta + 5 = 13 \sin 2\theta$ $4 \sin 2\theta(1 - \sin^2 2\theta) + 5 = 13 \sin 2\theta$ $4 \sin 2\theta - 4 \sin^3 2\theta + 5 = 13 \sin 2\theta$ $4 \sin^3 2\theta + 9 \sin 2\theta - 5 = 0$	B1 B1 B1 B1 [4]	Use $\tan 2\theta = \frac{\sin 2\theta}{\cos 2\theta}$ or $\tan 2\theta \cos 2\theta = \sin 2\theta$ Correct method to remove fraction(s) Use $\cos^2 2\theta = 1 - \sin^2 2\theta$ Obtain correct equation, from correct working Must be explicit, and correct notation when used Allow even if errors elsewhere in equation Any correct equation seen no longer containing fractions (allow recovery from a slip in notation) Must be explicit, and correct notation when used Allow even if errors elsewhere in equation Must be correct notation throughout Dependent on B1 B1 B1 awarded NB - must annotate answer space at top of pg12
	(b)	$(2 \sin 2\theta - 1)(2 \sin^2 2\theta + \sin 2\theta + 5) = \sin 2\theta = \frac{1}{2}$ $2\theta = \frac{1}{6}\pi, \frac{5}{6}\pi, \frac{13}{6}\pi, \frac{17}{6}\pi$ $\theta = \frac{1}{12}\pi, \frac{5}{12}\pi, \frac{13}{12}\pi, \frac{17}{12}\pi$	B1 M1 A1 A1 [4]	State that $\sin 2\theta = \frac{1}{2}$ oe Attempt to solve $\sin 2\theta = \pm \frac{1}{2}$ to find at least one root Obtain at least 2 correct roots Obtain 4 correct roots Could just be stated, or implied by later method Correct order of operations ie $\frac{1}{2} (\sin^{-1} \frac{1}{2})$ Allow M1 if angle(s) found in degrees ($15^\circ, 75^\circ$ etc) Must be in radians, and given in an exact form Allow recurring decimals, or mixed numbers Must be in radians, and given in an exact form Allow recurring decimals, or mixed numbers ISW any angles that come from an incorrect quadratic quotient, or an incorrect attempt to find the roots of the quadratic quotient A0 if any extras in range $[0, 2\pi]$ that are not clearly from their quadratic roots

APPENDIX 1**Guidance for marking C2****Accuracy**

Allow answers to 3sf or better, unless an integer is specified or clearly required.

Answers to 2 sf are penalised, unless stated otherwise in the mark scheme.

‘3sf’ is sometimes explicitly specified in a question - this is telling candidates that a decimal is required rather than an exact answer eg in logs, and more than 3sf should not be penalised unless stated in mark scheme.

If more than 3sf are given, allow the marks for an answer that falls within the guidance given in the mark scheme, with no obvious errors.

Extra solutions

Candidates will usually be penalised if any extra, incorrect, solutions are given. However, in trigonometry questions, only look at solutions in the given range and ignore any others, correct or incorrect.

Solving equations

With simultaneous equations, the method mark is given for eliminating one variable. Any valid method is allowed, such as balancing or substitution.

Solving quadratic equations

Factorising - candidates must get as far as factorising into two brackets which, on expansion, would give the correct coefficient of x^2 and at least one of the other two coefficients. This method is only credited if it is possible to factorise the quadratic – if the roots are surds then candidates are expected to use either the quadratic formula or complete the square.

Completing the square - candidates must get as far as $(x + p) = \pm \sqrt{q}$, with reasonable attempts at p and q .

Using the formula - candidates need to substitute values into the formula, with some attempt at evaluation (eg calculating $4ac$). The correct formula must be seen, either algebraic or numerical. If the algebraic formula is quoted then candidates are allowed to make one slip when substituting their values. The division line must extend under the entire numerator (seen or implied by later working). Condone not dividing by $2a$ as long as it has been seen earlier.

Solutions with no method shown

If a correct equation is seen, then the correct answers will imply that the method is correct – unless specified otherwise in the mark scheme, all answers to the equation must be given. So, if solving a quadratic, only the correct two roots will imply a correct method (NB on this paper the MS does identify an exception to this rule on Q1).

If an incorrect equation is seen, and no supporting method for solving it is shown, then examiners must not try to deduce the method used from the solutions provided.

OCR (Oxford Cambridge and RSA Examinations)
1 Hills Road
Cambridge
CB1 2EU

OCR Customer Contact Centre

Education and Learning

Telephone: 01223 553998

Facsimile: 01223 552627

Email: general.qualifications@ocr.org.uk

www.ocr.org.uk

For staff training purposes and as part of our quality assurance programme your call may be recorded or monitored

Oxford Cambridge and RSA Examinations
is a Company Limited by Guarantee
Registered in England
Registered Office; 1 Hills Road, Cambridge, CB1 2EU
Registered Company Number: 3484466
OCR is an exempt Charity

OCR (Oxford Cambridge and RSA Examinations)
Head office
Telephone: 01223 552552
Facsimile: 01223 552553

© OCR 2017

