


4753

Mark Scheme

June 2006

**Mark Scheme 4753**  
**June 2006**

<p><b>1</b>      <math> 3x - 2  = x</math>  <math>\Rightarrow 3x - 2 = x \Rightarrow 2x = 2 \Rightarrow x = 1</math>      or      <math>2 - 3x = x \Rightarrow 2 = 4x \Rightarrow x = \frac{1}{2}</math>      or      <math>(3x - 2)^2 = x^2</math>  <math>\Rightarrow 8x^2 - 12x + 4 = 0 \Rightarrow 2x^2 - 3x + 1 = 0</math>  <math>\Rightarrow (x - 1)(2x - 1) = 0,</math>  <math>\Rightarrow x = 1, \frac{1}{2}</math></p>	B1 M1 A1 M1 A1 A1 [3]	$x = 1$ solving correct quadratic
<p><b>2</b>      let <math>u = x</math>, <math>dv/dx = \sin 2x \Rightarrow v = -\frac{1}{2}\cos 2x</math>  <math>\Rightarrow \int_0^{\pi/6} x \sin 2x dx = \left[ x \cdot -\frac{1}{2} \cos 2x \right]_0^{\pi/6} + \int_0^{\pi/6} \frac{1}{2} \cos 2x \cdot 1 dx</math>  <math>= \frac{\pi}{6} \cdot -\frac{1}{2} \cos \frac{\pi}{3} - 0 + \left[ \frac{1}{4} \sin 2x \right]_0^{\pi/6}</math>  <math>= -\frac{\pi}{24} + \frac{\sqrt{3}}{8}</math>  <math>= \frac{3\sqrt{3} - \pi}{24} *</math></p>	M1 A1 B1ft M1 B1 E1 [6]	parts with $u = x$ , $dv/dx = \sin 2x$ ... + $\left[ \frac{1}{4} \sin 2x \right]_0^{\pi/6}$ substituting limits $\cos \pi/3 = \frac{1}{2}$ , $\sin \pi/3 = \sqrt{3}/2$ soi www
<p><b>3 (i)</b>      <math>x - 1 = \sin y</math>  <math>\Rightarrow x = 1 + \sin y</math>  <math>\Rightarrow dx/dy = \cos y</math></p> <p><b>(ii)</b> When <math>x = 1.5</math>, <math>y = \arcsin(0.5) = \pi/6</math></p> $\begin{aligned} \frac{dy}{dx} &= \frac{1}{\cos y} \\ &= \frac{1}{\cos \pi/6} \\ &= 2/\sqrt{3} \end{aligned}$	M1 A1 E1 M1 A1 M1 A1 [7]	www condone $30^\circ$ or 0.52 or better or $\frac{dy}{dx} = \frac{1}{\sqrt{1-(x-1)^2}}$ or equivalent, but must be exact
<p><b>4(i)</b>      <math>V = \pi h^2 - \frac{1}{3}\pi h^3</math>  <math>\Rightarrow \frac{dV}{dh} = 2\pi h - \pi h^2</math></p> <p><b>(ii)</b> <math>\frac{dV}{dt} = 0.02</math>  <math>\frac{dV}{dt} = \frac{dV}{dh} \cdot \frac{dh}{dt}</math>  <math>\Rightarrow \frac{dh}{dt} = \frac{0.02}{dV/dh} = \frac{0.02}{2\pi h - \pi h^2}</math></p> <p>When <math>h = 0.4</math>, <math>\Rightarrow \frac{dh}{dt} = \frac{0.02}{0.8\pi - 0.16\pi} = 0.0099 \text{ m/min}</math></p>	M1 A1 B1 M1 M1dep A1cao [6]	expanding brackets (correctly) or product rule oe soi $\frac{dV}{dt} = \frac{dV}{dh} \cdot \frac{dh}{dt}$ oe substituting $h = 0.4$ into their $\frac{dV}{dh}$ and $\frac{dV}{dt} = 0.02$ 0.01 or better or $1/32\pi$

<p><b>5(i)</b></p> $\begin{aligned} a^2 + b^2 &= (2t)^2 + (t^2 - 1)^2 \\ &= 4t^2 + t^4 - 2t^2 + 1 \\ &= t^4 + 2t^2 + 1 \\ &= (t^2 + 1)^2 = c^2 \end{aligned}$ <p><b>(ii)</b> <math>c = \sqrt{(20^2 + 21^2)} = 29</math> For example: <math>2t = 20 \Rightarrow t = 10</math> <math>\Rightarrow t^2 - 1 = 99</math> which is not consistent with 21</p>	M1 M1 E1  B1 M1 E1 [6]	substituting for $a$ , $b$ and $c$ in terms of $t$ Expanding brackets correctly www  Attempt to find $t$ Any valid argument or E2 'none of 20, 21, 29 differ by two'.
<p><b>6 (i)</b></p>  <p><b>(ii)</b> <math>\frac{M}{M_0} = e^{-0.000121 \times 5730} = e^{-0.6933\dots} \approx \frac{1}{2}</math></p> <p><b>(iii)</b> <math>\frac{M}{M_0} = e^{-kT} = \frac{1}{2}</math>  <math>\Rightarrow \ln \frac{1}{2} = -kT</math>  <math>\Rightarrow \ln 2 = kT</math>  <math>\Rightarrow T = \frac{\ln 2}{k} *</math></p> <p><b>(iv)</b> <math>T = \frac{\ln 2}{2.88 \times 10^{-5}} \approx 24000 \text{ years}</math></p>	B1 B1  M1 E1  M1 M1  E1  B1 [8]	Correct shape Passes through $(0, M_0)$  substituting $k = -0.000121$ and $t = 5730$ into equation (or ln eqn) showing that $M \approx \frac{1}{2} M_0$  substituting $M/M_0 = \frac{1}{2}$ into equation (oe) taking ln correctly  24 000 or better

## Section B

7(i) $x = 1$	B1 [1]	
(ii) $\begin{aligned} \frac{dy}{dx} &= \frac{(x-1)2x - (x^2 + 3).1}{(x-1)^2} \\ &= \frac{2x^2 - 2x - x^2 - 3}{(x-1)^2} \\ &= \frac{x^2 - 2x - 3}{(x-1)^2} \end{aligned}$ $\begin{aligned} dy/dx = 0 \text{ when } x^2 - 2x - 3 &= 0 \\ \Rightarrow (x-3)(x+1) &= 0 \\ \Rightarrow x = 3 \text{ or } -1 \\ \text{When } x = 3, y = (9+3)/2 &= 6 \\ \text{So P is } (3, 6) \end{aligned}$	M1  A1    M1 M1 A1 B1ft [6]	Quotient rule  correct expression   their numerator = 0 solving quadratic by any valid method $x = 3$ from correct working $y = 6$
(iii) $\begin{aligned} \text{Area} &= \int_2^3 \frac{x^2 + 3}{x-1} dx \\ u = x - 1 \Rightarrow du/dx &= 1, du = dx \\ \text{When } x = 2, u = 1; \text{ when } x = 3, u = 2 \\ &= \int_1^2 \frac{(u+1)^2 + 3}{u} du \\ &= \int_1^2 \frac{u^2 + 2u + 4}{u} du \\ &= \int_1^2 \left(u + 2 + \frac{4}{u}\right) du * \\ &= \left[\frac{1}{2}u^2 + 2u + 4\ln u\right]_1^2 \\ &= (2 + 4 + 4\ln 2) - (\frac{1}{2} + 2 + 4\ln 1) \\ &= 3\frac{1}{2} + 4\ln 2 \end{aligned}$	M1  B1  B1  E1  B1  M1 A1cao [7]	Correct integral and limits  Limits changed, and substituting $dx = du$  substituting $\frac{(u+1)^2 + 3}{u}$  www  [ $\frac{1}{2}u^2 + 2u + 4\ln u$ ]  substituting correct limits
(iv) $\begin{aligned} e^y &= \frac{x^2 + 3}{x-1} \\ \Rightarrow e^y \frac{dy}{dx} &= \frac{x^2 - 2x - 3}{(x-1)^2} \\ \Rightarrow \frac{dy}{dx} &= e^{-y} \frac{x^2 - 2x - 3}{(x-1)^2} \end{aligned}$  $\begin{aligned} \text{When } x = 2, e^y &= 7 \Rightarrow \\ \Rightarrow dy/dx &= \frac{1}{7} \cdot \frac{4-4-3}{1} = -\frac{3}{7} \end{aligned}$	M1  A1ft   B1 A1cao [4]	$e^y dy/dx = \text{their } f'(x)$ or $xe^y - e^y = x^2 + 3$ $\Rightarrow e^y + xe^y \frac{dy}{dx} - e^y \frac{dy}{dx} = 2x$ $\Rightarrow \frac{dy}{dx} = \frac{2x - e^y}{e^y(x-1)}$ $y = \ln 7 \text{ or } 1.95\dots \text{ or } e^y = 7$ or $\frac{dy}{dx} = \frac{4-7}{7(2-1)} = -\frac{3}{7}$ or $-0.43$ or better

<p><b>8 (i) (A)</b></p> <p><b>(B)</b></p>	B1 B1  M1 A1 [4]	Zeros shown every $\pi/2$ . Correct shape, from $-\pi$ to $\pi$  Translated in $x$ -direction $\pi$ to the left
<p><b>(ii)</b> <math>f'(x) = -\frac{1}{5}e^{-\frac{1}{5}x} \sin x + e^{-\frac{1}{5}x} \cos x</math></p> <p><math>f'(x) = 0</math> when <math>-\frac{1}{5}e^{-\frac{1}{5}x} \sin x + e^{-\frac{1}{5}x} \cos x = 0</math></p> <p><math>\Rightarrow \frac{1}{5}e^{-\frac{1}{5}x}(-\sin x + 5\cos x) = 0</math></p> <p><math>\Rightarrow \sin x = 5\cos x</math></p> <p><math>\Rightarrow \frac{\sin x}{\cos x} = 5</math></p> <p><math>\Rightarrow \tan x = 5^*</math></p> <p><math>\Rightarrow x = 1.37(34\dots)</math></p> <p><math>\Rightarrow y = 0.75</math> or <math>0.74(5\dots)</math></p>	B1 B1  M1 E1 B1 B1 [6]	$e^{-\frac{1}{5}x} \cos x$  $\dots -\frac{1}{5}e^{-\frac{1}{5}x} \sin x$  dividing by $e^{-\frac{1}{5}x}$  www 1.4 or better, must be in radians 0.75 or better
<p><b>(iii)</b> <math>f(x+\pi) = e^{-\frac{1}{5}(x+\pi)} \sin(x+\pi)</math></p> <p><math>= e^{-\frac{1}{5}x} e^{-\frac{1}{5}\pi} \sin(x+\pi)</math></p> <p><math>= -e^{-\frac{1}{5}x} e^{-\frac{1}{5}\pi} \sin x</math></p> <p><math>= -e^{-\frac{1}{5}\pi} f(x)^*</math></p> <p><math>\int_{\pi}^{2\pi} f(x) dx \quad \text{let } u = x - \pi, du = dx</math></p> <p><math>= \int_0^{\pi} f(u+\pi) du</math></p> <p><math>= \int_0^{\pi} -e^{-\frac{1}{5}u} f(u) du</math></p> <p><math>= -e^{-\frac{1}{5}\pi} \int_0^{\pi} f(u) du^*</math></p> <p>Area enclosed between <math>\pi</math> and <math>2\pi</math>  <math>= (-) e^{-\frac{1}{5}\pi} \times \text{area between } 0 \text{ and } \pi.</math></p>	M1 A1 A1 E1  B1 B1dep E1  B1 B1 [8]	$e^{-\frac{1}{5}(x+\pi)} = e^{-\frac{1}{5}x} \cdot e^{-\frac{1}{5}\pi}$  $\sin(x+\pi) = -\sin x$ www  $\int f(u+\pi) du$ limits changed  using above result or repeating work  or multiplied by 0.53 or better