

Write your name here

Surname

Other names

Pearson Edexcel
Level 3 GCE

Centre Number

--	--	--	--	--

Candidate Number

--	--	--	--

Chemistry

Advanced

Paper 2: Advanced Organic and Physical Chemistry

Sample Assessment Materials for first teaching September 2015

Time: 1 hour 45 minutes

Paper Reference

9CH0/02

You must have:

Data Booklet
Scientific calculator, ruler

Total Marks

--

Instructions

- Use **black** ink or ball-point pen.
- **Fill in the boxes** at the top of this page with your name, centre number and candidate number.
- Answer **all** questions.
- Answer the questions in the spaces provided
– *there may be more space than you need.*

Information

- The total mark for this paper is 90.
- The marks for **each** question are shown in brackets
– *use this as a guide as to how much time to spend on each question.*
- You may use a scientific calculator.
- For questions marked with an *, marks will be awarded for your ability to structure your answer logically showing the points that you make are related or follow on from each other where appropriate.
- A Periodic Table is printed on the back cover of this paper.

Advice

- Read each question carefully before you start to answer it.
- Try to answer every question.
- Check your answers if you have time at the end.
- Show all your working in calculations and include units where appropriate.

Turn over ►

S47554A

©2015 Pearson Education Ltd.



PEARSON

Answer ALL questions.

Write your answers in the spaces provided.

Some questions must be answered with a cross .
If you change your mind about an answer, put a line through the box
and then mark your new answer with a cross .


1 This question is about some hydrocarbons.

(a) (i) Which isomeric alkane has the lowest boiling temperature?

(1)

- A** 2,2-dimethylbutane
- B** hexane
- C** 3-methylpentane
- D** 2-methylpentane

(ii) What is the name of this alkane?



(1)

- A** 4-ethylpentane
- B** 2-ethylpentane
- C** 4-methylhexane
- D** 3-methylhexane

(b) Describe how the orbitals from carbon atoms interact to form the bonds in a benzene ring. You may include a labelled diagram to support your answer.

(3)

.....





.....

.....

.....

(Total for Question 1 = 5 marks)

2 The skeletal formulae of four compounds are shown.



(a) Which compound has the molecular formula $C_6H_{14}O$?

(1)

- A
- B
- C
- D

(b) Which compound forms a pale yellow precipitate when reacted with iodine and sodium hydroxide solution?

(1)

A

B

C

D

(c) Which compound forms a silver mirror when reacted with Tollens' reagent?

(1)

A

B

C

D

(Total for Question 2 = 3 marks)

3 This question is about amines.

(a) Amines can behave as bases.

What is meant by the term base?

(1)

.....

.....

.....

.....

(b) Which of these compounds has the highest pH when dissolved in water to form solutions of the same concentration?

(1)

- A** NH_3
- B** CH_3NH_2
- C** $\text{C}_6\text{H}_5\text{NH}_2$
- D** NH_4Cl

- (c) A student wrote a statement about the mechanism of the reaction between a primary chloroalkane and an amine.

The amine acts as a nucleophile due to its unpaired electron. It attacks the slightly positive carbon in the chloroalkane, forming an intermediate. The carbon is slightly positive as it is attached to an electropositive chlorine atom. The intermediate then loses a proton and a chloride ion, forming the product, an N-substituted amide.

Identify and correct **two** of the errors in the statement.

(2)

.....

.....


.....

.....

(Total for Question 3 = 4 marks)

4 This question is about the properties of amino acids and other organic compounds.

(a) A chromatogram of two amino acids, **A** and **B**, is shown.



The R_f value of amino acid A is

(1)

- A 0.51
- B 0.56
- C 0.63
- D 0.67

(b) Amino acid **B** has the structure $\text{H}_2\text{NCH}(\text{CH}_2\text{CH}(\text{CH}_3)_2)\text{COOH}$.

Predict the pH of the solution formed when it dissolves in water and justify your answer.

(2)

.....

.....

.....

.....

(c) Some data about different organic compounds is shown in the table.

Compound	Relative molecular mass	Melting temperature / °C
glycine (an amino acid)	75	233
butan-1-ol	74	-90
pentane	72	-130

Use information from the table to justify the types of intermolecular forces in each of these compounds.

(4)

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(Total for Question 4 = 7 marks)


5 This question is about the chemistry of iodine compounds and hydrocarbons.

(a) Iodine monochloride (ICl) reacts with unsaturated compounds such as sunflower oil.


(i) Which diagram shows the first step in the mechanism of the reaction between iodine monochloride and an unsaturated compound?

(1)


A




B



C



D




(ii) What is the name and type of the mechanism in the overall reaction?

(1)


- A electrophilic substitution
- B nucleophilic substitution
- C electrophilic addition
- D nucleophilic addition

(b) The number of grams of iodine from iodine monochloride that reacts with 100 g of an oil or fat is known as the iodine value and is used to compare levels of unsaturation in different oils and fats.

An excess of iodine monochloride, 11.0 g, was mixed with 6.40 g of sunflower oil. The remaining unreacted iodine monochloride was treated with excess potassium iodide to liberate iodine.



The liberated iodine was titrated with sodium thiosulfate solution of concentration $0.500 \text{ mol dm}^{-3}$. 21.20 cm^3 of this solution was required to completely react with the liberated iodine.



(i) Calculate the amount, in mol, of sodium thiosulfate that reacted with the liberated iodine and hence calculate the amount (in mol) of liberated iodine.

(2)

(ii) Use your answer to (i) and the initial mass of iodine monochloride, 11.0 g, to calculate the iodine value of the oil.

(5)

(c) Sunflower oil can be hydrogenated to form solid fats such as margarine. What is the **most** appropriate catalyst for this reaction?

(1)

- A nickel
- B iron
- C rhodium
- D vanadium(V) oxide

(Total for Question 5 = 10 marks)

BLANK PAGE

6 This question is about the kinetics of chemical reactions.

(a) The rate equation for the reaction between hydrogen and nitrogen monoxide is:

$$\text{rate} = k[\text{H}_2][\text{NO}]^2$$

By what factor does the rate increase when the concentration of hydrogen is tripled and that of nitrogen monoxide is doubled?

(1)

- A 5
- B 6
- C 12
- D 18

(b) The 'initial rates' method is used to investigate the orders of reaction with respect to reactants **A**, **B** and **C**. The table shows the results obtained.

Run	Initial concentration / mol dm ⁻³			Initial rate / mol dm ⁻³ s ⁻¹
	A	B	C	
1	0.32	0.080	0.16	2.4×10^{-3}
2	0.64	0.080	0.16	4.8×10^{-3}
3	0.32	0.16	0.16	9.6×10^{-3}
4	0.64	0.32	0.48	0.23

(i) Deduce the orders with respect to **A** and **B**.

(2)

A

B

(ii) Deduce the order with respect to **C** and justify your answer.

(2)

.....

.....

.....

.....

.....


(iii) Give the rate equation for the reaction.

(1)

(iv) Calculate the rate constant, k , to an appropriate number of significant figures.
Give units for your answer.

(3)

(c) A different reaction, between iodine and propanone, in the presence of hydrogen ions, has the rate equation:



(i) Give the overall order of the reaction.

(1)

.....

.....

.....

(ii) Explain, in terms of collision theory, why increasing the concentration of propanone changes the rate of reaction.

(2)

.....

.....

.....

.....

.....

.....

(d) Catalysts are sometimes used in reactions.

(i) Explain how a catalyst affects the rate of a reaction.

(3)

.....

.....

.....

.....

.....

.....

.....

.....


.....

.....

.....

(ii) Draw and label lines on the axes to show how an increase in temperature affects the number of particles with $E > E_a$.

(2)



(Total for Question 6 = 17 marks)

7 The compound 2-phenylethyl propanoate is a synthetic oil used in some fragrances due to its rose-like aroma.


(a) Combustion analysis of 1.56 g of 2-phenylethyl propanoate, in a laboratory, produced 4.26 g of $\text{CO}_2(\text{g})$ and 1.10 g of $\text{H}_2\text{O}(\text{l})$.

Show that the data is consistent with the molecular formula of 2-phenylethyl propanoate, $\text{C}_{11}\text{H}_{14}\text{O}_2$.

(4)

(b) Nuclear magnetic resonance (NMR) spectroscopy can be used to determine structural formulae.

An incomplete ^1H NMR spectrum for 2-phenylethyl propanoate shows a peak for the hydrogen atoms attached to the benzene ring.




Complete the spectrum for the protons attached to the carbon atoms **p**, **q**, **r** and **s**, giving the relative areas under each peak.


You should use the Data Booklet. Your spectrum does not have to be to scale.

(6)

Predicted spectrum:



(c) A possible synthetic route to prepare 2-phenylethyl propanoate is:



(i) The attacking species in Step 1 can be represented as $^+\text{CH}_2\text{CH}_2\text{OAlCl}_3^-$


What is the mechanism in this step?

(1)

- A electrophilic substitution
- B nucleophilic substitution
- C electrophilic addition
- D nucleophilic addition

(ii) Complete the diagram, including curly arrows, to show the mechanism for the reaction to form 2-phenylethanol.

(4)



(iii) In the synthetic route, what type of reaction is Step 2?

(1)

- A** hydrolysis
- B** esterification
- C** neutralisation
- D** hydrogenation

(iv) Name a suitable catalyst for Step 2.

(1)

.....

.....


.....

(Total for Question 7 = 17 marks)

BLANK PAGE

8 Propanoic acid can be synthesised from propene, a by-product of the petrochemical industry.

(a) Complete the equation to show the formation of propene and one other hydrocarbon.



(b) Propanoic acid can be synthesised from propene in 3 steps. A student proposed the following synthesis:

Step 1: propene → 1-bromopropane

Step 2: 1-bromopropane → **X**

Step 3: **X** → propanoic acid

(i) Identify **X** and give the conditions needed to carry out Step 2 and Step 3 in this synthesis.

(4)

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(ii) Explain, by considering the mechanism of the reaction, why the student's choice of reaction for Step 1 will lead to a low overall yield.

(2)

.....

.....

.....

.....

(c) Propene is used in the manufacture of polymers.

(i) Write an equation to show the formation of poly(propene) from propene. (2)

(ii) Describe how the chemical reaction to form the polymer nylon differs from the chemical reaction to form poly(propene). (2)

.....

.....

.....

.....

(iii) Give **two** ways in which chemists contribute to a more sustainable use of such materials as poly(propene) and nylon. (2)

.....

.....

.....

.....

(Total for Question 8 = 13 marks)

- 9 The table shows the formulae of three different halogenoalkanes and the optical activity of the products of their reaction with hydroxide ions in aqueous solution.

Reaction	Halogenoalkane reacting with hydroxide ions	Optical activity of product mixture
1	CH_3Cl	none
2	$\text{CH}_3\text{CHBrCH}_2\text{CH}_3$	significant
3	$\text{CH}_3\text{CH}_2\text{CH}_2\text{C}(\text{Br})(\text{CH}_3)\text{CH}_2\text{CH}_3$	no significant activity

(a) Draw the mechanism for reaction 1.

(4)

(b) The halogenoalkane in reaction 2 is 2-bromobutane.

(i) Name the halogenoalkane in reaction 3.

(1)

.....

.....

*(ii) Both halogenoalkanes used in reaction **2** and reaction **3** were a single optical isomer.

Explain the optical activity of the product mixture for reaction **2** and reaction **3**, in terms of the reaction mechanism.

(6)

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(c) Compounds containing the alkene functional group can show stereoisomerism.

(i) Draw the **two** structural isomers for an alkene that shows stereoisomerism.

(1)

(ii) Explain why stereoisomerism occurs in alkenes.

(2)

.....

.....

.....

.....

(Total for Question 9 = 14 marks)

TOTAL FOR PAPER = 90 MARKS

BLANK PAGE

The Periodic Table of Elements

1	2	3	4	5	6	7	0 (8)
6.9 Li lithium 3	9.0 Be beryllium 4	10.8 B boron 5	12.0 C carbon 6	14.0 N nitrogen 7	16.0 O oxygen 8	19.0 F fluorine 9	20.2 Ne neon 10
23.0 Na sodium 11	24.3 Mg magnesium 12	27.0 Al aluminium 13	28.1 Si silicon 14	31.0 P phosphorus 15	32.1 S sulfur 16	35.5 Cl chlorine 17	39.9 Ar argon 18
39.1 K potassium 19	40.1 Ca calcium 20	47.9 Ti titanium 22	54.9 Mn manganese 25	58.9 Co cobalt 27	65.4 Zn zinc 30	79.9 Br bromine 35	83.8 Kr krypton 36
85.5 Rb rubidium 37	87.6 Sr strontium 38	91.2 Zr zirconium 40	[98] Tc technetium 43	102.9 Rh rhodium 45	112.4 Cd cadmium 48	126.9 I iodine 53	131.3 Xe xenon 54
132.9 Cs caesium 55	137.3 Ba barium 56	178.5 Hf hafnium 72	186.2 Re rhenium 75	192.2 Ir iridium 77	200.6 Hg mercury 80	[210] At astatine 85	[222] Rn radon 86
[223] Fr francium 87	[226] Ra radium 88	138.9 La* lanthanum 57	183.8 W tungsten 74	192.2 Os osmium 76	204.4 Tl thallium 81	[209] Po polonium 84	
		89 Ac* actinium	[266] Sg seaborgium 106	[277] Hs hassium 108	207.2 Pb lead 82		
			[264] Bh bohrium 107	[268] Mt meitnerium 109			
			[147] Pm promethium 61	[271] Ds darmstadtium 110			
			150 Sm samarium 62	[272] Rg roentgenium 111			
			144 Nd neodymium 60				
			141 Pr praseodymium 59				
			140 Ce cerium 58				
			152 Eu europium 63				
			157 Gd gadolinium 64				
			163 Dy dysprosium 66				
			165 Ho holmium 67				
			167 Er erbium 68				
			169 Tm thulium 69				
			173 Yb ytterbium 70				
			175 Lu lutetium 71				
			232 Th thorium 90				
			[231] Pa protactinium 91				
			[237] Np neptunium 93				
			238 U uranium 92				
			[242] Pu plutonium 94				
			[243] Am americium 95				
			[247] Cm curium 96				
			[245] Bk berkelium 97				
			[251] Cf californium 98				
			[254] Es einsteinium 99				
			[253] Fm fermium 100				
			[256] Md mendelevium 101				
			[254] No nobelium 102				
			[257] Lr lawrencium 103				

1.0 H hydrogen 1

relative atomic mass
atomic symbol
name
atomic (proton) number

Elements with atomic numbers 112-116 have been reported but not fully authenticated

* Lanthanide series
* Actinide series