



# Mark Scheme (Results)

January 2021

Pearson Edexcel International GCSE  
In Chemistry (4CH1) Paper 2C

## **Edexcel and BTEC Qualifications**

Edexcel and BTEC qualifications are awarded by Pearson, the UK's largest awarding body. We provide a wide range of qualifications including academic, vocational, occupational and specific programmes for employers. For further information visit our qualifications websites at [www.edexcel.com](http://www.edexcel.com) or [www.btec.co.uk](http://www.btec.co.uk). Alternatively, you can get in touch with us using the details on our contact us page at [www.edexcel.com/contactus](http://www.edexcel.com/contactus).

## **Pearson: helping people progress, everywhere**

Pearson aspires to be the world's leading learning company. Our aim is to help everyone progress in their lives through education. We believe in every kind of learning, for all kinds of people, wherever they are in the world. We've been involved in education for over 150 years, and by working across 70 countries, in 100 languages, we have built an international reputation for our commitment to high standards and raising achievement through innovation in education. Find out more about how we can help you and your students at: [www.pearson.com/uk](http://www.pearson.com/uk)

January 2021

Publications Code 4CH1\_2C\_2101\_MS

All the material in this publication is copyright



© Pearson Education Ltd 2021

## General Marking Guidance

- All candidates must receive the same treatment. Examiners must mark the first candidate in exactly the same way as they mark the last.
- Mark schemes should be applied positively. Candidates must be rewarded for what they have shown they can do rather than penalised for omissions.
- Examiners should mark according to the mark scheme not according to their perception of where the grade boundaries may lie.
- There is no ceiling on achievement. All marks on the mark scheme should be used appropriately.
- All the marks on the mark scheme are designed to be awarded. Examiners should always award full marks if deserved, i.e. if the answer matches the mark scheme. Examiners should also be prepared to award zero marks if the candidate's response is not worthy of credit according to the mark scheme.
- Where some judgement is required, mark schemes will provide the principles by which marks will be awarded and exemplification may be limited.
- When examiners are in doubt regarding the application of the mark scheme to a candidate's response, the team leader must be consulted.
- Crossed out work should be marked UNLESS the candidate has replaced it with an alternative response.

Question number	Answer	Notes	Marks
1 (a) (i)	<p><b>D</b></p> <p><b>D</b> is the correct answer because protons occur in the nucleus and have a positive charge.</p> <p><b>A</b> is not the correct answer since electrons occur in the energy levels.</p> <p><b>B</b> is not the answer since ions do not occur in the nucleus.</p> <p><b>C</b> is not the correct answer since neutrons have no charge.</p>		1
(ii)	7		1
(iii)	lithium	<b>ALLOW</b> Li	1
(b) (i)	<p>M1 same number of protons</p> <p>M2 different number of neutrons</p>	<p><b>ALLOW</b> same number of electrons</p> <p><b>IGNORE</b> references to mass number and atomic number.</p>	2
5 marks			


Question number	Answer	Notes	Marks
2 (a) (i)	nitrogen	<b>ALLOW</b> N <sub>2</sub> /N	1
(ii)	carbon dioxide	<b>ALLOW</b> CO <sub>2</sub>	1
(iii)	argon	<b>ALLOW</b> Ar	1
(iv)	carbon dioxide	<b>ALLOW</b> CO <sub>2</sub>	1
(b)	lighted splint (produces squeaky) pop		1
5 marks			

Question number	Answer	Notes	Marks
3 (a) (i)	Any <b>two</b> from:  M1 volume of acid  M2 temperature  M3 mass / moles of magnesium  M4 surface area / size of pieces of magnesium		2
(ii)	so as little gas as possible escapes	<b>ALLOW</b> so no gas escapes <b>IGNORE</b> references to accuracy <b>REJECT</b> references to gas getting in	1
(b) (i)	M1 $(69 + 70 + 71) \div 3$  M2 70s	Answer of 70 with or without working scores 2  Answer of 76 or 75.8 or 75.75 with or without working scores 1	2
(ii)	as the (number of) carbons increases the time (to produce 10 cm <sup>3</sup> of hydrogen) increases ORA		1
(c)	M1 ester linkage as a displayed structure   M2 rest of molecule correct as a fully displayed structure 		2
8 marks			

Question number	Answer	Notes	Marks
4 (a) (i)	$2 \text{Na(s)} + 2 \text{H}_2\text{O(l)} \rightarrow 2 \text{NaOH(aq)} + \text{H}_2\text{(g)}$ M1 correct balancing numbers  M2 (s) and (aq) for state symbols	<b>ALLOW</b> multiples or fractions.	2
(ii)	hydroxide or $\text{OH}^-$	<b>REJECT</b> OH	1
(iii)	Any <b>three</b> from:  M1 the sodium moves (on the surface)  M2 effervescence or bubbles (of gas)  M3 (indicator or phenolphthalein or water) turns pink  M4 the sodium gets smaller  M5 the sodium melts or turns into a ball	<b>ALLOW</b> sodium floats  <b>IGNORE</b> gas or hydrogen produced  <b>IGNORE</b> initial colour of indicator  <b>ALLOW</b> the sodium disappears / (appears to) dissolve	3
(b)	M1 electron configuration of sodium is 2,8,1 and electron configuration of potassium is 2,8,8,1  M2 outer electron less attracted (to the nucleus of potassium)  M3 therefore (outer shell electron) is more easily lost	<b>ALLOW</b> the outer shell is further from the nucleus <b>ALLOW</b> potassium has more shells <b>ALLOW</b> larger atom / larger atomic radius  <b>ALLOW</b> reverse argument for sodium	3
9 marks			

Question number	Answer	Notes	Marks
5 (a) (i)	M1 layers / rows (of atoms / ions)  M2 can slide over one another	M2 is dependent on mention of layers / rows in M1	2
(ii)	M1 delocalised electrons  M2 can move / can flow / are free to move (throughout the structure)	<b>IGNORE</b> references to charge or current <b>IGNORE</b> free electrons M2 dependent on mention of electrons in M1	2
(b)	aluminium is more reactive than carbon	<b>ALLOW</b> references to position in reactivity series e.g. aluminium is higher in reactivity series than carbon. <b>ALLOW</b> carbon is less reactive than aluminium	1
(c) (i)	M1 aluminium / Al <sup>3+</sup> ions are attracted to the negative electrode / cathode (because they are positively charged)  M2 where they gain electrons (forming aluminium)	<b>ALLOW</b> Al <sup>3+</sup> + 3e <sup>-</sup> → Al <b>IGNORE</b> references to reduction	2
(ii)	2O <sup>2-</sup> → O <sub>2</sub> + 4e <sup>-</sup>	<b>ALLOW</b> 2O <sup>2-</sup> - 4e <sup>-</sup> → O <sub>2</sub>	1
(iii)	M1 electrodes are made of carbon  M2 which reacts with / burns in oxygen		2



(d) (i)	iron oxide loses oxygen	<b>IGNORE</b> references to electrons	1
(ii)	 <p>M1 right hand line below left hand line</p> <p>M2 correct name / formula of <b>both</b> reactants</p> <p>M3 correct name / formula of <b>both</b> products</p>	<p><b>IGNORE</b> horizontal axis drawn</p> <p><b>IGNORE</b> activation energy if shown</p> <p>If only use words <i>reactants</i> (on left) and <i>products</i> (on right) award 1 mark from M2 and M3</p> <p>M2 and M3 can be scored from an endothermic diagram</p>	3
14 marks			

Question number	Answer	Notes	Marks
6 (a)	<p>M1 mix / stir / add (silver nitrate and copper (II) chloride)</p> <p>M2 filter (the silver chloride)</p> <p>M3 wash with (deionised / distilled) water</p> <p>M4 dry in a warm oven <b>or</b> dry with filter paper <b>or</b> leave / allow to dry (on a windowsill) <b>or</b> dry in a desiccator</p>	<p><b>IGNORE</b> references to heating</p> <p>If evaporation is mentioned to form crystals max = 1</p>	4
(b)	<p>(i) M1 and M2 all points correctly plotted to <math>\pm</math> half a square</p> <p>(ii) M3 two straight lines of best fit which must meet at 3 cm and 6 cm<sup>3</sup></p> <p>(iii) Any <b>one</b> from:</p> <p>the precipitate wasn't left to settle (for long enough)</p> <p>the tube was on a slant</p> <p>not enough / less than 3.0 cm<sup>3</sup> of silver nitrate added</p> <p>(iv) all the copper [(II)] chloride has reacted</p>	<p>Deduct 1 mark for every incorrect point.</p> <p><b>ALLOW</b> measured the height too early</p> <p><b>IGNORE</b> references to human error unqualified</p> <p><b>ALLOW</b> the silver nitrate is in excess / not all the silver nitrate has reacted</p>	<p>2</p> <p>1</p> <p>1</p> <p>1</p>

Question number	Answer	Notes	Marks
(c) (i)	Any <b>one</b> from:  burette  (volumetric) pipette	<b>ALLOW</b> measuring cylinder  <b>REJECT</b> beaker	1
(ii)	Example calculation  M1 moles of copper chloride = $(25 \times 0.50) \div 1000$ <b>OR</b> 0.0125 moles  M2 moles of silver chloride = 0.0250  M3 mass of silver chloride = 3.59 g	<b>ALLOW</b> answer to M1 x 2  <b>ALLOW</b> answer to M1 or M2 $\times 143.5$ <b>ALLOW</b> 2 or more significant figures  Correct answer of 3.59 g scores 3 marks	3
(iii)	M1 $(0.744 \div 0.850) \times 100$  M2 87.5(%)	<b>ALLOW</b> 2 or more significant figures	2
			15 marks

Question number	Answer	Notes	Marks
7 (a)	<p>M1 crude oil is heated / vapourised</p> <p>M2 vapours / gases / compounds / hydrocarbons rise up the column</p> <p>M3 the column is hotter at the bottom than the top</p> <p>M4 vapours / compounds / hydrocarbons condense at their boiling point</p>	<p><b>ALLOW</b> boiled</p> <p><b>ALLOW</b> temperature gradient of the column</p> <p><b>ALLOW</b> vapours / compounds / hydrocarbons / condense at different heights</p> <p><b>ALLOW</b> the vapours / compounds / hydrocarbons / fractions have different boiling points.</p>	4
(b)	<p>M1 temperature of 600°C - 700°C</p> <p>M2 catalyst of silica / alumina</p>	<p><b>ALLOW</b> aluminosilicates / zeolites / silicon dioxide / aluminium oxide</p> <p><b>IGNORE</b> references to pressure</p>	2

Question number	Answer	Notes	Marks
(c)	(i) M1 nitrogen (from the air) reacts / combines with oxygen (from the air)  M2 at high temperatures (in the car engine)	<b>REJECT</b> any implication that oxygen or nitrogen come from the fuel.	2
	(ii) Any one from:  acid rain  respiratory problems		1
	(iii) Example calculation  M1 volume of carbon dioxide = $206\,000\text{ cm}^3 / 2.06 \times 10^5\text{ cm}^3 / 206\text{ dm}^3$  M2 volume of carbon dioxide per km = $51\,500\text{ cm}^3 / 5.15 \times 10^4\text{ cm}^3 / 51.5\text{ dm}^3$  M3 $(51\,500 \div 24\,000) = 2.15$ moles  M4 $M_r$ of carbon dioxide is 44  M5 mass of carbon dioxide per Km = 94.4 g	Division by 4 can happen in M1, M2, M3 or M5 <b>ALLOW</b> $M1 \div 4$  <b>ALLOW</b> M2 or $M1 \div 24\,000$  <b>ALLOW</b> 94 – 95 g <b>ALLOW</b> ecf from incorrect $M_r$  Correct answer of 94 – 95 g scores 5 marks.	5
			14 marks

