

**GCE**

**Physics B**

Unit **H557A/01**: Fundamentals of physics

Advanced GCE

**Mark Scheme for June 2017**

OCR (Oxford Cambridge and RSA) is a leading UK awarding body, providing a wide range of qualifications to meet the needs of candidates of all ages and abilities. OCR qualifications include AS/A Levels, Diplomas, GCSEs, Cambridge Nationals, Cambridge Technicals, Functional Skills, Key Skills, Entry Level qualifications, NVQs and vocational qualifications in areas such as IT, business, languages, teaching/training, administration and secretarial skills.

It is also responsible for developing new specifications to meet national requirements and the needs of students and teachers. OCR is a not-for-profit organisation; any surplus made is invested back into the establishment to help towards the development of qualifications and support, which keep pace with the changing needs of today's society.

This mark scheme is published as an aid to teachers and students, to indicate the requirements of the examination. It shows the basis on which marks were awarded by examiners. It does not indicate the details of the discussions which took place at an examiners' meeting before marking commenced.















All examiners are instructed that alternative correct answers and unexpected approaches in candidates' scripts must be given marks that fairly reflect the relevant knowledge and skills demonstrated.

Mark schemes should be read in conjunction with the published question papers and the report on the examination.

OCR will not enter into any discussion or correspondence in connection with this mark scheme.

© OCR 2017

Annotations available in RM Assessor

Annotation	Meaning
	Benefit of doubt given
	Contradiction
	Incorrect response
	Error carried forward
	Level 1
	Level 2
	Level 3
	Transcription error (in copying data from root of question – <b>ALLOW</b> method mark(s) if no further error but zero credit for evaluation)
	Benefit of doubt not given
	Power of 10 error
	Omission mark
	Error in number of significant figures
	Correct response
	Wrong physics or equation

## Section A: MCQs

Question	Answer	Marks	Guidance
----------	--------	-------	----------

Question			Answer	Marks	Guidance
1	D		D	1	
2	B		B	1	
3	A		A	1	
4	A		A	1	
5	B		B	1	
6	B		B	1	
7	A		A	1	
8	B		B	1	
9	C		C	1	
10	C		C	1	
11	A		A	1	
12	D		D	1	
13	C		C	1	
14	C		C	1	
15	D		D	1	
16	B		B	1	
17	D		D	1	
18	B		B	1	
19	D		D	1	
20	D		D	1	
21	A		A	1	
22	D		D	1	
23	C		C	1	
24	D		D	1	
25	B		B	1	
26	C		C	1	
27	B		B	1	
28	A		A	1	
29	B		B	1	
30	A		A	1	
			<b>Total</b>	<b>30</b>	

## Section B

Question		Answer	Marks	Guidance
31	(a)	(usually air is not a conductor) having no charge carriers (to support a current) / ✓  the ions and / or electrons provide charge carriers (for the air to conduct)	1	<b>expect</b> concept of charged particles free to move  <b>not</b> just air is an insulator or non-conductor / just ions are charged / just charges can conduct  <b>allow</b> charge carriers / delocalised or free electrons or charges / charges available to carry current / charges can flow
31	(b)	$(\Delta Q = I \Delta t) = 30 \times 10^3 \times 250 \times 10^{-6}$ ✓  $= 7.5 \text{ (C)}$ ✓	1  1	bare correct value scores both marks <b>ignore</b> units
<b>Total</b>			<b>3</b>	


Question		Answer	Marks	Guidance
32	(a)	$\frac{c/c_{\text{glass}}}{c/c_{\text{water}}}$ ✓	1	
32	(b)	$n_w = 1.3 / 1.6 = 0.81(3)$ ✓  $r = \sin^{-1}(\sin 30^\circ / 0.813) = 38^\circ$ ✓  OR $n_1 \sin \theta_1 = n_2 \sin \theta_2$ method  $r = \sin^{-1}(\sin 30^\circ \times 1.6 / 1.3)$ ✓ = $38^\circ$ ✓	1  1	first mark for evaluating / using correct index  <b>not</b> $24^\circ$ have inverted the index  <b>only</b> if this method is clear <b>allow</b> $n_1 / n_2 = 1.6 / 1.3 = 1.2(3)$ for first mark
<b>Total</b>			<b>3</b>	

Question			Answer	Marks	Guidance
33	(a)	(i)	$\Delta E = m c \Delta \theta$ $= 4200 \times 17 = 71.4$ (kJ)	✓ ✓	method evaluation <b>accept</b> 71 (kJ)
		(ii)	$\Delta t = m c \Delta \theta / I V$ or $= 71.4 \times 10^3 / (230 \times 46)$ $= 6.7(5)$ (s)	✓ ✓	method in rearranged algebra or numbers <b>accept</b> $t = E / P$ evaluation <b>not</b> 6.74 (s) RE <b>allow</b> ecf on value from (a)
33	(b)	$\Delta \theta$ doubled so flow (rate) or $\Delta m / \Delta t$ will have to halve	✓ ✓	1 1	<b>accept</b> mass flow rate drops from 0.15 to 0.075 kg s <sup>-1</sup> <b>not</b> $\Delta E$ doubles so time doubles / other time reasoning <b>max 1</b> for just flow (rate) less or slower
<b>Total</b>				<b>6</b>	

Question			Answer	Marks	Guidance	
34	(a)		$(4.5 \times 1 / 10) = 0.45$	✓	1	
			$(4.5 - 0.45) = 4.05$	✓	1	<b>ignore</b> 4.1
34	(b)	(i)	$\Delta Q \approx I \Delta t = V \Delta t / R$	✓	1	<b>accept</b> algebra expressed in words / = or $\approx$ symbols
			and $V = Q / C$ ( $\Rightarrow \Delta Q \approx Q \Delta t / R C$ )	✓	1	<b>not</b> credit for any exponential type reasoning
34	(b)	(ii)	assumes current / voltage / charge (on capacitor) is constant during $\Delta t$ (instead of continuously decaying)	✓	1	assumption <b>not</b> just rate is constant <b>accept</b> rate of charge flow is constant or rate of discharge is constant
			overcome by making $\Delta t$ smaller / as small as possible (as needed for better approximation ) <b>not</b> just make $\Delta t$ small	✓	1	how overcome for 1 standalone mark if no answer to assumption
<b>Total</b>				<b>6</b>		

Question		Answer	Marks	Guidance
35	(a)	$v = 0.24 \times 60 \times 3 \times 10^8 / \{60 \times 60 \times 24\}$ ✓ $= 5.0 \times 10^4 \text{ (m s}^{-1}\text{)}$ ✓	1 1	<b>accept</b> { $4.32 \times 10^9 \text{ m} \div 8.64 \times 10^4 \text{ s}$ }
35	(b)	range $R = 44.444 \times 60 \times 3 \times 10^8 = (8.0 \times 10^{11} \text{ m})$ ✓  $v_{\text{perp.}} = R \omega = 8.0 \times 10^{11} \times 1.8 \times 10^{-3} / (24 \times 3600)$ ✓ $= 1.66 \times 10^4 \text{ (m s}^{-1}\text{)}$ ✓  <b>OR alternative method for last 2 marks</b>  $s_{\text{perp.}} \approx R \theta = 8 \times 10^{11} \times 1.8 \times 10^{-3} = (1.44 \times 10^9 \text{ m})$  $v_{\text{perp}} = s_{\text{perp}} / t = 1.44 \times 10^9 / (24 \times 3600) = 1.66 \times 10^4 \text{ (m s}^{-1}\text{)}$	1  1 1	<b>accept</b> ranges based on either time or mean time of signal travel <b>all</b> give range = $8.0 \times 10^{11} \text{ m}$ (2 S.F.)  method  evaluation  method <b>accept</b> $\sin \theta \approx \tan \theta \approx \theta$ for small angle $\theta$ <b>n.b.</b> $s_{\text{perp.}} = 0.08$ light minutes can be credited  more method & evaluation  <b>allow</b> answers close to $280 \text{ m s}^{-1}$ to score 2/3 marks because light mins treated as light secs so 1/60 of correct answer so one small error
<b>Total</b> <b>Total section B</b>			<b>5</b> <b>23</b>	

## Section C

Question		Answer	Marks	Guidance
36	(a)	<p>velocity vector is changing direction constantly towards the centre of the orbit (magnitude / speed remains constant) ✓</p> <p>and acceleration = rate of change of velocity so there is an acceleration ✓</p> <p>OR (circular motion) requires a force towards the centre of the circle ( otherwise the mass will move in a straight line at a tangent to the circle) ✓</p> <p>and acceleration <math>\propto</math> force so there is an acceleration ( towards the centre ) ✓</p>	<p>1</p> <p>1</p>	<p><b>accept</b> in annotated diagram form</p>  <p>dependent on the first mark</p> <p><b>accept</b> (circular motion) requires centripetal force</p> <p><b>ignore</b> references to gravitational force of Earth on Moon cause Moon to accelerate towards earth</p> <p>dependent on the first mark <b>accept</b> <math>a = F / m</math> so there is an acceleration ( towards the centre )</p>
36	(b) (i)	<p><math>a = v^2 / R = \{2\pi R\}^2 / \{T^2 R\} = \dots\dots</math> ✓</p> <p>OR</p> <p><math>a = R \omega^2 = R \{2\pi / T\}^2 = \dots\dots</math></p>	1	<p>algebraic reasoning</p> <p><b>accept</b> using forces and <math>F = ma = m v^2 / R</math> and cancelling <math>m</math> and completing</p>
36	(b) (ii)	<p><math>4\pi^2 \times 3.84 \times 10^8 / (2.35 \times 10^6)^2 = 0.0027 \text{ m s}^{-2}</math> ✓</p>	1	<p>evaluation <b>accept</b> <math>2.74 \text{ mm s}^{-2} / 2.75 \text{ mm s}^{-2}</math> (<math>\pi \approx</math>)</p>
36	(b) (iii)	<p><math>g_{\text{at moon orbit}} = g_{\text{Earth surface}} / 60^2</math> ✓</p> <p><math>= 9.8 / 3600 = 2.7(2) \times 10^{-3} \text{ m s}^{-2}</math> ✓</p> <p>same value as (ii) ✓</p>	<p>1</p> <p>1</p> <p>1</p>	<p>method using inverse square law reasoning in numbers / words / algebra</p> <p>evaluation <b>accept</b> <math>g = 9.81 \text{ m s}^{-2}</math> / correct use of <math>a = G M / D^2</math></p> <p>comparison</p> <p><b>allow</b> ecf from (ii) if compared sensibly to <math>3 \text{ mm s}^{-2}</math></p>
<b>Total</b>			<b>7</b>	



Question			Answer	Marks	Guidance
37	(a)		$v_{\text{terminal}} = 0.65 \text{ (m s}^{-1}\text{)}$ ✓	1	$v_{\text{terminal}}$ read from graph <b>accept</b> in range 0.64 to 0.66 (m s <sup>-1</sup> )
			$\pm 0.02 \text{ (m s}^{-1}\text{)}$ ✓	1	uncertainty estimate apply SF penalty for 2 or 3 SF e.g. 0.019 or 0.0195 (m s <sup>-1</sup> ) scores 0
37	(b)		( at $t = 0.5 \text{ s}$ ) ball is accelerating (and $a$ is decreasing) ✓	1	<b>credit</b> numerical estimates of acceleration $\approx 0.55 \text{ m s}^{-2}$
			because downwards weight is larger than upwards drag force ✓	1	<b>accept</b> in algebra $W > D$ or $> (D + U)$ <b>accept</b> if upthrust $U$ is overlooked / air resistance or friction for drag <b>not</b> $U$ confused with $D$ <b>not</b> just because net force is downwards
37	(c)	(i)	temperature should be monitored or held constant ✓	1	<b>accept</b> density of the glycerol ; as it will affect upthrust
			since the viscosity / drag force will depend on $T$ ✓	1	<b>accept</b> viscosity of the glycerol ; as it will affect drag <b>accept</b> purity of the glycerol ; as moisture affects viscosity <b>accept</b> density of ball bearing ; as it will affect the weight <b>accept</b> mass of ball bearing ; as it will affect the weight <b>not</b> height drop or air bubbles in glycerol or keep same liquid <b>ignore</b> edge effect
37	(c)	(ii)	$D^2 / v_T = \text{constant}$ <b>OR</b> $v_T / D^2 = \text{constant}$ ✓	1	proposal <b>if</b> $v_T / D^2 = \text{constant}$ <b>accept</b> log / log graph <b>allow</b> $v_T = k D^2$
			$D^2 / v_T$ values: 144, 145, 144, 204, 221 (mm <sup>2</sup> m <sup>-1</sup> s) ✓✓ OR $v_T / D^2$ values (6.9, 6.9, 6.9, 4.9, 4.5) $\times 10^{-3}$ (m s <sup>-1</sup> mm <sup>-2</sup> ) ✓✓	2	working <b>expect</b> at least <b>2 data</b> tests for credit 1 mark and all <b>5 data</b> tested for 2 marks  <b>accept</b> table of $D^2$ values (for $v_T$ vs $D^2$ sketch graph) <b>accept</b> calculated log $v_T$ and log $D$ values same rule on data

Question	Answer	Marks	Guidance
	<p>noticing smallest three b.bs have almost constant <math>k</math> /</p> <p>largest two b.bs have a different sensible constant /</p> <p>smallest and largest b.bs <math>k</math> not constant</p> <p>.....</p> <p>consideration of <b>quantitative</b> uncertainty in <math>k</math></p> <p><math>D^2 / v_T = 144, 145, 144</math> constant to <math>\pm 0.3\%</math> /</p> <p><math>= 204, 221</math> constant to <math>\pm 4\%</math></p> <p>.....</p> <p>correct statement about <b>their test</b> showing proportionality or not showing proportionality</p> <p>.....</p> <p>If graphs sketched 2 marks from:</p> <p>sketches of log / log graph or sketches of <math>v_T</math> vs <math>D^2</math></p> <p>correct comment on gradient or linearity of <b>their</b> graph</p>	<p><b>2</b></p>	<p>conclusions <b>2 marks</b> available for <b>any sensible test involving 2 or more data points</b></p> <p><b>accept</b> any two or all three of smallest b.bs have almost constant <math>k</math></p> <p><b>accept</b> largest two b.bs have sensible constant <math>k</math> (if only 2 tested)</p> <p><b>accept</b> use of 3% uncertainty based on the uncertainty in <math>v_T</math> from (a)</p> <p>OR comment on differences in their <math>k</math> values 144, 221 show increase in <math>k</math> of about 50% / decrease of about 35%</p> <p><b>for graph method candidates</b></p>
	<p><b>Total</b></p>	<p><b>11</b></p>	


Question			Answer	Marks	Guidance
38	(a)		draw tangent and suitable large $\Delta$ at $t = 1$ or $3$ s ✓	1	method <b>accept</b> tangent and $\Delta s / \Delta t = 1.6 / 1.5$ <b>ignore</b> signs here award magnitude evaluation <b>accept</b> in range 1.0 to 1.2 ( $\text{m s}^{-1}$ )  alternative method for two marks
			1.1 ( $\text{m s}^{-1}$ ) ✓	1	
			OR identifying $A = 0.70$ m $f = 1/4$ Hz $v_{\max} = A \omega$ ;  $= 0.7 \times 2 \pi \times 1/4 = 1.1$ ( $\text{m s}^{-1}$ ) ✓		
38	(b)		- <b>sin</b> graph of period 4 s and shape by eye ✓ scaled to amplitude of $1.1 \text{ m s}^{-1}$ ✓	1 1	<b>accept</b> - sin graph scaled to agree with (a) ecf <b>ignore</b> shape of graph here just peak values
38	(c)	(i)	$L = T^2 g / 4\pi^2$ or equivalent using numbers ✓	1	method <b>not</b> just $T = 2 \pi \sqrt{L / g}$
			$= 4^2 \times 9.8 / 4\pi^2 = 3.97$ m ✓	1	evaluation <b>accept</b> $g = 9.81$ gives $L = 3.98$ m
<b>Total</b>				<b>6</b>	

Question			Answer	Marks	Guidance
39	(a)		<b>X</b> pure $\beta$ no $\gamma$ at background with 5 mm lead ✓	1	all identifications correct for <b>first mark</b> even if <b>no</b> explanations <b>total</b> zero if three sources incorrect <b>two</b> correct explanations for <b>second mark</b>  <b>three</b> correct explanations for <b>third mark</b>  <b>ignore</b> comments on $\beta$ as present in all three sources <b>not</b> credit for descriptions of data expect logical analysis
			<b>Y</b> $\alpha, \beta, \gamma$ large drop with paper $\therefore$ must have $\alpha$ ✓	1	
			<b>Z</b> $\beta, \gamma$ (no drop with paper $\therefore$ no $\alpha$ ) counts with lead so some $\gamma$ ✓	1	
39	(b)	(i)	$\Delta \log C / \Delta \log R$ or e.g. $(4.0 - 0.5) / (0.4 - 1.9) =$ ✓	1	method evidence of sensible gradient taken or tangent drawn or $\Delta$ constructed in downward section of graph even if sign is wrong  evaluation <b>accept</b> in range - 2.0 to - 2.5 must have correct sign here  <b>allow</b> both marks if bare answer in this range
			$= - 2.3$ ✓	1	
39	(b)	(ii)	✓ ✓ ✓ ✓  <b>Graph comments:</b>  $C = k / R^2$ gives $\log C = \log k - 2 \log R$ <b>OR</b> gradient close to 2 suggests $R^2$ variation and – sign indicates inverse relation $1 / R^2$  accuracy: - 2.3 is close but not perfect fit  for low range is not a good fit (log graph flat) <b>OR</b> higher range only is a reasonable fit	4	any 4 points from the list but must include a comment on graph <b>AND</b> a suggestion about radiation for full marks i.e. a <b>max 3</b> from each section  complete log analysis worth <b>2 marks accept</b> if $k$ taken as 1  conclusion <b>ecf</b> on their gradient value if outside range then not a good fit <b>accept</b> need to know $\pm$ uncertainties to estimate the significance of the small difference  i.e. recognising the significance of knee in graph

Question	Answer	Marks	Guidance
	<p><b>Suggestions about radiation:</b></p> <p><math>\alpha</math> attenuated by a few cms in air</p> <p>inverse square law applies to point sources, close to source will not be a good approximation / it will be more constant</p> <p><math>\gamma</math> should follow <math>1 / R^2</math> dilution</p> <p><math>\gamma</math> travel in straight lines from (point) source with little interaction / absorption by the air</p>		<p>accept <math>\beta</math> with explanation that these follow <math>1 / R^2</math> reasonably well up to this range</p>
	<b>Total</b>	<b>9</b>	

Question			Answer	Marks	Guidance
40	(a)		$(50 \text{ MPa} / 7 \times 10^{-4}) = 7.1(4) \times 10^{10} \text{ (Pa)}$ ✓	1	evaluation <b>accept</b> in range 7.0 to $7.3 \times 10^{10} \text{ (Pa)}$
40	(b)		<b>alloy</b> absorbs more energy (per volume) ✓ <b>alloy</b> is stronger / has higher breaking stress ✓	1 1	choice explained <b>accept</b> alloy because it is tougher <b>not</b> stiffer <b>not</b> any credit for pure metal takes greater strain and prolongs time of collision
40	(c)	(i)	method: any {scaled distance ÷ appropriate number of atoms} ✓  evaluation: e.g. 4 atoms per nm gives $0.25 \text{ nm} / 2.5 \times 10^{-10} \text{ (m)}$ ✓ OR 5 atoms per nm gives $0.20 \text{ nm} = 2.0 \times 10^{-10} \text{ (m)}$	1  1	<b>allow</b> atom counting angled to atomic rows <b>not</b> unreal estimates like 10 atoms per 1 nm  estimation <b>accept</b> in range $\{ 1.8 \text{ to } 2.7 \} \times 10^{-10} \text{ (m)}$ credit 2 marks for answer in range with no working
40	(c)	(ii)	a dislocation / edge dislocation ✓	1	<b>accept</b> extra half-plane of atoms
40	(d)		<b>Level 3 (5–6 marks)</b>  Marshals argument in a clear manner and includes clear <b>explanation of three strands</b> :  <ul style="list-style-type: none"> <li>• metallic bonding</li> <li>• structure of metal and alloy</li> <li>• elastic and plastic deformation</li> </ul> <p><i>There is a well-developed line of reasoning which is clear and logically structured. The information presented is relevant and substantiated.</i></p>	6	<b>Look for number of strands attempted to help decide the Level, then look at quality.</b>  <b>Indicative scientific points may include:</b>  <b>Metallic bonding</b> <ul style="list-style-type: none"> <li>• +ve ion lattice in sea / gas of free mobile electrons</li> <li>• non-directional strong electrostatic bond electron glue</li> <li>• similar for pure metal and alloy</li> </ul> <b>Structure of metal and alloy</b> <ul style="list-style-type: none"> <li>• ordered regular stacking of atoms in planes in metal</li> <li>• alloy has a few impurity metal atoms of different size</li> <li>• most metals are polycrystalline with grains and grain boundaries between crystals of different orientation</li> <li>• ions can slip and atomic planes move</li> </ul>

Question	Answer	Marks	Guidance
	<p><b>Level 2 (3–4 marks)</b></p> <p>Shows clear understanding of at least <b>two of the three strands</b> above to the argument <b>or</b> covers all three at a superficial manner and does not include enough indicative points for level 3.</p> <p><i>There is a line of reasoning presented with some structure. The information presented is in the most-part relevant and supported by some evidence.</i></p> <p><b>Level 1 (1–2 marks)</b></p> <p>Makes at least <b>two independent points</b> (possibly from one strand) that are relevant to the argument but does not link them together and shows only superficial engagement with the argument.</p> <p><i>There is an attempt at a logical structure with a line of reasoning. The information is in the most part relevant.</i></p> <p><b>0 marks</b></p> <p><i>no response worthy of credit remember No response at all record NR</i></p>		<ul style="list-style-type: none"> <li>• dislocations are stacking imperfections e.g. extra half plane reduce stress at which planes slip by localising stress</li> </ul> <p><b>Elastic and plastic deformation</b></p> <ul style="list-style-type: none"> <li>• elastic behaviour atoms return to original position when stress removed stretched stiff bonds spring back</li> <li>• metals and alloys are stiff and elastic for small strains</li> <li>• dislocation movement in pure metals allows slip and plastic deformation at relatively low stress this is permanent and ions do not return when stress removed</li> <li>• dislocations are pinned by impurity atoms in alloy which restricts slip giving a smaller plastic region at higher yield stress</li> <li>• reference to Fig. 40.2</li> <li>• <b>accept</b> well labelled diagrams throughout for credit if integrated into the explanation</li> </ul>
	<b>Total</b>	<b>12</b>	

Question		Answer	Marks	Guidance
41	(a)	. <b>N</b> on Fig. 41.1 ✓ ; . <b>V</b> on Fig. 41.2 ✓	2	<b>both</b> at mid-points of charges judged by eye and field lines (may need magnification to see amongst field lines)
41	(b) (i)		1 1	both sketches <b>any</b> 3 equipotentials of roughly correct shape judged by eye <b>accept</b> $\Delta V$ not equal (as diagram) <b>expect</b> attempt at orthogonality <b>accept</b> on Fig. 41.1 three equipotential loops surrounding both charges
41	(c)	$E = 2kQ/R^2$ or $= 2 \times 9 \times 10^9 \times 1 / 500^2$ ✓ $= 7.2 \times 10^4 \text{ (V m}^{-1}\text{)}$ ✓	1 1	method <b>must</b> have 2 factor for method mark evaluation <b>allow</b> 1 mark for $3.6 \times 10^4$ (½ correct value)
41	(d)	<b>Level 3 (5–6 marks)</b>  Marshals argument in a clear manner and includes clear explanation of <b>three strands</b> :  • work done • area under $E(R)$ field graph • gradient of $V(R)$ potential graph  <i>There is a well-developed line of reasoning which is clear and logically structured. The information presented is relevant and substantiated.</i>  <b>Level 2 (3–4 marks)</b>  Shows clear understanding of at least <b>two of the three strands</b> above to the argument <b>or</b> covers all three at a superficial manner and does not include enough indicative points for level 3.  <i>There is a line of reasoning presented with some structure. The information presented is in the most-part relevant and supported by some evidence.</i>	6	<b>Look for number of strands attempted to help decide the Level, then look at quality.</b>  <b>Do not penalise incorrect signs in this answer.</b>  <b>Indicative scientific points may include:</b>  <b>work done</b>  • work $W$ is done against electrical attraction of + and – charges which increases the electrical potential energy of the system • + charge is worked on in raising it up the potential well of the – charge • $W_{\text{total}} = \Sigma \Delta W = \Sigma F \Delta s = \Sigma F \Delta R$  <b>area under <math>E(R)</math> graph:</b>  <b>accept algebraic or numerical reasoning</b>  • $W_{\text{total}} = \Sigma \Delta W = \Sigma F \Delta s = \Sigma F \Delta R$ <b>only credit once</b> • $E_{\text{field}} = F / q$ but test charge is unit charge $q = 1 \text{ C}$ • in this example $E_{\text{field}} = F$



Question	Answer	Marks	Guidance
	<p><b>Level 1 (1–2 marks)</b></p> <p>Makes at least two independent points that are relevant to the argument but does not link them together and shows only superficial engagement with the argument.</p> <p><i>There is an attempt at a logical structure with a line of reasoning. The information is in the most part relevant.</i></p> <p><b>0 marks</b></p> <p>No response worthy of credit scores zero No response record NR</p>		<ul style="list-style-type: none"> <li>• <math>\therefore W =</math> Area under the field graph</li> <li>• 15 squares <math>\times 4 \times 10^6</math> J per square = 60 MJ C<sup>-1</sup> or MV</li> <li>• agrees with increase in potential from (-90 to -30) MV</li> <li>• recognising that <math>E = k Q / R^2</math></li> </ul> <p><b>gradient of <math>V(R)</math> graph</b></p> <ul style="list-style-type: none"> <li>• <math>E_{\text{field}} = -</math> gradient of <math>V(R) = - dV / dR = - \Delta V / \Delta R</math></li> <li>• tangent to graph drawn and shown = field</li> <li>• e.g. at <math>R = 200</math> m grad = (120 MV) / 600 m = <math>2 \times 10^5</math> V m<sup>-1</sup></li> <li>• agrees with the field at <math>R = 200</math> m of <math>20 \times 10^4</math> V m<sup>-1</sup></li> <li>• recognising that <math>V = k Q / R</math></li> </ul> <p><b>check</b> graphs for annotation credit</p>
	<b>Total</b>	<b>12</b>	
	<b>Total section C</b>	<b>56</b>	
	<b>Total sections B &amp; C</b>	<b>80</b>	

**OCR (Oxford Cambridge and RSA Examinations)**  
**1 Hills Road**  
**Cambridge**  
**CB1 2EU**

**OCR Customer Contact Centre**

**Education and Learning**

Telephone: 01223 553998

Facsimile: 01223 552627

Email: [general.qualifications@ocr.org.uk](mailto:general.qualifications@ocr.org.uk)

**[www.ocr.org.uk](http://www.ocr.org.uk)**

For staff training purposes and as part of our quality assurance programme your call may be recorded or monitored

**Oxford Cambridge and RSA Examinations**  
**is a Company Limited by Guarantee**  
**Registered in England**  
**Registered Office; 1 Hills Road, Cambridge, CB1 2EU**  
**Registered Company Number: 3484466**  
**OCR is an exempt Charity**

**OCR (Oxford Cambridge and RSA Examinations)**  
**Head office**  
**Telephone: 01223 552552**  
**Facsimile: 01223 552553**

© OCR 2017

