Mark Scheme

Mark Scheme 4752 MEI PURE MATHS C2 JANUARY 2005

Section A

	ion A		T -	
1	$6x^5 + \frac{1}{2}x^{-\frac{1}{2}}$ o.e.	B1	$6x^5$	
		B1	$x^{\frac{1}{2}}$ soi	
		B1	$\frac{1}{2}x^{-\frac{1}{2}}$ isw	3
2	$x^4/4$	B1	2	
	x^{-2}	D.0		
	$\overline{-2}$	B2	B1 for kx^{-2}	
	c	B1		4
3	At least 1 period of sine curve	G1		
	Sine curve from 0 to 360	G1	± 1 indicated	
	191.537 rot to 3 or more sf	B1	After B1 B1, -1 for extras in the range	
	348.463 rot to 3 or more sf	B1	SC1 for 192.8 and 347.2 (grads)	4
			SC1 for 180.2 and 359.8 (radians)	
4	9.0 or 8.96 or 8.960	В3	M1 for	
			$[BC^2=]6.8^2+4.1^2-2\times4.1\times6.8\times\cos108$	
	13.2577	B2	A1 for 80.2(8), 8.37(grads), 6.49 (rads) Correctly rounded to 3 or more sf	5
	13.2377		M1 for 0.5×4.1×6.8×sin108	
			For complete long methods using BC,	
			allow M1 and A1 for 13.2 to 13.3	[16]
5	a = 4, $r = 1/2$ identified	B1	Stated or identified by correct use	
	2 ⁻¹⁷	T2	M1 20 th term = their(a)x(their r) ¹⁹	
	8	S2	M1 S = their $(a)/(1$ -their (r))	5
6	4, 7, 10, 13, 16 ignore extras	B1	For showing 1 st four or 2 nd four terms	
	15250	B4	B1 for $d = 3$ soi	5
			B1 for $a = 4$ soi M1 for use of $100/2[2a + 99d]$ o.e.	5
7	2 12	В3	M1 for $30 = \frac{1}{2} \times 25 \times q$ o.e.	
-	(i) 2.4, $2\frac{2}{5}$, $\frac{12}{5}$		M1 for $q = (2 \times 30) / 5^2$	
				5
	(ii) 22	P2	M1 for $(arc =)5 x$ their 2.4	
8	(i) 2.5, 2.50, 2.500, 2.499	B2	M1 for log ₁₀ 316 or ln 316/ln 10	
	(ii) 6 www	В3	B2 for $6 \log_a a$ or $\log_a(a^6)$	
			Or B1 for $2\log_a(a)$ or $-\log_a a^{-4}$	5
			SC1 Using $a=10 \Rightarrow 6$	F4.03
			SC2 Using numerical a, not $10 \Rightarrow 6$	[20]
		1		1
			Total for section A	36

Mark Scheme 4752 MEI PURE MATHS C2 JANUARY 2005

Section B

9 iA 6.25 B2 M1 for $x = 5$ used to find \mathbf{B} (V =) area of cross-section × length $(\frac{100}{4})[\frac{10}{2}x^2 - \frac{1}{3}x^3]$ o.e.	and y
length	
length	
length	
$\left \frac{100}{100} \right \frac{10}{100} x^2 - \frac{1}{100} x^3 = 0.6.$	
M1 C-1	into their
[val at $x = 10$] – [val at $x = 0$] Subs of correct limits integrand	
4166 to 4167 or 4170 A2 A1 for 166.6 or 1666	66.6 or 5
41.6rot to 3 sf or me	ore
ii 52.62 B4 M3 for-	
$2/2 \times [2.15 \times 2 + 2(5.64 \times 2)]$	2+6.44x2)]oe
Or M2 if one slip	
Or M1 if 2 slips or one	e trap evaluated 5
	[12]
Their (5262) – their (4167) M1 Must be >0	
10 i $y' = 3x^2 - 12x$ B1B1	
use of $y' = 0$ M1	
x = 0 and 4 A1	
(0, 12) and $(4, -20)$ A1 Allow $y = 12$ and $y = -20$	-20
y'' = 6x - 12 used M1 y' used each side of TF	_
\int max when $x = 0$, min when $x = 4$ A1 Both stated, only one is	needs testing
ii when $x = 2$ $y' = -12$ B1	
grad of normal = $1/12$ B1ft from their y'	
y + 4 = 1/12(x - 2) M1ft accept any numerical r	
$Or -4 = their(m) \times 2 +$	
$y = \frac{1}{12}x - 4\frac{1}{6}$ Any recognisable 25/6	, at worst 4.1
11 i Excess temperature B1	2
At t = 0 oe B1	
ii $\log z = \log z_0 + \log (10^{-kt})$ B1	
$= \log z_0 - kt \log 10$ B1 nb AG	2
iii Z= 46 31 20 14 9 If z= 68, 53P1, L1,N	M1, M1, M1
available	
log z : 1.66 1.49 1.30 1.15 0.95 T1	
correctly plotted P1 ft their values, within 2	
line of best fit L1 Ruled, using their point	
k = 0.017 to 0.019 or 0.02 G2 M1 for attempting +/-	_
$z_0 = 66 \text{ to } 73$ B2 M1 for (log) $z_0 = 1.82 \text{ to}$	to 1.86
1 1 1 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2	$\times 10^{-70 \text{ x their } k}$ [13]
temp of drink = 25 to 27	[[[[[[[[[[[[[[[[[[[