



---

# LEVEL 2 CERTIFICATE FURTHER MATHEMATICS 8360/2

Paper 2      Calculator

---

Mark scheme

June 2019

---

Version: 1.0 Final

\*jun1983602/MS\*

Mark schemes are prepared by the Lead Assessment Writer and considered, together with the relevant questions, by a panel of subject teachers. This mark scheme includes any amendments made at the standardisation events which all associates participate in and is the scheme which was used by them in this examination. The standardisation process ensures that the mark scheme covers the students' responses to questions and that every associate understands and applies it in the same correct way. As preparation for standardisation each associate analyses a number of students' scripts. Alternative answers not already covered by the mark scheme are discussed and legislated for. If, after the standardisation process, associates encounter unusual answers which have not been raised they are required to refer these to the Lead Assessment Writer.

It must be stressed that a mark scheme is a working document, in many cases further developed and expanded on the basis of students' reactions to a particular paper. Assumptions about future mark schemes on the basis of one year's document should be avoided; whilst the guiding principles of assessment remain constant, details will change, depending on the content of a particular examination paper.

Further copies of this mark scheme are available from [aqa.org.uk](http://aqa.org.uk)

## Glossary for Mark Schemes

GCSE examinations are marked in such a way as to award positive achievement wherever possible. Thus, for GCSE Mathematics papers, marks are awarded under various categories.

If a student uses a method which is not explicitly covered by the mark scheme the same principles of marking should be applied. Credit should be given to any valid methods. Examiners should seek advice from their senior examiner if in any doubt.

<b>M</b>	Method marks are awarded for a correct method which could lead to a correct answer.
<b>M dep</b>	A method mark dependent on a previous method mark being awarded.
<b>A</b>	Accuracy marks are awarded when following on from a correct method. It is not necessary to always see the method. This can be implied.
<b>B</b>	Marks awarded independent of method.
<b>B dep</b>	A mark that can only be awarded if a previous independent mark has been awarded.
<b>ft</b>	Follow through marks. Marks awarded following a mistake in an earlier step.
<b>SC</b>	Special case. Marks awarded within the scheme for a common misinterpretation which has some mathematical worth.
<b>oe</b>	Or equivalent. Accept answers that are equivalent. eg, accept 0.5 as well as $\frac{1}{2}$
<b>[a, b]</b>	Accept values between $a$ and $b$ inclusive.
<b>3.14...</b>	Accept answers which begin 3.14 eg 3.14, 3.142, 3.1416

Examiners should consistently apply the following principles.

***Diagrams***

Diagrams that have working on them should be treated like normal responses. If a diagram has been written on but the correct response is within the answer space, the work within the answer space should be marked. Working on diagrams that contradicts work within the answer space is not to be considered as choice but as working, and is not, therefore, penalised.

***Responses which appear to come from incorrect methods***

Whenever there is doubt as to whether a candidate has used an incorrect method to obtain an answer, as a general principle, the benefit of doubt must be given to the candidate. In cases where there is no doubt that the answer has come from incorrect working then the candidate should be penalised.

***Questions which ask candidates to show working***

Instructions on marking will be given but usually marks are not awarded to candidates who show no working.

***Questions which do not ask candidates to show working***

As a general principle, a correct response is awarded full marks.

***Misread or miscopy***

Candidates often copy values from a question incorrectly. If the examiner thinks that the candidate has made a genuine misread, then only the accuracy marks (A or B marks), up to a maximum of 2 marks are penalised. The method marks can still be awarded.

***Further work***

Once the correct answer has been seen, further working may be ignored unless it goes on to contradict the correct answer.

***Choice***

When a choice of answers and/or methods is given, mark each attempt. If both methods are valid then M marks can be awarded but any incorrect answer or method would result in marks being lost.

***Work not replaced***

Erased or crossed out work that is still legible should be marked.

***Work replaced***

Erased or crossed out work that has been replaced is not awarded marks.

***Premature approximation***

Rounding off too early can lead to inaccuracy in the final answer. This should be penalised by 1 mark unless instructed otherwise.

***Continental notation***

Accept a comma used instead of a decimal point (for example, in measurements or currency), provided that it is clear to the examiner that the candidate intended it to be a decimal point.

Q	Answer	Mark	Comments	
<b>1(a)</b>	<b>Alternative method 1</b>			
	$3a = 4(2a + 3)$ or $3a = 8a + 12$ or $5a = 4b$	M1	oe equation	
	$(a =) -\frac{12}{5}$ or $-2\frac{2}{5}$ or $-2.4$	A1		
	$(b =) -3$	A1ft	ft their $a \times 1.25$ evaluated or $\frac{\text{their } 5a}{4}$ evaluated $a \neq 0$ if awarding ft M1 is implied	
	<b>Alternative method 2</b>			
	$3a + 5a = 4(2a + 3) + 4b$ or $8a = 8a + 12 + 4b$ or $4b = -12$	M1	oe equation	
	$(b =) -3$	A1		
	$(a =) -\frac{12}{5}$ or $-2\frac{2}{5}$ or $-2.4$	A1ft	ft their $b \times 0.8$ evaluated or $\frac{\text{their } 4b}{5}$ evaluated $b \neq 0$ if awarding ft M1 is implied	
	<b>Additional Guidance</b>			
	Alt 1 $a = 2$ $b = 2.5$ ( $5a = 4b$ is implied)			M1A0A1ft
	$\begin{pmatrix} 3a \\ 5a \end{pmatrix} = \begin{pmatrix} 8a + 12 \\ 4b \end{pmatrix}$			M0
Accept $-\frac{12}{5}$ or $\frac{12}{-5}$ for $-\frac{12}{5}$ (apply throughout scheme for values)				
Only solutions seen with one correct and the other incorrect (or missing)			2 marks	

Q	Answer	Mark	Comments
1(b)	$2m + 2 = 1$ or $2m + 1 = 0$ or $\frac{1-2}{2}$ or $\begin{pmatrix} 2m+2 & 2m+1 \\ 0 & 1 \end{pmatrix} = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$	M1	oe equation or calculation
	$-\frac{1}{2}$ or $-0.5$	A1	
	<b>Additional Guidance</b>		
	Condone missing brackets in $\begin{pmatrix} 2m+2 & 2m+1 \\ 0 & 1 \end{pmatrix} = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$		
	Allow $\begin{pmatrix} 2m+2 & 2m+1 \\ 2-2 & 2-1 \end{pmatrix} = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$		
	Mark positively eg error in matrix multiplication but $2m + 2 = 1$ and answer $-0.5$	M1A1	
	More than one answer given is A0 eg $m + 2 = 1$ and $2m + 1 = 0$ (mark positively) Answer $-1$ and $-0.5$	M1 A0	

Q	Answer	Mark	Comments
2	$\left(\frac{4+6}{2}, \frac{1+9}{2}\right)$ or (5, 6)	M1	oe eg $\left(4 + \frac{6-4}{2}, 1 + \frac{11-1}{2}\right)$ may be on diagram
	$\frac{1--3}{4-10}$ or $\frac{4}{-6}$ or $\frac{0-\text{their } 6}{14-\text{their } 5}$ or $\frac{-6}{9}$	M1	oe method for at least one gradient or at least one unsimplified gradient seen eg $\frac{-3-1}{10-4}$ or $\frac{-4}{6}$ or $\frac{\text{their } 6-0}{\text{their } 5-14}$ or $\frac{6}{-9}$  $\frac{6-0}{5-14}$ or $\frac{6}{-9}$ is M1M1
	$\frac{1--3}{4-10}$ or $\frac{4}{-6}$ and $\frac{0-6}{14-5}$ or $\frac{-6}{9}$ and shows that the gradients are equal	A1	oe method for both gradients or two unsimplified gradients seen and gradients shown to be equal eg $\frac{4}{-6}$ and $\frac{-6}{9}$ and these are both $-\frac{2}{3}$  SC2 (5, 6) and at least one gradient given as $-\frac{2}{3}$  SC1 at least one gradient given as $-\frac{2}{3}$

Additional Guidance is on the next page

Q	Answer	Mark	Comments
---	--------	------	----------

<b>Additional Guidance</b>			
<b>2 cont</b>	Mark intention for 1st M1 eg condone 5, 6	M1	
	$\frac{4}{-6} = -\frac{2}{3}$ and $\frac{-6}{9} = -\frac{2}{3}$	M2A1	
	$\frac{1-3}{4-10} = -\frac{2}{3}$ and $\frac{0-6}{14-5} = -\frac{2}{3}$	M2A1	
	$\frac{4}{-6} = \frac{-6}{9}$	M2A1	
	$\frac{4}{-6}$ and $\frac{-6}{9}$ and parallel	M2A0	
	$\frac{4}{6}$ is 2nd M0 unless recovered to $-\frac{4}{6}$		
	$\frac{4}{6}$ recovered to $-\frac{4}{6}$ and $\frac{6}{9}$ recovered to $-\frac{6}{9}$ could go on to score full marks		
	both gradients = $-\frac{2}{3}$ with no method or unsimplified gradients seen cannot score the A mark		
	$\frac{4}{-6}x$ or $\frac{-6}{9}x$ do not score 2nd M1 unless recovered		
	Equation of a line does not score 2nd M1 unless a method or unsimplified gradient seen		
	Using the reciprocals of gradients can score a maximum of M1M0A0		
	Allow $-0.66\dots$ or $-0.67$ for $-\frac{2}{3}$ and $\frac{4}{-6}$ etc Ignore conversion attempt after a correct fraction is seen		
	or method for $\frac{4}{-6}$ $1 = 4m + c$ and $-3 = 10m + c$ $4 = -6m$ $\frac{4}{-6} = m$ (similar method possible for $\frac{-6}{9}$ )	(2nd) M1	



Q	Answer	Mark	Comments
3	$a^2 < 0$ <input type="checkbox"/> <input type="checkbox"/> <input checked="" type="checkbox"/>	B4	B1 for each correct row
	$-1 < b^3 < 1$ <input checked="" type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/>		
	$\frac{b}{a} < 0$ <input type="checkbox"/> <input checked="" type="checkbox"/> <input type="checkbox"/>		
	$a - b > 0$ <input type="checkbox"/> <input checked="" type="checkbox"/> <input type="checkbox"/>		
	<b>Additional Guidance</b>		
	Two boxes ticked in a row with other 3 rows fully correct	B3	
One row correct, two rows blank, all three boxes ticked in another row	B1		
Only crosses used instead of ticks eg cross in all 4 correct boxes with all other boxes blank	B4		
Ticks and crosses used – only mark the ticks for that row eg Top row has X X ✓ scores B1 for that row Second row has X ✓ X scores B0 for that row			

Q	Answer	Mark	Comments
<b>4</b>	$(y =) \frac{3}{2}x \dots$ or $(y =) 1.5x \dots$ or $\frac{3}{2}$ or 1.5	M1	oe eg $(y =) \frac{3x-9}{2}$
	$\frac{x^5 - 17}{10} = \frac{3}{2}$	M1dep	oe implies M2
	$x^5 = \frac{3}{2} \times 10 + 17$ or $\sqrt[5]{32}$ or correctly rearranges $\frac{x^5 - 17}{10} = k$ to the form $x^5 =$ (k any non-zero value)	M1	oe eg $x^5 = 15 + 17$ or $x^5 = 32$ or $\sqrt[5]{15+17}$ must rearrange to the form $x^5 =$
	2	A1	

**Additional Guidance is on the next page**

Q	Answer	Mark	Comments
---	--------	------	----------

<b>Additional Guidance</b>				
<b>4 cont</b>	Condone error seen in rearrangement of $3x - 2y = 9$ if gradient is $\frac{3}{2}$ May go on to score M3A1			
	$\frac{x^5 - 17}{10} = \frac{3}{2}x$	M1M0M0A0		
	(gradient =) 3 $\frac{x^5 - 17}{10} = 3$ $x^5 = 30 + 17 \quad (3\text{rd M is not dependent})$ 2.16	M0M0dep		M1 A0
	$\frac{3}{2}$ $\frac{x^5 - 17}{10} = -\frac{2}{3}$ $x^5 = -\frac{2}{3} \times 10 + 17 \quad (3\text{rd M is not dependent})$ 1.595	M1  M0		M1  A0
	Condone answer (2, ...)			
	2 embedded			M3A0

Q	Answer	Mark	Comments
5(a)	$a^{-2}$	B2	B1 applies an index law or changes fourth root to fractional or decimal power in a correct expression eg $\sqrt[4]{a^{-8}}$ or $(a^{-8})^{\frac{1}{4}}$ or $(a^8)^{-\frac{1}{4}}$ or $a^{-\frac{8}{4}}$ or $\frac{1}{a^{\frac{8}{4}}}$ or $\frac{1}{a^2}$ or $a^{\frac{1}{4}} \times a^{-\frac{9}{4}}$ or $a^{\frac{1}{4}} \times \frac{1}{a^{\frac{9}{4}}}$ or $\left(\frac{1}{a^8}\right)^{\frac{1}{4}}$ or $\sqrt[4]{\frac{1}{a^8}}$ or $\sqrt[4]{a \times \frac{1}{a^9}}$ or $(a \times a^{-9})^{\frac{1}{4}}$ or $\left(a \times \frac{1}{a^9}\right)^{\frac{1}{4}}$
	<b>Additional Guidance</b>		
	$a^{-\frac{8}{4}}$ or $a^{\frac{8}{-4}}$	B1	
	$a^{-2}$ in working with $-2$ on answer line	B1	
	$a^{-2}$ in working with $\frac{1}{a^2}$ on answer line	B1	
	B1 response followed by further work is still awarded B1		
	Allow 0.25 for $\frac{1}{4}$ etc		
Allow recovery of missing brackets			

Q	Answer	Mark	Comments
5(b)	$32c^2d^2$ or $32(cd)^2$	B3	B2 (numerator =) $64c^3d^6$ or single term answer with two of 32, $c^2$ and $d^2$ (not in a denominator)  B1 single term answer with one of 32, $c^2$ and $d^2$ (not in a denominator)  SC2 factorised correct expression eg $16cd(2cd)$
	<b>Additional Guidance</b>		
	$2c^2d^2$ or $32c^2d$ or $32c^2$ or $\frac{32d^2}{c^3}$ or $\frac{c^2d^2}{32}$ or $64(cd)^2$ etc	B2	
	$32c^3d$ or $c^2$ or $\frac{d^2}{c}$ or $\frac{c^2d}{32}$ or $\frac{32}{c^2}$ etc		B1
	$\frac{32c^2d^2}{1}$ or $\frac{32(cd)^2}{1}$	B2	
	Allow denominator of 1 in a B2 or B1 answer eg $\frac{32c^2d}{1}$	B2	
	Multiplication signs in a correct expression eg $32 \times c^2 \times d^2$	B2	
	Allow multiplication signs in a B2, SC2 or B1 answer eg $32 \times c^3 \times d$	B1	
	Do not accept $2^5$ for 32 eg $2^5c^2d$	B1	
	If answer line scores B1 or B0 check working lines for possible response for up to 2 marks		
$32c^2d^2$ in working with different answer on answer line	B2		

Q	Answer	Mark	Comments
6(a)	$A(-\frac{3}{2}, 0)$ and $B(2, 0)$	B2	oe B1 $A(-\frac{3}{2}, 0)$ oe or $B(2, 0)$ SC1 $A(2, 0)$ and $B(-\frac{3}{2}, 0)$ oe
	<b>Additional Guidance</b>		
	Ignore the diagram		

6(b)	$-\frac{3}{2} < x < 2$ or $2 > x > -\frac{3}{2}$	B1ft	oe correct or ft their values from (a) must be a single inequality in x
------	--	------	---

**Additional Guidance is on the next page**

Q	Answer	Mark	Comments
---	--------	------	----------

<b>Additional Guidance</b>			
<b>6(b) cont</b>	$-\frac{3}{2} \leq x < 2$	B0	
	$-\frac{3}{2} > x < 2$	B0	
	$-\frac{3}{2} < x$ and $x < 2$	B0	
	their (a) $A(-2, 0)$ and $B(\frac{3}{2}, 0)$ (B0 in (a)) (b) $-2 < x < \frac{3}{2}$	B1ft	
	their (a) $A(2, 0)$ and $B(-\frac{3}{2}, 0)$ (SC1 in (a)) (b) $2 < x < -\frac{3}{2}$	B0ft	
	their (a) $A(-3, 0)$ and $B(2, 0)$ (B1 in (a)) (b) $2 < x < -3$	B0ft	
	their (a) $A(4, 0)$ and $B(-2, 0)$ (B0 in (a)) (b) $-2 < x < 4$	B1ft	
	Only one value in (a) can only score in (b) for $-\frac{3}{2} < x < 2$ or $2 > x > -\frac{3}{2}$		

Q	Answer	Mark	Comments
---	--------	------	----------

7(a)	Horizontal straight line	B1	mark intention
	<b>Additional Guidance</b>		
	Ignore any attempt at an equation		
	Mark the entire graph on the grid		
	Ignore any graph not on the grid		
	Line clearly drawn on the x-axis	B1	
	Line does not need to start from the y-axis		
	Ignore any points plotted		

7(b)	Straight line with gradient $> 0$	B1	mark intention
	<b>Additional Guidance</b>		
	Ignore any attempt at an equation		
	Mark the entire graph on the grid		
	Ignore any graph not on the grid		
	Vertical line	B0	
	A straight line joined to another line with a different gradient	B0	
	Line does not need to start at (0, 0)		
Ignore any points plotted			



Q	Answer	Mark	Comments	
8(a)	$7 + 12\sqrt{5} + 6(9 - 2\sqrt{5})$ or $12\sqrt{5} + 6(-2\sqrt{5}) = 0$ or $12\sqrt{5} \div 2\sqrt{5} = 6$ or states that need to add 6 lots of $(9 - 2\sqrt{5})$ or 7th term	M1	oe eg $7 + 6 \times 9$ or $7 + 54$ or $6 \times -2 = -12$  allow $7 + 12\sqrt{5} + (n-1)(9 - 2\sqrt{5})$ with $n = 7$  allow $7 + 12\sqrt{5} + n(9 - 2\sqrt{5})$ with $n = 6$	
	61		A1	
	<b>Additional Guidance</b>			
	61 in working lines with 7(th) on answer line			M1A0
	If repeatedly adding $(9 - 2\sqrt{5})$ they must stop after adding 6 lots or clearly select the relevant one			
	Answer 6 or 6th term with M1 not seen			M0A0
	Ignore any conversions to decimals			
Beware $(9 - 2\sqrt{5})(9 + 2\sqrt{5}) = 61$			M0A0	

Q	Answer	Mark	Comments	
8(b)	$\frac{29}{5}$ or $5\frac{4}{5}$ or 5.8	B2	oe eg $5\frac{8}{10}$ B1 any two of 1, $\frac{11}{5}$ , $\frac{26}{10}$ oe values	
	<b>Additional Guidance</b>			
	Terms must be evaluated for B1 unless correct answer seen			
	eg1 $\frac{3-1}{1+1} + \frac{12-1}{4+1} + \frac{27-1}{9+1}$			B0
	eg2 $\frac{3-1}{1+1} + \frac{12-1}{4+1} + \frac{27-1}{9+1} = 5.8$			B2
1 7 2.6			B1	
Ignore conversion attempts after a correct value seen				

Q	Answer	Mark	Comments
8(c)	<b>Alternative method 1</b>		
	(Second differences =) 4 or $2n^2$	M1	second differences seen at least once and not contradicted may be seen by the sequence
	$-3 - 2 \quad 3 - 8 \quad (13 - 18 \quad 27 - 32)$ or $-5 \quad -5 \quad (-5 \quad -5)$	M1dep	subtracts $2n^2$ from the given terms
	$2n^2 - 5$	A1	oe eg $2n^2 + 0n - 5$ does not need terms collected
	<b>Alternative method 2</b>		
	(Second differences =) 4 or $2n^2$	M1	second differences seen at least once and not contradicted may be seen by the sequence
	$3a + b = 3 - -3$ and substitutes $a = 2$ or $b = 0$	M1dep	oe
	$2n^2 - 5$	A1	oe eg $2n^2 + 0n - 5$ does not need terms collected

Mark scheme and Additional Guidance continue on the next page

Q	Answer	Mark	Comments
<b>8(c) cont</b>	<b>Alternative method 3</b>		
	Any three of $a + b + c = -3$ $4a + 2b + c = 3$ $9a + 3b + c = 13$ $16a + 4b + c = 27$	M1	
	$3a + b = 3 - -3$ and $5a + b = 13 - 3$ or $a = 2$ and $b = 0$	M1dep	oe obtains two correct equations in same two variables from their equations
	$2n^2 - 5$	A1	oe eg $2n^2 + 0n - 5$ does not need terms collected
	<b>Alternative method 4</b>		
	(Second differences =) 4 or $2n^2$	M1	second differences seen at least once and not contradicted may be seen by the sequence
	$2 \times 1^2 + b \times 1 - 5 = -3$ or $2 + b - 5 = -3$ or $b = 0$	M1dep	$2n^2 + bn - 5 = -3$ with $n = 1$ substituted oe eg $2 \times 2^2 + b \times 2 - 5 = 3$
	$2n^2 - 5$	A1	oe eg $2n^2 + 0n - 5$ does not need terms collected
	<b>Additional Guidance</b>		
	Condone working in a different variable		
	Alt 1 2nd M1 Subtracting given terms from $2n^2$ leading to $5 \ 5 \ (5 \ 5)$ must be recovered eg final answer $2n^2 - 5$ ( $2n^2 - 5n$ or $2n^2 - 5n - 5$ is not a recovery)		
	Answer $2n^2$ ..... scores at least M1		
Condone $n = 2n^2 - 5$ or $2n^2 - 5 = 0$		M2A1	

Q	Answer	Mark	Comments
<b>9</b>	$(p + 6)^{10}(p + 5)$ or $(p + 5)(p + 6)^{10}$	B2	B1 $(p + 6)^{10}(p + 6 - 1)$ or states $x = p + 6$ and $x^{10}(x - 1)$ (any letter for $x$ other than $p$ ) or correct partial factorisation eg $(p + 6)[(p + 6)^{10} - (p + 6)^9]$ or $(p + 6)^2[(p + 6)^9 - (p + 6)^8]$
	<b>Additional Guidance</b>		
	Any shape of bracket may be used		
	$(p + 6)^{10}((p + 6) - 1)$		B1
	Missing brackets must be recovered eg $p + 5 (p + 6)^{10}$ not recovered and B1 response not seen		B0
	Condone $(p + 6)^{10}(p + 5$		B2
	Condone $(p + 6)^{10}(p + 6 - 1$		B1
	$(p + 6)^{10}(p + 5)$ followed by expansion attempt		B1
	B1 response followed by expansion attempt		B1
	$(p + 6)^{10} \times (p + 5)$		B1
	Condone multiplication signs for B1 eg $(p + 6)^{10} \times (p + 6 - 1)$		B1
$(p + 6)^{11} \left[ 1 - \frac{1}{p + 6} \right]$		B1	

Q	Answer	Mark	Comments
10(a)	$f(x) \leq 25$ or $25 \geq f(x)$	B2	B1 $f(x) < 25$ or $k \leq f(x) \leq 25$ or $k < f(x) \leq 25$ where k is any number $< 25$  SC1 $\leq 25$ or $x \leq 25$
	<b>Additional Guidance</b>		
	Condone $f(x)$ replaced by eg y or f or fx or $F(x)$ or F or Fx or $x^3 - 2$ in B2 or B1 responses		
	Equivalent inequalities may be seen $25 > f(x)$	B1	
	Allow $-\infty < f(x) \leq 25$	B2	
	Condone $-\infty \leq f(x) \leq 25$	B2	
	$-\infty < f(x) < 25$ or $-\infty \leq f(x) < 25$	B1	
	$[-\infty, 25]$ or $(-\infty, 25]$	B1	
	$(-\infty, 25)$	B0	
	Condone $f(x) = \leq 25$	B2	
	Condone $f(x) = < 25$	B1	
	Condone $f(x) = x \leq 25$	SC1	
	$f(x) \leq 25$ in working with list of integers on answer line	B1	
	Only a list of integers	B0	

Q	Answer	Mark	Comments
10(b)	$1 \leq g(x) \leq 5$ or $5 \geq g(x) \geq 1$	B2	B1 $1 \leq g(x) < 5$ or $1 < g(x) \leq 5$ or $1 < g(x) < 5$ or $g(x) \geq 1$ and $g(x) \leq 5$ or $1 \leq g(x) \leq k$ where k is a constant $> 1$ or $p \leq g(x) \leq 5$ where p is a constant $< 5$ SC1 $1 \leq x \leq 5$
	<b>Additional Guidance</b>		
	Condone $g(x)$ replaced by eg y or g or gx or $f(x)$ or f or fx or $5 - x^2$ in B2 or B1 responses		
	Equivalent inequalities may be seen eg $5 \geq g(x) > 1$		B1
	Only $g(x) \geq 1$ given as the answer		B0
	Only $g(x) \leq 5$ given as the answer		B0
	$1 \leq g(x) \leq 4$		B1
	$1 \leq g(x) < 4$		B0
	$0 \leq g(x) \leq 5$		B1
	$0 < g(x) \leq 5$		B0
	Invalid statements do not score eg1 $1 \leq g(x) \geq 5$ eg2 $1 \geq g(x) \leq 5$ eg3 $6 \leq g(x) \leq 5$		B0 B0 B0
	$[1, 5]$		B1
	$[1, 5)$ or $(1, 5]$ or $(1, 5)$ or $1 - 5$ or $5 - 1$		B0
	$1 \leq g(x) \leq 5$ in working with list of integers on answer line		B1
	Only a list of integers		B0

Q	Answer	Mark	Comments
11	$x = -2$	B1	
	<b>Additional Guidance</b>		
12	<b>Alternative method 1</b>		
	$\frac{1}{2} \times \frac{4}{3} \times \pi \times (6a)^3$ or $\frac{2}{3} \times \pi \times 216a^3$ or $144\pi a^3$	M1	oe eg $\frac{1}{2} \times \frac{4}{3} \times \pi \times \left(\frac{12a}{2}\right)^3$ or $\frac{2}{3} \times \pi \times (6a)^3$
	$a^3 = \frac{486\pi}{144\pi}$ or $a^3 = \frac{27}{8}$ or $a^3 = 486 \div \left(\frac{2}{3} \times 6^3\right)$ or $a^3 = 3.375$ or $\sqrt[3]{3.375}$	A1	oe equation of form $a^3 =$ or calculation allow $(6a)^3 = 729$ or $6a = 9$
$\frac{3}{2}$ or $1\frac{1}{2}$ or 1.5	A1	SC1 answer 0.75 oe or answer 1.19... or answer 4.95...	

**Mark scheme and Additional Guidance continue on the next page**



Q	Answer	Mark	Comments
<b>12 cont</b>	<b>Alternative method 2</b>		
	$r^3 = \frac{486\pi}{\frac{2}{3}\pi}$ or $r^3 = 729$ or $\sqrt[3]{729}$ or 9	M1	oe equation of form $r^3 =$ or calculation
	$6a = \sqrt[3]{\frac{486\pi}{\frac{2}{3}\pi}}$ or $6a = 9$ or $9 \div 6$	A1	oe equation or calculation allow $(6a)^3 = 729$
	$\frac{3}{2}$ or $1\frac{1}{2}$ or 1.5	A1	SC1 answer 0.75 oe or answer 1.19... or answer 4.95...
	<b>Additional Guidance</b>		
	Allow recovery of missing brackets		
Allow use of $\pi = [3.14, 3.142]$			

Q	Answer	Mark	Comments
13	$x(1 - x^2)$ or $2x(1 + x)$ or $x(2 + 2x)$ or $\frac{1 - x^2}{2 + 2x}$	M1	implied by 2nd M1 oe factorisation eg $-x(x^2 - 1)$
	$x(1 + x)(1 - x)$ or $\frac{x(1 - x^2)}{2x(1 + x)}$ or $\frac{1 - x^2}{2(1 + x)}$ or $\frac{(1 + x)(1 - x)}{2 + 2x}$	M1dep	implies M2 oe factorisation eg $-x(x + 1)(x - 1)$
	$\frac{x(1 + x)(1 - x)}{2x(1 + x)}$ or $\frac{(1 + x)(1 - x)}{2(1 + x)}$ or $\frac{x(1 - x)}{2x}$	M1dep	implies M3 oe factorisation eg $\frac{-x(x + 1)(x - 1)}{2x(1 + x)}$
	$\frac{1 - x}{2}$ with M3 seen	A1	oe simplest form eg $\frac{1}{2}(1 - x)$ or $\frac{1}{2} - \frac{1}{2}x$ or $\frac{-x + 1}{2}$
	<b>Additional Guidance</b>		
	$\frac{x(1 + x)(1 - x)}{2x(1 + x)}$ or $\frac{(1 + x)(1 - x)}{2(1 + x)}$ or $\frac{x(1 - x)}{2x}$ is sufficient working	M3	
	$2(x + x^2)$ with no further work	M0	
$\frac{x - 1}{-2}$ with M3 seen or $-\frac{1}{2}(x - 1)$ with M3 seen or $\frac{-(x - 1)}{2}$ with M3 seen	M3A1		

Q	Answer	Mark	Comments	
14	$a^2 + (3a)^2 - 2 \times a \times 3a \times \cos 120$ or $\cos 120 = \frac{a^2 + (3a)^2 - b^2}{2 \times a \times 3a}$	M1	oe eg may substitute $\cos 120 = -0.5$ may be seen in a square root	
	$b^2 = a^2 + 9a^2 + 3a^2$ or $b^2 = 13a^2$ or $-b^2 = -13a^2$ or $b = \sqrt{13} a$	A1	oe equation of the form $b^2 =$ or $b =$ with brackets expanded and terms fully simplified $\cos 120 = -0.5$ substituted	
	13 : 1	A1	SC1 7 : 1	
	<b>Additional Guidance</b>			
	Allow recovery of missing brackets			
	$a^2 = a^2 + (3a)^2 - 2 \times a \times 3a \times \cos 120$ not recovered			M1M0A0
	$b^2 = 10a^2 - - 3a^2$			M1A0A0
$b^2 = 10a^2 + 3a^2$			M1A1A0	

Q	Answer	Mark	Comments
15	<b>Alternative method 1</b> $3mp = 3(2p + 1) + p + 5$ or $(m =) \frac{3(2p+1)}{3p} + \frac{p+5}{3p}$ or $(m =) \frac{6p+3+p+5}{3p}$	M1	oe fractions eliminated or common denominator eg $(m =) \frac{3p(2p+1)}{3p^2} + \frac{p(p+5)}{3p^2}$ or $(m =) \frac{6p^2+3p+p^2+5p}{3p^2}$
	$3mp = 6p + 3 + p + 5$ or $3mp = 7p + 8$	M1dep	oe brackets expanded and fractions eliminated eg $3mp^2 = 7p^2 + 8p$ implies M2
	$3mp - 7p = 8$ or $\frac{8}{3m-7}$ or $\frac{-8}{7-3m}$	M1dep	oe terms collected eg $p(3m-7) = 8$ or $7p - 3mp = -8$ implies M3
	$p = \frac{8}{3m-7}$ or $p = \frac{-8}{7-3m}$	A1	oe eg $\frac{8}{3m-7} = p$

**Mark scheme and Additional Guidance continue on the next page**

Q	Answer	Mark	Comments
<b>15 cont</b>	<b>Alternative method 2</b>		
	$(m =) \frac{3(2p+1)}{3p} + \frac{p+5}{3p}$ or $(m =) \frac{6p+3+p+5}{3p}$	M1	oe common denominator eg $(m =) \frac{3p(2p+1)}{3p^2} + \frac{p(p+5)}{3p^2}$ or $(m =) \frac{6p^2+3p+p^2+5p}{3p^2}$
	$m = \frac{7p+8}{3p}$ and $m = \frac{7}{3} + \frac{8}{3p}$ and $m - \frac{7}{3} = \frac{8}{3p}$	M1dep	simplifies numerator and isolates term in p eg $m = \frac{7p^2+8p}{3p^2}$ and $m = \frac{7}{3} + \frac{8}{3p}$ and $m - \frac{7}{3} = \frac{8}{3p}$ implies M2
	$\frac{3m-7}{3} = \frac{8}{3p}$	M1dep	converts $m - \frac{7}{3}$ to a single fraction implies M3
	$p = \frac{8}{3m-7} \text{ or } p = \frac{-8}{7-3m}$	A1	oe eg $\frac{8}{3m-7} = p$
	<b>Additional Guidance</b>		
	$p = \frac{8}{3m-7}$ in working but $\frac{8}{3m-7}$ on answer line		M3A1
	Allow recovery of missing brackets		
	$p = \frac{8}{3m-7}$ followed by incorrect further work		M3A0
	Allow equivalences for A1 eg $p = \frac{\frac{8}{3}}{\frac{3m-7}{3}}$		M3A1
Do not regard eg $3m(p) = 7p + 8$ as having unexpanded brackets		M1M1dep	

Q	Answer	Mark	Comments	
<b>16</b>	$\frac{8-5}{2} = \sqrt{1-a} \quad \text{or} \quad \frac{3}{2} = \sqrt{1-a}$ or $3^2 = 2^2(1-a) \quad \text{or} \quad 9 = 4(1-a)$	M1		
	$1-a = \left(\frac{3}{2}\right)^2$ or $1-a = \frac{9}{4}$ or $9 = 4 - 4a$ or $\frac{4-9}{4}$	M1dep	oe equation or calculation eg $1-a = \left(\frac{8-5}{2}\right)^2$ or $1-a = 2.25$ or $\frac{9-4}{-4}$ implies M2	
	$-\frac{5}{4} \quad \text{or} \quad -1.25 \quad \text{or} \quad -1\frac{1}{4}$	A1		
	<b>Additional Guidance</b>			
	$3 = 2\sqrt{1-a}$			M0
	Allow recovery of missing brackets			

Q	Answer	Mark	Comments
<b>17</b>	$x^2 + 3x + x + 3$ with three terms correct or $x^2 + 4x + k$ where k is a non-zero constant	M1	oe expansion attempt of one pair of brackets eg1 $x^2 + 4x + 3x + 12$ with three terms correct or $x^2 + 7x + k$ where k is a non-zero constant eg2 $x^2 + 4x + x + 4$ with three terms correct or $x^2 + 5x + k$ where k is a non-zero constant
	$x^3 + 3x^2 + x^2 + 3x$ or $x^3 + 4x^2 + 3x$ or $4x^2 + 12x + 4x + 12$ or $4x^2 + 16x + 12$	M1dep	attempt at a full expansion with correct multiplication of their 3 or 4 terms by one of the terms in the remaining bracket oe eg $x^3 + 4x^2 + 3x^2 + 12x$ or $x^3 + 7x^2 + 12x$ or $x^2 + 4x + 3x + 12$ or $x^2 + 7x + 12$ ( $x^2 + 7x + 12$ must be from an attempt at a full expansion) or $x^3 + 4x^2 + x^2 + 4x$ or $x^3 + 5x^2 + 4x$ or $3x^2 + 12x + 3x + 12$ or $3x^2 + 15x + 12$
	$x^3 + 8x^2 + 19x + 12$	A1	fully correct expansion allow if terms not collected eg $x^3 + 3x^2 + x^2 + 3x + 4x^2 + 12x + 4x + 12$ or $x^3 + 4x^2 + 3x + 4x^2 + 16x + 12$
	$x^2 + 8x + 12$	A1ft	ft M2A0 full simplification of their $(x^3 + 8x^2 + 19x + 12) - x^3 - 7x^2 - 11x$ their $(x^3 + 8x^2 + 19x + 12)$ must be a cubic
	$x^2 + 8x + 12$ and $(x + 6)(x + 2)$ or $(x + 2)(x + 6)$	A1	oe product of brackets

**Additional Guidance is on the next page**

Q	Answer	Mark	Comments
---	--------	------	----------

17 cont	<b>Additional Guidance</b>		
	1st M1 Do not allow omissions or extras eg1 $x^2 + 3x + 3$ eg2 $x^2 + 3x + x + 3 + x^2$	M0 M0	
	For the first 2 marks terms may be seen in a grid		
	If 1st A1 has been awarded with terms not collected, A1ft can still be awarded using their simplified cubic eg $x^3 + 4x^2 + 3x + 4x^2 + 16x + 12$ $= x^3 + 8x^2 + 18x + 12$ $x^3 + 8x^2 + 18x + 12 - x^3 - 7x^2 - 11x$ $= x^2 + 7x + 12$	M1M1A1  A1ftA0	
	First A1 may be seen embedded eg $x^3 + 8x^2 + 19x + 12 - x^3 + 7x^2 - 11x$	M1M1A1	
	If an attempt at the expansion of all three brackets in one go is made it must be fully correct to gain M2A1, otherwise M0M0A0 eg $x^2 + 3x + x + 3 + x^2 + 4x$	M0M0A0	
	Allow recovery of missing brackets when subtracting $x^3 + 7x^2 + 11x$ from their cubic		
	For final A1 allow $x^2 + 8x + 12$ and $a = 6$ $b = 2$ or $x^2 + 8x + 12$ and $a = 2$ $b = 6$		
	Ignore equating to zero and/or any 'solving' of an equation		



Q	Answer	Mark	Comments
<b>18</b>	<b>Alternative method 1</b>		
	$x - 5 = \frac{k}{2}$ or $x - 5 = -\frac{k}{2}$ or $2(x - 5) = k$ or $2x - 10 = k$ or $2(x - 5) = -k$ or $2x - 10 = -k$	M1	oe linear equation eg $x - 5 = \sqrt{\frac{k^2}{4}}$ or $x = \frac{k}{2} + 5$ or $\sqrt{4}(x - 5) = \sqrt{k^2}$
	$x - 5 = \frac{k}{2}$ and $x - 5 = -\frac{k}{2}$ or $2(x - 5) = k$ and $2(x - 5) = -k$ or $2x - 10 = k$ and $2x - 10 = -k$	A1	oe eg $x - 5 = \pm \frac{k}{2}$ square root(s) must be processed implied by final A1
	$\frac{k}{2} + 5$ and $-\frac{k}{2} + 5$	A1	oe simplest form eg $\frac{10+k}{2}$ and $\frac{10-k}{2}$ or $\frac{k+10}{2}$ and $\frac{k-10}{-2}$ or $5 \pm 0.5k$
	<b>Alternative method 2</b>		
	$4x^2 - 40x + 100 - k^2 (= 0)$	M1	expands and collects terms
	$\frac{- -40 \pm \sqrt{(-40)^2 - 4 \times 4 \times (100 - k^2)}}{2 \times 4}$	A1	oe eg $\frac{40 \pm \sqrt{16k^2}}{8}$ or $\frac{40 \pm 4k}{8}$ implied by final A1
	$\frac{k}{2} + 5$ and $-\frac{k}{2} + 5$	A1	oe simplest form eg $\frac{10+k}{2}$ and $\frac{10-k}{2}$ or $\frac{k+10}{2}$ and $\frac{k-10}{-2}$ or $5 \pm 0.5k$
<b>Additional Guidance</b>			
Allow recovery of missing brackets			

Q	Answer	Mark	Comments
19(a)	$\sqrt{(3x)^2 + (4x)^2} (= 5x)$ or $\sqrt{9x^2 + 16x^2} (= 5x)$ or $(3x)^2 + (4x)^2 = (5x)^2$ or 3x, 4x, 5x triangle	B1	may be seen in stages eg $9x^2 + 16x^2 = 25x^2$ and $\sqrt{25x^2} (= 5x)$
	<b>Additional Guidance</b>		
	Only $\sqrt{25x^2} (= 5x)$ seen	B0	
	Pythagorean triple 3x, 4x, 5x	B1	
	Pythagorean triple 3, 4, 5	B0	
	Missing brackets can <b>not</b> be recovered eg1 $\sqrt{3x^2 + 4x^2} = 5x$	B0	
	eg2 $3x^2 + 4x^2 = 9x^2 + 16x^2 = 25x^2$ and $\sqrt{25x^2} (= 5x)$	B0	
	Incorrect statements are B0 (mark the full response) eg1 $9x^2 + 16x^2 = 25x^2 = \sqrt{25x^2} (= 5x)$ eg2 $9x^2 + 16x^2 = 25x^2 \quad \sqrt{25} x^2 (= 5x)$ eg3 $\sqrt{(3x)^2 + (4x)^2} = 5x$ and $3x + 4x = 5x$	B0 B0 B0	
Only uses values for x	B0		

Q	Answer	Mark	Comments
<b>19(b)</b>	<b>Alternative method 1</b>		
	$0.5 \times 4x \times 3x$ or $6x^2$	M1	oe may be seen on the diagram
	$(6.5x)^2 - (2.5x)^2$ or $42.25x^2 - 6.25x^2$ or $36x^2$	M1	oe eg $(6.5x)^2 - \left(\frac{5x}{2}\right)^2$
	$\sqrt{\text{their } 36x^2}$ or $6x$	M1dep	dep on 2nd M1 may be seen on the diagram
	$0.5 \times 5x \times \text{their } 6x$ or $15x^2$	M1dep	oe dep on 2nd and 3rd M1
	$21x^2$	A1	allow $p = 21$ if areas $6x^2$ and $15x^2$ seen
	<b>Alternative method 2</b>		
	$0.5 \times 4x \times 3x$ or $6x^2$	M1	oe may be seen on the diagram
	$\cos ACD = \frac{2.5x}{6.5x}$ or $\cos ACD = \frac{5}{13}$	M1	oe
	$\cos^{-1} \frac{2.5x}{6.5x}$ or $67(.3\dots)$ or $67.4$	M1dep	oe eg $\cos^{-1} \frac{(6.5x)^2 + (5x)^2 - (6.5x)^2}{2 \times 6.5x \times 5x}$  dep on 2nd M1
	$0.5 \times 5x \times 6.5x \times \sin \text{their } 67(.3\dots)$ or $15x^2$	M1dep	oe dep on 2nd and 3rd M1
	$21x^2$	A1	allow $p = 21$ if areas $6x^2$ and $15x^2$ seen

**Mark scheme and Additional Guidance continue on the next page**

Q	Answer	Mark	Comments
19(b) cont	<b>Alternative method 3</b>		
	$0.5 \times 4x \times 3x$ or $6x^2$	M1	oe may be seen on the diagram
	$(5x)^2 = (6.5x)^2 + (6.5x)^2$ $- 2 \times 6.5x \times 6.5x \times \cos D$	M1	oe
	$\cos^{-1} \frac{(6.5x)^2 + (6.5x)^2 - (5x)^2}{2 \times 6.5x \times 6.5x}$ or $\cos^{-1} \frac{119}{169}$ or 45(.2...)	M1dep	oe dep on 2nd M1
	$0.5 \times 6.5x \times 6.5x \times \sin$ their 45(.2...) or $15x^2$	M1dep	oe dep on 2nd and 3rd M1
	$21x^2$	A1	allow $p = 21$ if areas $6x^2$ and $15x^2$ seen
	<b>Additional Guidance</b>		
	Allow recovery of algebra eg1 $0.5 \times 4 \times 3 = 6$ is 1st M0 but if recovered to $6x^2$ scores 1st M1 eg2 Alt 1 $\sqrt{42.25 - 6.25} = 6$ is 2nd M0 and 3rd M0 but if recovered to $6x$ scores 2nd M1 and 3rd M1		
	Do not allow final mark if an incorrect area is seen eg do not allow answer $21x^2$ if their two areas are $6x^2$ and $15x$		
	Answer $21x^2$ with no incorrect working eg fully correct working with numbers and final answer $21x^2$		5 marks
Allow recovery of missing brackets			
Choose the scheme that favours the student			

Q	Answer	Mark	Comments
20(a)	<b>Alternative method 1</b>		
	Full method leading to angle $BCD = 180 - 2x$	M1	eg angle $CFE = x$ and angle $FCE = 180 - 2x$ and angle $BCD = 180 - 2x$
	Full reasoning for their method	A1	eg (base angles of) isosceles (triangle are equal) and (sum of) angles in a triangle (is 180) and (vertically) opposite angles
	angle $BAD = 2x$ and (opposite angles of) cyclic quadrilateral (add to 180)	A1	must see M1
	<b>Alternative method 2 Working out angle <math>DCF</math> using angle at centre</b>		
	angle $DCF = 2x$	M1	
	angle at centre (is double angle at circumference)	A1	
	Full method leading to angle $BAD = 2x$ and full reasoning for their method	A1	must see M1 eg angle $BCD = 180 - 2x$ and angle $BAD = 2x$ and angles on a (straight) line (add to 180) and (opposite angles of) cyclic quadrilateral (add to 180)

Mark scheme and Additional Guidance continue on the next three pages

Q	Answer	Mark	Comments
---	--------	------	----------

<b>20(a) cont</b>	<b>Alternative method 3 Working out angle <math>DCF</math> not using angle at centre</b>		
	Full method leading to angle $DCF = 2x$	M1	eg angle $CFE = x$ and angle $DCF = 2x$
	Full reasoning for their method	A1	eg (base angles of) isosceles (triangle are equal) and exterior angle (of triangle is sum of interior opposite angles)
Full method leading to angle $BAD = 2x$ and full reasoning for their method	A1	must see M1 eg angle $BCD = 180 - 2x$ and angle $BAD = 2x$ and angles on a (straight) line (add to 180) and (opposite angles of) cyclic quadrilateral (add to 180)	

**Mark scheme and Additional Guidance continue on the next two pages**

Q	Answer	Mark	Comments
---	--------	------	----------

20(a) cont	<b>Alternative method 4</b>		
	Full method leading to angle $DFC = 90 - x$ and angle $ABC = 90 - x$	M1	eg angle $CFE = x$ and angle $DFE = 90$ and angle $DFC = 90 - x$ and angle $CDF = 90 - x$ and angle $ADC = 90 + x$ and angle $ABC = 90 - x$
	Full reasoning for their method	A1	eg (base angles of) isosceles (triangle are equal) and (angle in a) semicircle (is 90) and (sum of) angles in a triangle (is 180) and angles on a (straight) line (add to 180) and (opposite angles of) cyclic quadrilateral (add to 180)
angle $BAD = 2x$ and (sum of) angles in a triangle (is 180)	A1	must see M1	

**Additional Guidance is on the next page**

Q	Answer	Mark	Comments
---	--------	------	----------

	<b>Additional Guidance</b>		
<b>20(a) cont</b>			
	It is possible to score M1A1A0 or M1A0A1		
	Do not award any marks from angles on the diagram		
	Angles must be stated unambiguously eg condone angle <i>B</i> but do not condone angle <i>D</i>		
	'angle' may be missing or replaced by a symbol - mark intention		
	angle <i>CFE</i> may be seen as angle <i>EFC</i> or angle <i>BFE</i> etc		
	For (base angles of) isosceles (triangle are equal) allow radii (are equal)		
	For (sum of) angles in a triangle (is 180) allow triangle is 180		
	Use judgement when considering wording of reasons and allow abbreviations		
	Alt 2 Final A1 reason may be exterior angle of cyclic quadrilateral (equals interior opposite angle)		
	Choose the scheme that favours the student		
	Ignore angles that are not needed for their scheme even if incorrect		
	Allow recovery of missing brackets		
	Starting with angle <i>BAD</i> = 2x		M0A0A0



Q	Answer	Mark	Comments
20(b)	30	B2	B1 correct equation or calculation eg $90 + 2x + x = 180$ or $90 - x = 2x$ or $3x = 90$ or $6x = 180$ or $90 \div 3$
	<b>Additional Guidance</b>		
	Ignore any expressions for angles and any other calculated angles		
	Ignore any reasons		

Q	Answer	Mark	Comments
21	$8^2 + 12^2$ or $64 + 144$ or $208$ or $8^2 + 12^2 + 15^2$ or $64 + 144 + 225$ or $433$	M1	$HC^2$ or $CE^2$ implied by 2nd M1
	$\sqrt{8^2 + 12^2}$ or $\sqrt{208}$ or $4\sqrt{13}$ or $14.4\dots$ or $\sqrt{8^2 + 12^2 + 15^2}$ or $\sqrt{433}$ or $20.8\dots$	M1dep	oe may be on diagram fully correct trigonometry method leading to $14.4\dots$ or $20.8\dots$ can score M2 eg $8 \div \sin\left(\tan^{-1}\frac{8}{12}\right)$ or $8 \div \sin\left(\tan^{-1}\frac{8}{\sqrt{12^2 + 15^2}}\right)$
	$\tan x = \frac{15}{\sqrt{8^2 + 12^2}}$ or $\cos x = \frac{\sqrt{8^2 + 12^2}}{\sqrt{8^2 + 12^2 + 15^2}}$ or $\sin x = \frac{15}{\sqrt{8^2 + 12^2 + 15^2}}$	M1dep	oe eg $\tan x = [1.04, 1.042]$ or or $\cos x = [0.69, 0.6934]$ or $\sin x = [0.72, 0.7212]$ or $90 - \tan^{-1}\frac{\sqrt{8^2 + 12^2}}{15}$ dep on M2 any letter
	46(.1...)	A1	
	<b>Additional Guidance</b>		
	3rd M1 If using sine rule or cosine rule, must be in the form $\cos x =$ or $\sin x =$ eg $\cos x = \frac{20.8^2 + 14.4^2 - 15^2}{2 \times 20.8 \times 14.4}$ (oe eg $\cos^{-1} [0.69, 0.6934]$ )		
3rd M1 Condone $\tan = \frac{15}{\sqrt{8^2 + 12^2}}$ etc		M3	
Allow the first 2 M marks even if not subsequently used			
Allow recovery of missing brackets			

Q	Answer	Mark	Comments
<b>22(a)</b>	<b>Alternative method 1</b>		
	$2\sin^2x - 1 + 1 - \sin^2x$ or $2\sin^2x - (\sin^2x + \cos^2x) + \cos^2x$ or $2\sin^2x - \sin^2x - \cos^2x + \cos^2x$ or $2\sin^2x - \sin^2x$ or $\sin^2x - \cos^2x + \cos^2x$ or $1 + \sin^2x - 1$	M1	use of $\sin^2x + \cos^2x = 1$ in numerator ignore any denominator
	$\frac{\sin^2x}{\sin x \cos x}$ with M1 seen	$\frac{\sin^2x}{\tan x \cos^2x}$ with M1 seen	M1dep
$\frac{\sin x}{\cos x}$ and $\tan x$ with M2 seen	$\frac{\tan^2x}{\tan x}$ and $\tan x$ with M2 seen	A1	SC3 equates given expression to $\tan x$ and cross multiplies to show equivalence with full working shown

**Mark scheme and Additional Guidance continue on the next two pages**

Q	Answer	Mark	Comments	
<b>22(a) cont</b>	<b>Alternative method 2</b>			
	$2(1 - \cos^2 x) - 1 + \cos^2 x$ or $2 - 2\cos^2 x - 1 + \cos^2 x$	M1	use of $\sin^2 x + \cos^2 x = 1$ in numerator ignore any denominator	
	$\frac{1 - \cos^2 x}{\sin x \cos x}$ and $\frac{\sin^2 x}{\sin x \cos x}$ with M1 seen	$\frac{1 - \cos^2 x}{\sin x \cos x}$ and $\frac{\sin^2 x}{\tan x \cos^2 x}$ with M1 seen	M1dep	simplification to one step from $\frac{\sin x}{\cos x}$ or simplification to one step from $\frac{\tan^2 x}{\tan x}$
	$\frac{\sin x}{\cos x}$ and $\tan x$ with M2 seen	$\frac{\tan^2 x}{\tan x}$ and $\tan x$ with M2 seen	A1	SC3 equates given expression to $\tan x$ and cross multiplies to show equivalence with full working shown
	<b>Alternative method 3</b>			
	$\frac{2\sin x}{\cos x} - \frac{\sin^2 x}{\sin x \cos x}$	M1	from $\frac{2\sin^2 x}{\sin x \cos x} - \frac{1 - \cos^2 x}{\sin x \cos x}$	
	$2\tan x - \frac{\sin^2 x}{\sin x \cos x}$ or $\frac{2\sin x}{\cos x} - \frac{\sin x}{\cos x}$ with M1 seen	M1dep	simplification to one step from $2\tan x - \tan x$	
	$2\tan x - \tan x$ and $\tan x$ with M2 seen	A1	SC3 equates given expression to $\tan x$ and cross multiplies to show equivalence with full working shown	

**Additional Guidance is on the next page**



Q	Answer	Mark	Comments
<b>22(b)</b>	135 and 315 with no other solutions [0, 360]	B2	B1 135 with no other solutions [0, 360] or 315 with no other solutions [0, 360]  SC1 135 and 315 with one other solution [0, 360]
	<b>Additional Guidance</b>		
	Mark the answer line unless blank eg 135 and 315 in working with 135 on answer line	B1	
	–45 and 135 and 315	B2	
	–45 and 135	B1	
	Ignore incorrect solutions outside the range [0, 360] eg 135 and 315 and –90	B2	
	135 and 225 and 315	SC1	
	Both answers embedded ie tan 135 tan 315	B1	
	0 and 135 and 225 and 315	B0	
	45 and 135	B0	
	225 and 315	B0	
<b>23(a)</b>	(1, –3)	B1	
	<b>Additional Guidance</b>		
	Mark intention eg condone 1, –3	B1	

Q	Answer	Mark	Comments
23(b)	<b>Alternative method 1</b>		
	$-3 + \sqrt{25} (= 2)$ or $-3 + 5 (= 2)$	B1	oe eg $5 - 3 (= 2)$ or $2 + 3 = 5$
	<b>Alternative method 2</b>		
	$(y + 3)^2 = 25$ and $y = 2$ or $y + 3 = 5$ and $y = 2$ or $(2 + 3)^2 = 25$	B1	oe eg $(1 - 1)^2 + (y + 3)^2 = 25$ and $y = 2$
	<b>Additional Guidance</b>		
	$(1, -3) + (0, 5) = (1, 2)$ so $y = 2$		B0
	Allow $-3 +$ radius of 5		
	$2 = 0x + c$ $c = 2$ so $y = 2$		B0

Q	Answer	Mark	Comments
23(c)	<b>Alternative method 1 Using equation <math>PR</math></b>		
	$\frac{-7 - \text{their } -3}{4 - \text{their } 1}$ or $-\frac{4}{3}$	M1	oe grad $PC$ their $-3$ and their $1$ from (a)
	$-1 \div \text{their } -\frac{4}{3}$ or $\frac{3}{4}$	M1	oe grad $PR$ their $-\frac{4}{3}$ must be a value (gradient $PR =$ ) $\frac{3}{4}$ is M2
	$2 - -7 = \text{their } \frac{3}{4}(x - 4)$	M1dep	oe equation $PR$ with $y = 2$ substituted eg $2 = \frac{3}{4}x - 10$ dep on 2nd M1
	16	A1ft	only ft their $-3$ and their $1$ from (a)
	<b>Alternative method 2 Using <math>RC^2 = CP^2 + PR^2</math> or <math>PR^2 = QR^2</math> with <math>R(x, 2)</math></b>		
	$(x - \text{their } 1)^2 + (2 - \text{their } -3)^2$ $= (2 - \text{their } -3)^2 + (x - 4)^2 + (2 - -7)^2$	M1	oe eg $(x - 1)^2 = (x - 4)^2 + (2 - -7)^2$ their $-3$ and their $1$ from (a)
	$x^2 - 2x + 1 + 25$ $= 25 + x^2 - 8x + 16 + 81$	M1dep	oe brackets expanded
	$96 = 6x$ or $96 \div 6$	M1dep	oe linear equation or calculation dep on M2
	16	A1ft	only ft their $-3$ and their $1$ from (a)

Mark scheme and Additional Guidance continue on the next three pages



Q	Answer	Mark	Comments
23(c) cont	<b>Alternative method 3 Using equation CR</b>		
	$\frac{-7-2}{4 - \text{their } 1}$ or $-3$	M1	oe grad $PQ$ their 1 from (a)
	$-1 \div \text{their } -3$ or $\frac{1}{3}$	M1	oe grad $CR$ their $-3$ must be a value (gradient $CR =$ ) $\frac{1}{3}$ is M2
	$2 - \text{their } -3 = \text{their } \frac{1}{3}(x - \text{their } 1)$	M1dep	oe equation $CR$ with $y = 2$ substituted eg $2 = \frac{1}{3}x - \frac{10}{3}$ dep on 2nd M1
	16	A1ft	only ft their $-3$ and their 1 from (a)
	<b>Alternative method 4 Using equation MR where M is the midpoint of PQ</b>		
	$\frac{-7-2}{4 - \text{their } 1}$ or $-3$	M1	oe grad $PQ$ their 1 from (a)
	$-1 \div \text{their } -3$ or $\frac{1}{3}$	M1	oe grad $MR$ their $-3$ must be a value (gradient $MR =$ ) $\frac{1}{3}$ is M2
	$\left(\frac{4 + \text{their } 1}{2}, \frac{-7 + 2}{2}\right)$ or $(2.5, -2.5)$ and $2 - \text{their } -2.5 = \text{their } \frac{1}{3}(x - \text{their } 2.5)$	M1dep	oe midpoint of $PQ$ and equation $MR$ with $y = 2$ substituted eg $2 = \frac{1}{3}x - \frac{10}{3}$ dep on 2nd M1
	16	A1ft	only ft their 1 from (a)

Mark scheme and Additional Guidance continue on the next two pages

Q	Answer	Mark	Comments
<b>23(c) cont</b>	<b>Alternative method 5 Using equation <math>MC</math> where <math>M</math> is the midpoint of <math>PQ</math></b>		
	$\left(\frac{4 + \text{their } 1}{2}, \frac{-7 + 2}{2}\right) \text{ or } (2.5, -2.5)$	M1	oe midpoint of $PQ$ their 1 from (a)
	$\frac{\text{their } -3 - \text{their } -2.5}{\text{their } 1 - \text{their } 2.5} \text{ or } \frac{1}{3}$	M1dep	oe grad $MC$
	$2 - \text{their } -3 = \text{their } \frac{1}{3}(x - \text{their } 1)$ or $2 - \text{their } -2.5 = \text{their } \frac{1}{3}(x - \text{their } 2.5)$	M1dep	oe equation $MC$ with $y = 2$ substituted eg $2 = \frac{1}{3}x - \frac{10}{3}$ dep on M2
	16	A1ft	only ft their $-3$ and their 1 from (a)
	<b>Alternative method 6 Using trigonometry where <math>M</math> is the midpoint of <math>PQ</math></b>		
	$(QM =) \frac{1}{2} \sqrt{(4 - \text{their } 1)^2 + (-7 - 2)^2}$ or $\frac{1}{2} \sqrt{90}$ or 4.74...	M1	
	$\sin^{-1}\left(\frac{\text{their } 4.74\dots}{5}\right)$ or (angle $QCM =$ ) 71.5... or 71.6	M1dep	oe angle $QCM$
	$\tan(\text{their } 71.5\dots) = \frac{x - \text{their } 1}{5}$	M1dep	using triangle $QCR$
	16	A1ft	only ft their 1 from (a)

**Additional Guidance is on the next page**

Q	Answer	Mark	Comments
---	--------	------	----------

23(c) cont	Additional Guidance		
	Allow (16, ...) to imply answer 16		
	Alt 1 $-\frac{4}{3}x$ is M0 unless recovered		
	(a) (1, -2) grad $PC = -\frac{5}{3}$ grad $PR = \frac{3}{5}$ Answer 19 (3rd M1 can be implied by A1ft answer)		M1M1 M1A1ft

24	$3x^4$ or $4x^3$	M1	oe eg $5 \times \frac{3}{5}x^{5-1}$
	$3x^4 + 4x^3$	A1	
	$x^3(3x + 4) (= 0)$	M1dep	allow partial factorisation of their $3x^4 + 4x^3$ if at least x is taken as a factor ft their two terms if M1 scored
	$x^3(3x + 4) (= 0)$ and (x =) 0 and (x =) $-\frac{4}{3}$ with no other solutions	A1	allow partial factorisation if at least x is taken as a factor
	Additional Guidance		
	$3x^4 + 4x^3 = 0$ $x = 0$ and $x = -\frac{4}{3}$		M1A1 M0A0
	Condone $y = 3x^4 + 4x^3$		M1A1
	Ignore higher derivatives		
	Condone (0, ...) and $(-\frac{4}{3}, \dots)$ for (x =) 0 and (x =) $-\frac{4}{3}$		
	Allow $-1.33\dots$ for $-\frac{4}{3}$ (ignore any incorrect conversion attempt after $-\frac{4}{3}$ seen)		

Q	Answer	Mark	Comments
<b>25</b>	<b>Alternative method 1</b>		
	$(-c)^3 - 10(-c) - c (= 0)$ or $-c^3 + 10c - c (= 0)$ or $-c^3 + 9c (= 0)$	M1	oe
	$c(9 - c^2) (= 0)$ or $c(3 + c)(3 - c) (= 0)$ or $c^2 = 9$	M1dep	oe factorised expression or quadratic equation
	3 with no other value(s)	A1	SC2 answer 3 with one or both of -3 and 0 and no other value
	<b>Alternative method 2</b>		
	$(x + c)(x^2 - cx - 1)$	M1	
	$-1 - c^2 = -10$	M1dep	oe quadratic equation
	3 with no other value(s)	A1	SC2 answer 3 with one or both of -3 and 0 and no other value
	<b>Additional Guidance</b>		
	$(-3)^3 - 10(-3) - 3 = 0$ and Answer 3 (no part marks)		M2A1
	$(-3)^3 - 10(-3) - -3 = 0$ and Answer 3		Zero
	$3^3 - 10(3) - -3 = 0$ and Answer 3		Zero
	Answer 3 with no incorrect working		M2A1
	Allow recovery of missing brackets		

Q	Answer	Mark	Comments
26	<b>Alternative method 1</b>		
	$(x + 3)^2 \dots$	M1	
	$(x + 3)^2 - 3^2 - a$ or $(x + 3)^2 - 3^2 \geq a$ or $(x + 3)^2 \geq a + 3^2$	M1dep	oe expression or inequality eg $(x + 3)^2 \geq 9 + a$ allow $\geq$ to be any inequality symbol or = eg allow $(x + 3)^2 - 9 = a$ implies M2
	$-3^2 - a \geq 0$ or $-3^2 - a > 0$	M1dep	oe inequality eg $-9 - a \geq 0$ or $-9 - a > 0$ or $a < -9$ implies M3
	$a \leq -9$ or $-9 \geq a$	A1	SC1 $x^2 + 6x - a \geq 0$ oe inequality (may be seen in working lines)
	<b>Alternative method 2</b>		
	$2x + 6 = 0$	M1	must have = 0
	(minimum at) $x = -3$	M1dep	implies M2 $x = -3$ must be the only value or be clearly chosen
	$(-3)^2 + 6 \times (-3) - a \geq 0$ or $(-3)^2 + 6 \times (-3) - a > 0$	M1dep	oe inequality eg $9 - 18 - a \geq 0$ or $9 - 18 - a > 0$ or $a < -9$ implies M3
	$a \leq -9$ or $-9 \geq a$	A1	SC1 $x^2 + 6x - a \geq 0$ oe inequality (may be seen in working lines)

Mark scheme and Additional Guidance continue on the next page

Q	Answer	Mark	Comments
<b>26 cont</b>	<b>Alternative method 3</b>		
	$6^2 - 4 \times 1 \times -a$	M1	$b^2 - 4ac$ must be selected if seen in quadratic formula
	$6^2 - 4 \times 1 \times -a \leq 0$ or $6^2 - 4 \times 1 \times -a < 0$	M1dep	oe inequality implies M2
	$36 + 4a \leq 0$ or $36 + 4a < 0$	M1dep	oe inequality eg $4a \leq -36$ implies M3
	$a \leq -9$ or $-9 \geq a$	A1	SC1 $x^2 + 6x - a \geq 0$ oe inequality (may be seen in working lines)
	<b>Additional Guidance</b>		
	Alt 1 2nd M1 Any inequality symbol or = allowed 3rd M1 Only the inequality symbols shown are allowed (do not allow =)		
Allow $(x + 3)(x + 3)$ for $(x + 3)^2$			

Q	Answer	Mark	Comments
27	<b>Alternative method 1</b>		
	Shows substitution of a value of $x < -2$ into $\frac{dy}{dx}$ and shows substitution of a value of $x > -2$ into $\frac{dy}{dx}$	M1	eg $(-3 + 2)^6 + (-3 + 2)^4$ and $(-1 + 2)^6 + (-1 + 2)^4$ allow $(-1)^6 + (-1)^4$ with $x = -3$ stated and $(1)^6 + (1)^4$ with $x = -1$ stated
	Evaluates both correctly or states that each is positive with M1 seen	M1dep	eg $(-3 + 2)^6 + (-3 + 2)^4 = 2$ and $(-1 + 2)^6 + (-1 + 2)^4 = 2$ allow $(-1)^6 + (-1)^4 = 2$ with $x = -3$ stated and $(1)^6 + (1)^4 = 2$ with $x = -1$ stated
Statement with M2 seen	A1	eg either side of $P \frac{dy}{dx} > 0$ with M2 seen  SC2 states two values of $x$ (one $< -2$ and one $> -2$ ) and shows the correct value of $\frac{dy}{dx}$ for each and makes a statement  SC1 states two values of $x$ (one $< -2$ and one $> -2$ ) and shows the correct value of $\frac{dy}{dx}$ for each	


Mark scheme and Additional Guidance continue on the next three pages

Q	Answer	Mark	Comments
<b>27 cont</b>	<b>Alternative method 2</b>		
	$x < -2$ and $(-)^6 + (-)^4$ and $x > -2$ and $(+)^6 + (+)^4$	M1	allow without brackets allow less than for $<$ etc
	$x < -2$ and $(-)^6 + (-)^4 > 0$ and $x > -2$ and $(+)^6 + (+)^4 > 0$	M1dep	allow without brackets allow $= +$ for $> 0$
	Statement with M2 seen	A1	eg either side of $P \frac{dy}{dx} > 0$ with M2 seen  SC2 states two values of $x$ (one $< -2$ and one $> -2$ ) and shows the correct value of $\frac{dy}{dx}$ for each and makes a statement  SC1 states two values of $x$ (one $< -2$ and one $> -2$ ) and shows the correct value of $\frac{dy}{dx}$ for each

**Additional Guidance is on the next two pages**



Q	Answer	Mark	Comments
---	--------	------	----------

<b>Additional Guidance</b>									
For A1 a clear statement is needed after M2 scored									
Examples of acceptable statements with M2 seen eg1 For $x < -2$ gradient is + and for $x > -2$ gradient is + eg2 To the left of $P$ $m > 0$ To the right of $P$ $m > 0$ eg3 (When both of their substitutions correctly evaluate to the same value) They are the same <b>positive</b> value eg4 <b>Both</b> gradients are the same sign eg5 $m$ is + <b>both</b> times eg6 Gradient is <b>always</b> positive (apart from at $P$ ) eg7 <b>Function (or curve)</b> is increasing (either side of $P$ )									
Allow a statement to be made using a diagram with M2 seen eg accept for eg2 above									
Allow a statement to be made using a table with M2 seen eg accept for eg1 above		<table border="1" data-bbox="641 1348 890 1482"> <tbody> <tr> <td style="text-align: center;">-3</td> <td style="text-align: center;">-2</td> <td style="text-align: center;">-1</td> </tr> <tr> <td style="text-align: center;">+</td> <td style="text-align: center;">0</td> <td style="text-align: center;">+</td> </tr> </tbody> </table>		-3	-2	-1	+	0	+
-3	-2	-1							
+	0	+							
When both of their substitutions correctly evaluate to the same positive value condone for the statement with M2 seen Gradients are the same (implies both positive)									
Do not accept for the statement eg1 Gradient is increasing eg2 $m$ is positive eg3 Gradient is positive eg4 $P$ is a point of inflection									

**Additional Guidance continues on the next page**

Q	Answer	Mark	Comments
---	--------	------	----------

<b>Additional Guidance</b>								
<b>27 cont</b>	Allow gradient or m for $\frac{dy}{dx}$							
	For evaluations allow rounding or truncating to 1 sf or better							
	Ignore higher derivatives							
	Ignore substitution of $x = -2$							
	$x = -3$ gradient = 2 $x = -1$ gradient = 2 either side of $P$ gradient $> 0$		SC2					
	<table border="1" data-bbox="252 929 499 1064" style="display: inline-table; vertical-align: middle;"> <tr> <td style="text-align: center;">-3</td> <td style="text-align: center;">-2</td> <td style="text-align: center;">-1</td> </tr> <tr> <td style="text-align: center;">2</td> <td style="text-align: center;">0</td> <td style="text-align: center;">2</td> </tr> </table> gradient is positive both times		-3	-2	-1	2	0	2
-3	-2	-1						
2	0	2						
<table border="1" data-bbox="252 1142 499 1276" style="display: inline-table; vertical-align: middle;"> <tr> <td style="text-align: center;">-3</td> <td style="text-align: center;">-2</td> <td style="text-align: center;">-1</td> </tr> <tr> <td style="text-align: center;">2</td> <td style="text-align: center;">0</td> <td style="text-align: center;">2</td> </tr> </table>		-3	-2	-1	2	0	2	SC1
-3	-2	-1						
2	0	2						