

# F321 Atoms, Bonds and Groups



Question		Expected Answers	Marks	Additional Guidance
1	(a)	<p>Mass of the <b>isotope</b> compared to 1/12th  <b>OR</b>  <b>mass</b> of the <b>atom</b> compared to 1/12th ✓</p> <p>(the mass of a) carbon-12 <b>OR</b> <math>^{12}\text{C}</math> (atom) ✓</p>	2	<p><b>IGNORE</b> Reference to average <b>OR</b> weighted mean            (i.e. correct definition of relative atomic mass will score both marks)</p> <p><b>ALLOW</b> mass of a <b>mole</b> of the isotope/atom with 1/12th the mass of a <b>mole OR 12 g</b> of carbon-12 for two marks.</p> <p><b>ALLOW 2 marks for:</b>            'Mass of the isotope <b>OR</b> mass of the atom compared to <math>^{12}\text{C}</math> atom given a mass of 12.0'            i.e. 'given a mass of 12' <b>OR</b> C12 is 12 communicates the same idea as 1/12th.'</p> <p><b>ALLOW</b> 12C <b>OR</b> C12</p> <p><b>ALLOW 2 marks for:</b>  <math display="block">\frac{\text{mass of the isotope}}{\text{mass of 1/12th mass of carbon - 12}}</math>           i.e. fraction is equivalent to 'compared to'</p> <p><b>ALLOW 1 mark for</b> a mix of mass of atom and mass of mole of atoms, i.e. 'mass of the isotope/mass of an atom compared with 1/12th the mass of a <b>mole OR 12 g</b> of carbon-12.'</p> <p><b>DO NOT ALLOW</b> mass of 'ions' <b>OR</b> mass of element</p>
	(b)	<p><math display="block">\frac{(151 \times 47.77) + (153 \times 52.23)}{100}</math></p> <p><b>OR</b>            72.1327 + 79.9119  <b>OR</b>            152.0446 (calculator value) ✓  <math>A_r = 152.04</math> ✓</p>	2	<p><b>ALLOW</b> Correct answer for two marks</p> <p><b>ALLOW</b> One mark for ECF from transcription error in first sum provided final answer is to 2 decimal points and is to between 151 and 153 and is a correct calculation of the transcription</p>


Question		Expected Answers	Marks	Additional Guidance
	(c) (i)	$^{153}\text{Eu}$ has (2) more neutrons <b>OR</b> $^{153}\text{Eu}$ has 90 neutrons <b>AND</b> $^{151}\text{Eu}$ has 88 neutrons ✓	1	<b>ALLOW</b> There are a different number of neutrons <b>IGNORE</b> Correct references to protons / electrons <b>DO NOT ALLOW</b> Incorrect references to protons / electrons
	(ii)	(It has the) same number of protons <b>AND</b> electrons <b>OR</b> Both have 63 protons and 63 electrons ✓	1	<b>ALLOW</b> Same number of protons <b>AND</b> same electron configuration <b>DO NOT ALLOW</b> 'Same number of protons' without reference to electrons (and vice versa)

Question	Expected Answers	Marks	Additional Guidance
(d)	<p>Xe has a bigger atomic radius <b>OR</b> Xe has more shells ✓</p> <p>Xe has <b>more</b> shielding ✓</p> <p>The nuclear attraction decreases  <b>OR</b> Outermost electrons of Xe experience less attraction (to nucleus)  <b>OR</b> Increased shielding / distance outweighs the increased nuclear charge ✓  ORA throughout</p>	3	<p><b>ALLOW</b> Xe has more energy levels  <b>ALLOW</b> Xe has electrons in higher energy level  <b>ALLOW</b> Xe has electrons further from nucleus  <b>IGNORE</b> Xe has more orbitals <b>OR</b> more sub-shells  <b>DO NOT ALLOW</b> 'different shell' or 'new shell'</p> <p><b>ALLOW More</b> screening  There must be a clear comparison ie <b>more</b> shielding <b>OR</b> <b>increased</b> shielding.  i.e. <b>DO NOT ALLOW</b> Xe 'has shielding'  <b>ALLOW</b> Xe has <b>more</b> electron repulsion from inner shells</p> <p><b>ALLOW</b> Xe has less nuclear pull  <b>IGNORE</b> Xe has less effective nuclear charge  <b>DO NOT ALLOW</b> nuclear charge for nuclear attraction</p>
	<b>Total</b>	<b>9</b>	

Question			Expected Answers	Marks	Additional Guidance
2	(a)	(i)	The H <sup>+</sup> ion in an (nitric) acid has been replaced by a metal ion <b>OR</b> by a Ca <sup>2+</sup> ion ✓	1	<b>DO NOT ALLOW</b> it has been produced by the reaction of an acid and a base as this is stated in the question.  <b>IGNORE</b> references to replacement by NH <sub>4</sub> <sup>+</sup> ions or positive ions. <b>ALLOW</b> H <b>OR</b> Hydrogen for H <sup>+</sup> ; <b>DO NOT ALLOW</b> Hydrogen atoms <b>ALLOW</b> Ca <b>OR</b> Calcium for Ca <sup>2+</sup> . <b>DO NOT ALLOW</b> Calcium atoms <b>ALLOW</b> 'metal' for 'metal ion'
		(ii)	2HNO <sub>3</sub> (aq) + Ca(OH) <sub>2</sub> (aq) → Ca(NO <sub>3</sub> ) <sub>2</sub> (aq) + 2H <sub>2</sub> O(l) Formulae ✓ Balance <b>AND</b> states ✓	2	<b>ALLOW</b> multiples <b>ALLOW</b> (aq) <b>OR</b> (s) for Ca(OH) <sub>2</sub>
		(iii)	Accepts a <b>proton OR</b> accepts H <sup>+</sup> ✓	1	<b>ALLOW</b> H <sup>+</sup> + OH <sup>-</sup> → H <sub>2</sub> O <b>ALLOW</b> OH <sup>-</sup> reacts with H <sup>+</sup> <b>OR</b> OH <sup>-</sup> takes H <sup>+</sup> <b>ALLOW</b> OH <sup>-</sup> 'attracts' H <sup>+</sup> if 'to form water' is seen  <b>DO NOT ALLOW</b> OH <sup>-</sup> neutralises H <sup>+</sup> ('neutralises' is in the question)
	(b)	(i)	Calculates correctly $\frac{0.0880 \times 25.0}{1000} = 2.20 \times 10^{-3}$ mol <b>OR</b> 0.00220 mol ✓	1	<b>ALLOW</b> 0.0022 <b>OR</b> $2.2 \times 10^{-3}$ mol
		(ii)	Calculates correctly $\frac{0.00220}{2} = 1.10 \times 10^{-3}$ mol <b>OR</b> 0.00110 mol ✓	1	<b>ALLOW</b> 0.0011 <b>OR</b> $1.1 \times 10^{-3}$ mol  <b>ALLOW</b> ECF for answer (i)/2 as calculator value or correct rounding to 2 significant figures or more but ignore trailing zeroes
		(iii)	$\frac{0.00110 \times 1000}{17.60} = 0.0625$ mol dm <sup>-3</sup> <b>OR</b> $6.25 \times 10^{-2}$ mol dm <sup>-3</sup> ✓	1	<b>ALLOW</b> 0.063 <b>OR</b> $6.3 \times 10^{-2}$ mol dm <sup>-3</sup>  <b>ALLOW</b> ECF for answer (ii) × 1000/17.60 <b>OR</b> ECF from (i) for answer (i)/2 × 1000/17.60 as calculator value or correct rounding to 2 significant figures or more but ignore trailing zeroes

	(c)	(i)	(The number of) Water(s) of crystallisation ✓	1	<b>IGNORE</b> hydrated <b>OR</b> hydrous
		(ii)	142.1 ✓  $x = \frac{(322.1 - 142.1)}{18.0} = 10$ ✓	2	<b>ALLOW</b> 142 <b>ALLOW</b> $M_r$ expressed as a sum  <b>ALLOW</b> ECF from incorrect $M_r$ and $x$ is <b>calculated correctly</b>  <b>ALLOW</b> ECF values of $x$ from nearest whole number to calculator value  <b>ALLOW</b> 2 marks if final answer is 10 <b>without any working</b>
			<b>Total</b>	<b>10</b>	


Question			Expected Answers	Marks	Additional Guidance
3	(a)	(i)	(Electrostatic) <b>attraction</b> between oppositely charged ions. ✓	1	<b>IGNORE</b> force <b>IGNORE</b> references to transfer of electrons <b>MUST</b> be ions, not particles
		(ii)	Mg shown with either 8 or 0 electrons <b>AND</b> S shown with 8 electrons <b>with</b> 2 crosses and 6 dots (or vice versa) ✓  Correct charges on both ions ✓  	2	Mark charges on ions and electrons independently <b>For first mark</b> , if 8 electrons are shown around the Mg then 'extra electrons' around S must match the symbol chosen for electrons around Mg  Shell circles not required  <b>IGNORE</b> inner shell electrons  Brackets are not required
	(b)	(i)	Electron pairs in covalent bonds shown correctly using dots and crosses in a molecule of the F <sub>2</sub> O ✓  Lone pairs correct on O and both F atoms ✓  	2	Must be 'dot-and-cross' circles for outer shells <b>NOT</b> needed <b>IGNORE</b> inner shells  Non-bonding electrons of O do not need to be shown as pairs  Non-bonding electrons of F do not need to be shown as pairs
		(ii)	Predicted bond angle 104–105°. ✓  There are 2 bonded pairs and 2 lone pairs ✓ Lone pairs repel more than bonded pairs ✓	3	<b>ALLOW</b> 103–105° (103° is the actual bond angle)  <b>ALLOW</b> responses equivalent to second marking point. e.g. There are 4 pairs of electrons and 2 of these are lone pairs <b>ALLOW</b> 'bonds' for 'bonded pairs' <b>DO NOT ALLOW</b> 'atoms repel' <b>DO NOT ALLOW</b> electrons repel <b>ALLOW</b> LP for 'lone pair' <b>ALLOW</b> BP for bonded pair <b>ALLOW</b> LP repel more if bonded pairs have already been mentioned

Question	Expected Answers	Marks	Additional Guidance
(c) (i)	<p>(At least) two <math>\text{NH}_3</math> molecules with correct dipole shown with at least one H with <math>\delta^+</math> and one N with <math>\delta^-</math> ✓</p> <p>(Only) one hydrogen bond from N atom on one molecule to a H atom on another molecule ✓</p> <p>Lone pair shown on the N atom and hydrogen bond must hit the lone pair ✓</p> 	3	<p><b>DO NOT ALLOW</b> first mark for ammonia molecules with incorrect lone pairs</p> <p><b>DO NOT ALLOW</b> first mark if <math>\text{H}_2\text{O}</math>, <math>\text{NH}_2</math> or <math>\text{NH}</math> is shown</p> <p><b>ALLOW</b> hydrogen bond need not be labelled as long as it clear the bond type is different from the covalent N–H bond</p> <p><b>ALLOW</b> a line (i.e. looks like a covalent bond) as long as it is labelled 'hydrogen bond'</p> <p><b>ALLOW</b> 2-D diagrams</p> <p><b>ALLOW</b> two marks if water molecules are used. One awarded for a correct hydrogen bond and one for the involvement of lone pair</p>
(ii)	<p>Liquid <math>\text{H}_2\text{O}</math> is denser than solid ✓</p> <p>In solid state <math>\text{H}_2\text{O}</math> molecules are held apart by hydrogen bonds <b>OR</b> ice has an open lattice ✓</p> <p><b>OR</b></p> <p><math>\text{H}_2\text{O}</math> has a relatively high boiling point <b>OR</b> melting point ✓</p> <p>(relatively strong) hydrogen bonds need to be broken <b>OR</b> a lot of energy is needed to overcome hydrogen bonds <b>OR</b> hydrogen bonds are strong ✓</p>	2	<p>ORA</p> <p><b>ALLOW</b> ice floats for first mark</p> <p><b>ALLOW</b> higher melting <b>OR</b> boiling point than expected</p> <p><b>DO NOT ALLOW</b> <math>\text{H}_2\text{O}</math> has a high melting / boiling point</p> <p><b>ALLOW</b> other properties caused by hydrogen bonding not mentioned within the specification</p> <p>E.g. high surface tension – strong hydrogen bonds on the surface</p>
	<b>Total</b>	<b>13</b>	

Question		Expected Answers	Marks	Additional Guidance
4	(a)	<p><i>Advantage</i> removes or kills bacteria <b>OR</b> kills germs <b>OR</b> kills micro-organisms <b>OR</b> make it safe to drink <b>OR</b> sterilises water <b>OR</b> disinfects water ✓</p> <p><i>Disadvantage</i> it is toxic <b>OR</b> poisonous <b>OR</b> could form chlorinated hydrocarbons ✓</p>	2	<p><b>ALLOW</b> to make water potable <b>IGNORE</b> virus <b>IGNORE</b> 'purifies water' <b>DO NOT ALLOW</b> 'antiseptic'</p> <p><b>ALLOW forms</b> carcinogens <b>OR</b> forms toxins <b>IGNORE</b> harmful <b>DO NOT ALLOW</b> 'it causes cancer' <b>DO NOT ALLOW</b> "It kills you"</p>
	(b)	$3d^{10} 4s^2 4p^5$ ✓	1	<p><b>ALLOW</b> <math>4s^2 3d^{10} 4p^5</math> <b>ALLOW</b> subscripts or <math>3D^{10}</math> <b>ALLOW</b> answers with <math>1s^2 2s^2 2p^6 3s^2 3p^6</math> appearing twice</p>
	(c) (i)	$Cl_2 + 2Br^- \rightarrow Br_2 + 2Cl^-$ ✓	1	<p><b>IGNORE</b> state symbols <b>ALLOW</b> any correct multiple including fractions</p>
	(ii)	Yellow / orange / red / brown ✓	1	<b>ALLOW</b> any combination of these, but no others
	(d) (i)	Disproportionation ✓	1	<p><b>ALLOW</b> versions which sound the same</p> <p><b>DO NOT ALLOW</b> disproportional <b>OR</b> disproportionate <b>OR</b> disproportion</p>
	(ii)	<p><math>Cl_2 + 2NaOH \rightarrow NaClO + NaCl + H_2O</math> ✓</p> <p><math>3Cl_2 + 6NaOH \rightarrow NaClO_3 + 5NaCl + 3H_2O</math></p> <p><math>Cl_2</math> and NaOH as reactants <b>AND</b> <math>NaClO_3</math> and NaCl as products ✓</p> <p>Rest of the equation ✓</p>	3	<p><b>ALLOW</b> multiples for either equation</p> <p><b>ALLOW</b> <math>3Cl_2 + 6NaOH \rightarrow 2NaClO_3 + 4NaCl + 3H_2</math></p>
	(iii)	$NaClO_4$ ✓	1	<b>ALLOW</b> $Na_3ClO_5$ etc
<b>Total</b>			<b>10</b>	



Question			Expected Answers	Marks	Additional Guidance
5	(a)	(i)	Potassium <b>AND</b> argon ✓	1	<b>ALLOW</b> K and Ar
		(ii)	They are arranged in increasing atomic number <b>OR</b> Neither would show properties <b>OR</b> trends of rest of group <b>OR</b> Neither would show properties <b>OR</b> trends of rest of period <b>OR</b> They are arranged by electron configuration ✓	1	<b>ALLOW</b> any correct property difference e.g. This would place a reactive metal in the same group as noble gases  <b>ALLOW</b> they do not fit in with the rest of the group
	(b)	(i)	$2\text{Mg} + \text{O}_2 \rightarrow 2\text{MgO}$ ✓	1	<b>ALLOW</b> multiples. Correct species must be seen <b>IGNORE</b> state symbols
		(ii)	Fizzes <b>OR</b> bubbles <b>OR</b> gas produced <b>OR</b> effervescing ✓  Mg dissolves <b>OR</b> Mg disappears <b>OR</b> a solution is formed ✓	2	<b>DO NOT ALLOW</b> 'carbon dioxide gas produced' <b>DO NOT ALLOW</b> 'hydrogen produced' without 'gas'  <b>ALLOW</b> 'it for Mg' <b>IGNORE</b> Mg reacts <b>IGNORE</b> temperature change <b>IGNORE</b> steam produced
		(iii)	Quicker <b>OR</b> more vigorous <b>OR</b> gets hotter	1	<b>MUST</b> be a comparison of a reaction observation, not just 'more reactive'  <b>ALLOW</b> any comparison of greater rate including more bubbles etc. <b>DO NOT ALLOW</b> more gas produced

Question	Expected Answers	Marks	Additional Guidance
(c)	<p>Mg has a <b>giant</b> structure ✓</p> <p>Mg has <b>metallic</b> bonding OR description of metallic bonding as positive ions and <b>delocalised</b> electrons ✓</p> <p>(There is electrostatic attraction between) positive ions and electrons ✓</p> <p>Cl has a simple molecular <b>OR</b> simple covalent (lattice) ✓</p> <p>Cl has van der Waals' forces (between molecules) <b>OR</b> Cl has instantaneous dipole–induced dipoles <b>OR</b> temporary dipole–temporary dipole ✓</p>	6	<p><b>Metallic OR delocalised</b> seen spelt correctly at least <b>ONCE</b></p> <p><b>DO NOT ALLOW</b> as label nuclei <b>OR</b> protons for positive ions</p> <p><b>ALLOW</b> labelled diagram of metallic bonding for second and third marks</p>  <p>The diagram shows a lattice of positive ions (red spheres) with delocalised electrons (small black dots) between them. Arrows point from the labels 'positive ions' and 'delocalised electrons' to the respective parts of the diagram.</p> <p>Lattice must have at least two rows of positive ions. If a Mg ion is shown it must correct charge</p> <p><b>ALLOW</b> for labels: + ions, positive ions, cations</p> <p><b>DO NOT ALLOW</b> as label nuclei <b>OR</b> protons for positive ions</p> <p><b>ALLOW</b> e<sup>-</sup> or e as label for electron</p> <p><b>DO NOT ALLOW</b> '-' without label for electron</p> <p><b>Covalent OR molecule OR molecular</b> seen spelt correctly at least <b>ONCE</b></p> <p><b>ALLOW</b> Cl is a (covalent) <b>molecule</b></p> <p><b>IGNORE</b> Cl has intermolecular bonding</p>

		<p>van der Waals' forces are weak <b>and</b> metallic bonds are strong  <b>OR</b>  van der Waals' forces are weaker than metallic bonds  <b>OR</b>  Less energy is needed to overcome van der Waals' than metallic bonds ✓</p>		<p><b>ALLOW</b> ECF from incorrect descriptions of giant structure with strong bonds; e.g. Mg has giant ionic structure  <b>ALLOW</b> ECF from any incorrect intermolecular forces e.g. permanent dipole –dipole from marking point 5    <b>ALLOW</b> vdW easier to break  ORA</p>
(d)	(i)	<p>O goes from –2 to 0 ✓    N goes from +5 to +4 ✓    N is reduced <b>AND</b> O is oxidised ✓</p>	3	<p>Oxidation numbers may be seen with equation    Third mark is dependent upon seeing a reduction in oxidation number of N and an increase in oxidation number of O    <b>ALLOW</b> ECF for third mark for N is oxidised <b>and</b> O is reduced if incorrect oxidation numbers support this    <b>IGNORE</b> references to strontium  <b>IGNORE</b> references to electron loss <b>OR</b> gain    <b>DO NOT ALLOW</b> 'One increases and one decreases'</p>

	<b>(d)</b>	<b>(ii)</b>	<p>Calculates correctly:  Mol of <math>\text{Sr}(\text{NO}_3)_2 = \frac{5.29}{211.6} = 0.0250 \checkmark</math></p> <p>Calculates correctly:  Mol of gas = <math>5/2 \times 0.0250 = 0.0625 \checkmark</math></p> <p>Calculates correctly:  Volume of gas = <math>24.0 \times 0.0625 = 1.50 \text{ dm}^3 \checkmark</math></p>	<b>3</b>	<p><b>ALLOW</b> 0.025</p> <p><b>ALLOW</b> ECF for first answer <math>\times 2.5</math> as calculator value or correct rounding to 2 significant figures or more but ignore trailing zeroes</p> <p><b>ALLOW</b> ECF for second answer <math>\times 24(.0)</math> as calculator value or correct rounding to 2 significant figures or more but ignore trailing zeroes</p> <p><b>DO NOT ALLOW</b> ECF of first answer <math>\times 24(.0)</math> (which gives <math>0.6(0) \text{ dm}^3</math>) as this has not measured the volume of any gas, simply <math>0.0250 \text{ mol}</math> of solid <math>\text{Sr}(\text{NO}_3)_2</math> converted into a gas  i.e. This answer would give <b>one</b> mark</p> <p><b>ALLOW</b> <math>1.5 \text{ dm}^3</math></p> <p><b>ALLOW</b> ECF producing correct volume of <math>\text{NO}_2</math> only  i.e. <math>1.2(0) \text{ dm}^3</math> would give <b>two</b> marks</p> <p><b>OR</b></p> <p><b>ALLOW</b> ECF producing correct volume of <math>\text{O}_2</math> only  i.e. <math>0.3(0) \text{ dm}^3</math> would give <b>two</b> marks</p>
<b>Total</b>			<b>18</b>		