

Mathematics

Advanced Subsidiary GCE

Unit **4728**: Mechanics 1

Mark Scheme for June 2011

OCR (Oxford Cambridge and RSA) is a leading UK awarding body, providing a wide range of qualifications to meet the needs of pupils of all ages and abilities. OCR qualifications include AS/A Levels, Diplomas, GCSEs, OCR Nationals, Functional Skills, Key Skills, Entry Level qualifications, NVQs and vocational qualifications in areas such as IT, business, languages, teaching/training, administration and secretarial skills.

It is also responsible for developing new specifications to meet national requirements and the needs of students and teachers. OCR is a not-for-profit organisation; any surplus made is invested back into the establishment to help towards the development of qualifications and support which keep pace with the changing needs of today's society.

This mark scheme is published as an aid to teachers and students, to indicate the requirements of the examination. It shows the basis on which marks were awarded by Examiners. It does not indicate the details of the discussions which took place at an Examiners' meeting before marking commenced.

All Examiners are instructed that alternative correct answers and unexpected approaches in candidates' scripts must be given marks that fairly reflect the relevant knowledge and skills demonstrated.

Mark schemes should be read in conjunction with the published question papers and the Report on the Examination.

OCR will not enter into any discussion or correspondence in connection with this mark scheme.

© OCR 2011

Any enquiries about publications should be addressed to:

OCR Publications
PO Box 5050
Annesley
NOTTINGHAM
NG15 0DL

Telephone: 0870 770 6622
Facsimile: 01223 552610
E-mail: publications@ocr.org.uk

Question			Expected Answer	Mark	Rationale/Additional Guidance
1			$R^2 = 8^2 + 15^2$ $R = 17 \text{ N}$ $\cos\theta = 15/17$ $\theta = 28.1^\circ$	M1 A1 M1 A1 [4]	Uses Pythagoras 3 squared terms, addition Uses trig appropriately and targets either angle Accept 28° , 0.49 rad
2	i	Also if in ii	$T - 0.45g = 0.45 \times 0.98$ $T = 4.85(1) \text{ N}$	M1 A1 [2]	N2L on 0.45 kg, weight - tension and $\pm 0.98\text{m}$ Not 4.9, 4.8 (4.851 is exact, but 4.85 acceptable) $\{g=9.81 \rightarrow T=4.85 \text{ or } 4.86 \text{ or better}\}$
	ii	Also If in i	$mg - 4.85(1) = 0.98\text{m}$ $m = 4.85(1)/(9.8-0.98) \text{ or } m(g - 0.98) = 4.85(1)$ $m = 0.55$ OR $0.98 = g(m-0.45)/(m+0.45)$ $m = (g+0.98)/(g-0.98) \times 0.45$ $m = 0.55$	M1 A1ft A1 [3] M1 A1 A1	N2L on Q, weight - tension, tension=T(i), and 0.98m Simplified to a single term in m, ft cv(T(i)) art 0.550 $\{g=9.81 \rightarrow m=0.55(0) \text{ or better}\}$ $a = g \times \Delta(\text{masses})/\Sigma(\text{masses})$
	iii		$v^2 = (0 +)^2 + 2 \times 0.98 \times 0.36$ $v = 0.84 \text{ ms}^{-1}$	M1 A1 [2]	Uses $v^2 = u^2 + 2as$, a not 9.8, $2as > 0$, $u = 0$ or omitted
	iv		$0 = 0.84^2 - 2 \times 9.8s$ $(s = 0.036)$ $S = 0.036 + 2 \times 0.36 = 0.756 \text{ m}$	M1 A1 A1 [3]	$0 = (cv(\text{iii}))^2 - 2gs$, or $t=cv(\text{iii})/g$ and $s = ut + \frac{1}{2}gt^2$ May be implied by final answer (eg 0.396) Must be 3 sf (exact) $\{g=9.81 \rightarrow s=0.756 \text{ or better}\}$

			Frequent mis-read "horizontal/vertical" MR version in {}		Allow all A1 marks in (i) and (ii) <i>except final A1 in (ii).</i>
3	i		$R = 0.8g - 6\cos 60$ { $R = 0.8g - 6\sin 60$ } $R = 4.84$ { $R = 2.64$ }	M1 A1 [2]	Resolves vertically, (R=) difference of 2 forces inc. component of 6 Accept 4.8 {2.6} { $g=9.81 \rightarrow R=4.848$ {2.65}; accept 4.8 {2.6 or 2.7} }
	ii		$Fr = 0.2 \times 4.84 (=0.968)$ { $Fr = 0.2 \times 2.64.. (=0.5287..)$ } $6\sin 60 - 0.968 = 0.8a$ { $6\cos 60 - 0.5287.. = 0.8a$ } $a = 5.29 \text{ ms}^{-2}$ { $a = 3.09 \text{ ms}^{-2}$ A0}	M1 M1 A1 A1 [4]	Uses $F=0.2(cv(i))$ or $F=0.2 \times (R \text{ found in (ii)})$ by a method which would be given M1 in (i) Uses N2L, 3 terms inc. component of 6 Fr need not be evaluated Accept 5.3 { $g=9.81 \rightarrow a=5.28$ {3.09 A0} Accept 5.3 {3.1 A0}
	iii		$Fr = 0.2 \times 0.8 \times 9.8 (= 1.568)$ $0.8a = -0.2 \times 0.8 \times 9.8$ $0 = 4.9 - 1.96t$ $t = 2.5 \text{ s}$	B1 M1* D*M1 A1 [4]	Uses $Fr = 0.2 \times 0.8g$ N2L, Fr only, accept use of Fr from (ii) Accept $0.8a = 0.2 \times 0.8 \times 9.8$, ($a = (-)1.96$) Accept 4.9/1.96, not $0 = 4.9 + 1.96t$ Accept art 2.50 { $g=9.81 \rightarrow t=2.50$ Accept art 2.50}
4	i		$a = 15/6$ or $d = 15/2$ $a = 2.5 \text{ ms}^{-2}$ $d = 7.5 \text{ ms}^{-2}$	M1 A1 A1 [3]	Uses $a = \text{speed change/time}$ Accept -7.5
	ii		$T = 6+11+2 (=19)$ $x = 15(11+19)/2$ or $15 \times 6/2 + 15 \times 11 + 15 \times 2/2$ $x = 225 \text{ m}$	M1 M1 A1 [3]	Accounts for totality of car journey (may be implied) Idea area = distance SR Accept $15 \times (13+17)/2$ M1M1
	iii		Walks = $20 \times (-)2 = (-)40 \text{ m}$ Jogs = $40/5 = 8 \text{ s}$ $T_s = 60 - \{(6+11+2) + 20 + 8\}$ $T_s = 13 \text{ s}$	M1 A1 M1 A1 [4]	Finds distance walked $T_s + \{(6+11+2) + 20 + 8\} = 60$, needs all time elements

5	i	$V_P = 3 - 2.5 \times 0.4 (= 2)$ $V_Q = 2.5 \times 0.4 (= 1)$ $+/- (0.5 \times 2 - 0.2 \times 1) (= +/- 0.8)$ $0.5 \times 2 - 0.2 \times 1 = 0.5v + 0.2 \times 3.2$ $(v = 0.32) 0.32 \text{ ms}^{-1} \text{ up}$	M1 A1 B1 M1 A1 [5]	Calculation of either speed, either directions, $ a =2.5$ Both magnitudes correct (disregard signs) Momentum before Uses conservation of momentum in collision (not both $v_P = 3$ and $v_Q = 0$) Accept "same", value positive
	ii	$V_Q = 3.2 - 2.5 \times 0.6 (= 1.7)$ $V_R = 2.5 \times (0.4 + 0.6) (= 2.5)$ $0.2 \times 1.7 - 0.3 \times 2.5 = (0.2 + 0.3)v$ $(v = -0.82) 0.82 \text{ ms}^{-1} \text{ down}$	M1 A1 M1 A1ft A1 [5]	Calculation of either speed with its correct time, $ a =2.5$ Both magnitudes correct (disregard signs) Uses momentum conservation in collision (not both $v_Q = 3.2$ and $v_R = 0$) LHS different signs, RHS same signs, ft cv(speeds Q, R) Value positive
6	i	"...smooth ring...", "...no friction at ring.."	B1 [1]	If a variety of reasons is offered, "smooth ring" must be the last
	ii	$T \cos \theta + 5 = T \cos(90 - \theta)$ $T \cos \theta + 5 = T \sin \theta \dots\dots\dots(a)$ $T \sin \theta + T \sin(90 - \theta) = 7$ $T \sin \theta + T \cos \theta = 7 \dots\dots\dots (b)$	M1 A1 M1 A1 [4]	"Resolves horiz" equation, needs TCorS θ , 3 terms, 2 of which are T resolved "Resolves vert" equation, needs TCorS θ , 3 terms, 2 of which are T resolved {Allow candidates solving for (iii) to begin in (ii)}
	iii	uses (b)+(a) and (b)-(a) for example $T \sin \theta = 6$ or $2T \sin \theta = 12$, $T \cos \theta = 1$ or $2T \cos \theta = 2$ $T^2 = 6^2 + 1^{(2)}$ $T = 6.08 \text{ N}$ $\tan \theta = 6/(1)$ $\theta = 80.5^\circ$ OR (b) gives $T = 7/(\sin \theta + \cos \theta)$, subs in (a) for example $12 \cos \theta = 2 \sin \theta$ then mark as 6(iii) below for D*M1 A1 D*M1 A1	M1* A1 D*M1 A1 D*M1 A1 [6] M1* A1	Attempts to solve 2 equations in 2 unknowns Both terms have values correct Accept $\sqrt{37}$, 6.1 Uses a correct trig identity Accept 81° , 1.4 rad, 1.41 rad Attempts to solve 2 equations in 2 unknowns Correct two term equation in one variable

7	i	$v = dx/dt$ $v = 0.3t^2 - 0.6t + 0.2$ $a = dv/dt$ $a = 0.6t - 0.6$	M1 A1 M1 A1ft [4]	Uses differentiation of x Uses differentiation of v Correct differentiation of candidate's v(t)
	ii	$0.6t - 0.6 = 0$ (t = 1) $x(1) = 0.1x1^3 - 0.3x1^2 + 0.2x1$ $x(1) = 0$ AG OR $0.1t^3 - 0.3t^2 + 0.2t = 0$ (t=1, and disregard others) $a(1) = 0.6x1 - 0.6$ $a(1) = 0$	M1* D*M1 A1 [3]	Attempts to solve a=0 Puts solution in x formula Attempts to solve x=0 Puts solution in a formula
	iii	$0.3t^2 - 0.6t + 0.2 = 0$ t = 0.423 s t = 1.58 s	M1 A1 A1 [3]	Attempts to solve 3 term QE v = 0, accept imperfect attempt at formula, completing square or factorisation Accept 1 - 1/√3, 0.42, 0.422, or better Accept 1 + 1/√3, 1.6, 1.57, or better
	iv	$x = \int 0.2t^2 - 0.4dt$ $x = 0.2t^3/3 - 0.4t (+k)$ $0.1t^3 - 0.3t^2 + 0.2t = 0.2t^3/3 - 0.4t (+k)$ $t^3 - 9t^2 + 18t = 0$ $t^2 - 9t + 18 = 0$ AG $(t-3)(t-6) = 0$ T = 3 s	M1* A1 D*M1 D*M1 A1 M1 A1 [7]	Uses integration, ignore omission of k $x = 2t^3/30 - 4/10 t (+k)$, or coeff t ³ 0.067 or better Equates expressions for distance 3 terms with different powers of t, no constant Explains T is non-zero, or explains division by t Tries to solve given quadratic, accept imperfect attempt at completing square, formula or factorisation, and chooses smaller positive root
		Total	[72]	

Continued

Question 6 specifies the method students are likely to find most helpful. A more sophisticated approach, resolving parallel and perpendicular to the string, is mathematically valid, and leads to correct solutions. If seen it should be marked according to the following scheme, and no penalty levied.

The final 4 marks, in 6(iii), use the same mathematics as may be encountered when choosing an unorthodox method for solving the two simultaneous equations generated in 6(ii) of the specified method (see 6(iii) above).

		<i>OR</i>		
6	i	"...smooth ring...", "...no friction at ring.."		B1 [1] If a variety of reasons is offered, "smooth ring" must be the last
	ii	$T = 7\cos\theta + 5\sin\theta$(a) $T = 7\sin\theta - 5\cos\theta$(b)	M1 A1 M1 A1 [4]	Resolves //AR (Need not create $T\cos/\sin\theta$) Resolves //BR (Need not create $T\cos/\sin\theta$)
	iii	Equating expressions for T from (a) and (b) $2\sin\theta = 12\cos\theta$ $\tan\theta = 6(/1)$ $\theta = 80.5^\circ$ $T = 7\cos 80.5 + 5\sin 80.5$ or $7\sin 80.5 - 5\cos 80.5$ $T = 6.08$	M1* A1 D*M1 A1 D*M1 A1 [6]	Attempts to solve 2 equations in 2 unknowns Correct two term equation in one variable Uses a correct trig identity Accept 81° , 1.4 rad, 1.41 rad Accept $\sqrt{37}$, 6.1

OCR (Oxford Cambridge and RSA Examinations)
1 Hills Road
Cambridge
CB1 2EU

OCR Customer Contact Centre

14 – 19 Qualifications (General)

Telephone: 01223 553998

Facsimile: 01223 552627

Email: general.qualifications@ocr.org.uk

www.ocr.org.uk

For staff training purposes and as part of our quality assurance programme your call may be recorded or monitored

Oxford Cambridge and RSA Examinations
is a Company Limited by Guarantee
Registered in England
Registered Office; 1 Hills Road, Cambridge, CB1 2EU
Registered Company Number: 3484466
OCR is an exempt Charity



OCR (Oxford Cambridge and RSA Examinations)
Head office
Telephone: 01223 552552
Facsimile: 01223 552553