

A Level Chemistry A

H432/02 Synthesis and analytical techniques

Tuesday 12 June 2018 – Afternoon

Time allowed: 2 hours 15 minutes

You must have:

 the Data Sheet for Chemistry A (sent with general stationery)

You may use:

· a scientific or graphical calculator




| First name | |
|---------------|------------------|
| Last name | |
| Centre number | Candidate number |

INSTRUCTIONS

- Use black ink. You may use an HB pencil for graphs and diagrams.
- Complete the boxes above with your name, centre number and candidate number.
- · Answer all the questions.
- Write your answer to each question in the space provided. If additional space is required, use the lined page(s) at the end of this booklet. The question number(s) must be clearly shown.
- Do **not** write in the barcodes.

INFORMATION

- The total mark for this paper is 100.
- The marks for each question are shown in brackets [].
- Quality of extended responses will be assessed in questions marked with an asterisk (*).
- · This document consists of 32 pages.



SECTION A

You should spend a maximum of 20 minutes on this section.

Write your answer to each question in the box provided.

Answer all the questions.

- 1 Which compound is used as a standard for NMR chemical shift measurements?
 - \mathbf{A} Si(CH₃)₄
 - **B** $CDCl_3$
 - \mathbf{C} D_2O
 - \mathbf{D} CCl_{Δ}

Your answer

[1]

- 2 Which compound is a secondary amide?
 - A HO—C—CH₂—N
 - $\mathbf{B} \qquad \text{HO} \mathbf{C} \mathbf{CH_2} \mathbf{N} \qquad \mathbf{CH_3} \\ \mathbf{CH_3} \qquad \mathbf{CH_3}$
 - **c** H₃C—C—N

Your answer

| _ | | | | | | | | |
|---|----------|----------|------|-----|-------|-------|---------|------------|
| 2 | \//hich | compound | dooc | not | roact | with | nuclear | shilac? |
| 3 | VVIIICII | compound | uues | HOL | reaci | willi | HUCIEUL | 기 1111년5 (|


- A CH₃CH₂CHO
- B CH₃CHCH₂
- C CH₃CH₂COCH₃
- $\mathbf{D} \quad \mathrm{CH_3CH_2CH_2C}\mathit{l}$

| Your answer | |
|-------------|--|
|-------------|--|


[1]

4 Which structure represents an alicyclic compound?







В



С



D



Your answer

[1]

| | | 4 | |
|---|------|---|-----|
| 5 | Whi | ch molecule is not planar? | |
| | Α | C_2H_4 | |
| | В | C_2H_6 | |
| | С | H ₂ CO | |
| | D | HCN | |
| | You | r answer | [1] |
| 6 | Wha | at is the number of peaks in the ¹ H NMR spectrum of HOOCCH ₂ CHOHCH ₂ COOH? | |
| | Α | 3 | |
| | В | 4 | |
| | С | 5 | |
| | D | 6 | |
| | You | r answer | [1] |
| 7 | Etha | anol can be prepared by different reactions. | |
| | Whi | ch reaction has the lowest atom economy? | |
| | Α | $C_6H_{12}O_6 \rightarrow 2C_2H_5OH + 2CO_2$ | |
| | В | $C_2H_4 + H_2O \rightarrow C_2H_5OH$ | |
| | С | $C_2H_5Br + H_2O \rightarrow C_2H_5OH + HBr$ | |
| | D | $CH_3COOC_2H_5 + H_2O \rightarrow C_2H_5OH + CH_3COOH$ | |
| | | | |

[1]

Your answer

8 The breakdown of ozone is catalysed by NO radicals.


Which equation is a propagation step in the mechanism for this process?

- $A \qquad \mathsf{NO} + \mathsf{O}_2 \to \mathsf{N} + \mathsf{O}_3$
- $\mathbf{B} \quad \mathsf{NO} + \mathsf{O_2} \to \mathsf{NO_2} + \mathsf{O}$
- $\mathbf{C} \qquad \mathsf{N} + \mathsf{O}_3 \to \mathsf{NO} + \mathsf{O}_2$
- $\mathbf{D} \quad \operatorname{NO}_2 + \operatorname{O} \rightarrow \operatorname{NO} + \operatorname{O}_2$

Your answer

[1]

9 Which compound could have produced the IR spectrum below?




- ${\bf A} \quad {\rm CH_3CH_2OH}$
- B CH₃CHOHCN
- C CH₃COOH
- D CH₃CONH₂

Your answer

[1]

10 Benzene reacts with an organic reagent in the presence of a halogen carrier to form phenylethanone.



Which organic reagent is required?


- A CH₃CH₂OH
- B CH₃CHO
- C CH₃COC1
- D CH₃COOH

| Your answer | |
|-------------|------------------|
| | ¹ [1] |

- 11 How many straight-chain structural isomers of $C_7H_{15}Cl$ contain a chiral carbon atom?
 - **A** 1
 - **B** 2
 - **C** 3
 - **D** 4

| Your answer | |
|-------------|-----|
| | [1] |

12 The mass spectrum of (CH₃)₂CHCH₂OH is shown below.



Which ion is responsible for the peak with the greatest relative intensity?

- A CHCH₂OH⁺
- B CH₃CH₂CH⁺
- \mathbf{C} $(CH_3)_2CH^+$
- D CH₃CO⁺

| Your answer | |
|-------------|--|

[1]

13 Which statement(s) support(s) the delocalised model for the structure of benzene?

- 1 All carbon–carbon bonds have the same length.
- 2 The enthalpy change of hydrogenation of benzene is less exothermic than expected.
- 3 Bromine reacts with benzene less readily than with cyclohexene.
- **A** 1, 2 and 3
- B Only 1 and 2
- C Only 2 and 3
- D Only 1

| Your answer | |
|-------------|--|
|-------------|--|

[1]

Turn over

14 A solid organic compound can be purified by recrystallisation.


Which statement(s) about recrystallisation is/are true?

- 1 The organic compound is more soluble in hot solvent.
- 2 The hot solution is cooled before the purified organic compound is collected.
- 3 The melting point of the purified organic compound is lower than the impure compound.
- **A** 1, 2 and 3
- B Only 1 and 2
- C Only 2 and 3
- D Only 1

Your answer

[1]

15 Which of the following could react with the compound below to form a carbon–carbon bond?



- 1 CH_3Cl and $AlCl_3$
- 2 KCN in ethanol
- 3 CH $_3$ OH and H $_2$ SO $_4$
- **A** 1, 2 and 3
- B Only 1 and 2
- C Only 2 and 3
- **D** Only 1

Your answer

[1]

9


BLANK PAGE

PLEASE DO NOT WRITE ON THIS PAGE

SECTION B

Answer all the questions.

- 16 This question is about reactions of organic compounds containing carbon, hydrogen and oxygen.
 - (a) A chemist investigates two reactions of alcohol **A**, shown below.



| (i) | What is the systematic name of alcohol A | ν.? | |
|-------|---|--|-----|
| (ii) | | | [1] |
| | | | [1] |
| (iii) | The chemist heats alcohol A with an adalkenes. | cid catalyst to form a mixture containing tv | vo |
| | Draw the structures of the two alkenes fo | ormed in this reaction. | |
| | | | |
| | | | |
| | | | |
| | | | |

[2]


(iv) The chemist heats alcohol A with sodium chloride and sulfuric acid.

Construct a balanced equation for this reaction. Show structures for the organic compounds in your equation. **(b)** Compound **B**, shown below, is refluxed with excess acidified potassium dichromate(VI) to form a single organic product.

Complete the equation for this reaction.

(c) The flowchart below shows some reactions of compound C.

In the boxes, draw the organic products of these reactions.



| | | 12 |
|----|-----|--|
| 17 | The | general formula of an α -amino acid is RCH(NH $_2$)COOH. |
| | (a) | The α -amino acid cysteine (R = CH $_2$ SH) shows optical isomerism. |
| | | Draw 3-D diagrams to show the optical isomers of cysteine. |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | [0] |
| | | [2] |
| | (b) | The α -amino acid lysine (R = (CH ₂) ₄ NH ₂) reacts with an excess of dilute hydrochloric acid to form a salt. |
| | | Draw the structure of the salt formed in this reaction. |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | [2] |
| | | |
| | | |
| | | |
| | | |

(c) α -Amino acids can react to form proteins.

A short section of a protein chain is shown below.


HO
$$CH_2$$
 H O CH_2 $CH_$

A student hydrolyses the protein with hot NaOH(aq).

Draw the structures of the organic products formed from this section of the protein.

- 18 This question is about aromatic compounds.
 - (a) Phenol undergoes nitration more readily than benzene.
 - (i) A student carries out the nitration of phenol with dilute nitric acid to produce 2-nitrophenol and 4-nitrophenol.

A small amount of 3-nitrophenol is also produced.



The student thought that $^{13}\mathrm{C}$ NMR spectroscopy could be used to distinguish between these three nitrophenols.

| Explain whether the student is correct. | |
|---|-----|
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | [3] |

| (ii) | Explain why phenol is nitrated more readily than benzene. |
|------|---|
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | [3] |

(b) Methylbenzene reacts with sulfur trioxide, ${\rm SO_3}$, to form ${\bf D}$, shown below.

The electrophile in this reaction is ${\rm SO}_3$.

Complete the mechanism for the formation of ${\bf D}$. Show curly arrows and the structure of the intermediate.

19 This question is about the hydrolysis of haloalkanes.

| (a) | The rate of hydrolysis of a haloalkane depends on the halogen present. |
|-----|---|
| | State and explain how the halogen in the haloalkane affects the rate of hydrolysis. |
| | |
| | |
| | |
| | |
| | [2] |

(b) Chlorocyclohexane is hydrolysed with aqueous sodium hydroxide.

Outline the mechanism for this reaction.

Show curly arrows, relevant dipoles and the products.

[3]

- (c) A student hydrolyses a haloalkane, **E**, using the following method.
 - 0.0100 mol of haloalkane **E** is refluxed with excess NaOH(aq) to form a reaction mixture containing an organic product **F**.
 - The reaction mixture is neutralised with dilute nitric acid.
 - Excess AgNO₃(aq) is added to the reaction mixture. 1.88 g of a precipitate **G** forms.

Organic product, **F**, has a molar mass of 74.0 g mol⁻¹ and has a chiral carbon atom.

(i) Draw a **labelled** diagram to show how the student would carry out the hydrolysis of haloalkane **E**.

[2]

(ii) Analyse the information to identify E, F and G.

Show your working.

20 Cinnamaldehyde and methylcinnamaldehyde are naturally occurring organic compounds.

cinnamaldehyde

methylcinnamaldehyde

| (a) | Methylcinnamaldehyde is an <i>E</i> stereoisomer. |
|-----|--|
| | Explain this statement in terms of the Cahn-Ingold-Prelog (CIP) rules. |
| | |
| | |
| | |
| | |

(b) A student plans to carry out some chemical tests on both cinnamaldehyde and methylcinnamaldehyde.


cinnamaldehyde

methylcinnamaldehyde

| (i) | Suggest a suitable chemical test to confirm that both compounds contain an unsaturated carbon chain. |
|-------|--|
| | Your answer should include the reagent and observations. |
| | |
| | [1] |
| (ii) | Describe a chemical test to confirm that both compounds contain an aldehyde functional group. |
| | Your answer should include the reagent and observations. |
| | |
| | [1] |
| (iii) | Describe a chemical test to confirm that cinnamaldehyde and methylcinnamaldehyde contain a carbonyl group. |
| | How could the products of this test be used to distinguish between the two compounds? |
| | Your answer should not include spectroscopy. |
| | |
| | |
| | |
| | |
| | |
| | |
| | [3] |

(c) The flowchart below shows some reactions starting with cinnamaldehyde.

Draw the structures of the missing organic compounds in the boxes and add the missing reagent(s) on the dotted line.



(d)* Methylcinnamaldehyde reacts with iodine monochloride, IC*l*, by electrophilic addition. The reaction produces a mixture containing two different organic products.

methylcinnamaldehyde

The electronegativity values of chlorine and iodine are given in the table below.

| Pauling electronegativity value | |
|---------------------------------|-----|
| Cl | 3.0 |
| I | 2.5 |

Outline the mechanism, using the 'curly arrow' model, for the formation of **one** of the organic products and explain which of the two possible organic products is more likely to be formed.


| In your mechanism, you can show the phenyl group as $\mathrm{C_6H_5}$. | [6] |
|---|-----|
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |

.....

| Additional answer space if required. |
|--------------------------------------|
| |
| |
| |
| |
| |
| |
| |
| |
| |


- 21 This question is about aromatic carboxylic acids and their derivatives.
 - (a) The flowchart below shows some reactions of compound ${\bf H}$.

In the boxes, draw the organic products of these reactions.



(b) Compound **H** is used in the synthesis of polymer **I**, as shown in the flowchart below.

Complete the flowchart by drawing the structure of the acyl chloride and two repeat units of polymer I, and stating the formula of the reagent(s) required for the first stage on the dotted line.




© OCR 2018 Turn over

[4]

(c) A cosmetic product containing four esters, J, K, L and M, is analysed by gas chromatography and mass spectrometry. The results are shown below.

Gas chromatogram



The numbers by the peaks are the relative molar proportions of the compounds in the mixture.

Mass spectrometry

| ester | m/z of molecular ion peak | |
|-------|---------------------------|--|
| J | 152 | |
| K | 166 | |
| L | 180 | |
| M | 180 | |


(i) The concentration of ester **K** in the cosmetic product is $9.13 \times 10^{-2} \, \mathrm{g} \, \mathrm{dm}^{-3}$.

Using the results, calculate the concentration, in $mol\,dm^{-3}$, of ester ${\bf M}$ in the cosmetic product.

Give your answer to two significant figures.

concentration of ester $\mathbf{M} = \dots \mod \mathrm{dm}^{-3}$ [2]

(ii) A general structure for esters J, L and M is shown below.



Where 'R' is an alkyl group.

Use the mass spectrometry results to deduce possible structures for esters ${\bf J},\,{\bf L}$ and ${\bf M}.$

| | I . | |
|---|------------|-------|
| | | |
| | | |
| | | |
| | | |
| | I . | |
| | I . | |
| | I . | |
| | I . | |
| | I . | |
| | I . | |
| | I . | |
| | I . | |
| | · • | |
| | I I | M |
| _ | | IVI |
| | · | 1 141 |
| , | <u> </u> | === |
| | | |
| | | |

[3]

22 The relative molecular masses and boiling points of some fuels are shown in **Table 22.1**.

| Fuel | Relative molecular mass | Boiling point/°C |
|-------------|-------------------------|------------------|
| hexane | 86 | 69 |
| pentan-1-ol | 88 | 138 |
| heptane | 100 | 98 |

Table 22.1

| (a) | Write an equation for the incomplete combustion of heptane. | |
|-----|--|-------|
| (b) | Explain the difference in the boiling points of the fuels in Table 22.1 . | . [1] |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |

- (c) Fuel additives are often used to improve the combustion of a fuel.
 - (i) Compound N is a fuel additive containing carbon, hydrogen and oxygen only.

Complete combustion of 1.71g of compound $\bf N$ produces 2.97g of ${\rm CO_2}$ and 1.62g of ${\rm H_2O}$. The relative molecular mass of compound $\bf N$ is 76.0.

Calculate the molecular formula of ${\bf N}$ and suggest a possible structure for the compound.

compound N

© OCR 2018 Turn over

[5]

(ii) Solketal has been investigated as a potential fuel additive.

Solketal is synthesised from propane-1,2,3-triol and a carbonyl compound.

Construct a balanced equation for this synthesis. Show structures for the organic compounds in your equation.

[2]

(d)* A scientist is researching compounds that might be suitable as fuel additives. One of the compounds gives the analytical results below.


Elemental analysis by mass:

C: 54.54%; H: 9.10%; O: 36.36%

Mass spectrum:

Molecular ion peak at m/z = 132.0

¹H NMR spectrum in D₂O



The numbers by the peaks are the relative peak areas.

When the spectrum is run without D_2O , there are **two** additional peaks with the same relative peak areas at 11.0 ppm and 3.6 ppm.

| Use the information provided to suggest a structure for the compound. | | | |
|---|-----|--|--|
| Show all your reasoning. | [6] | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| Additional answer space if required. | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |

ADDITIONAL ANSWER SPACE

| If additiona must be cle | I space is required, you should use the following lined page(s). arly shown in the margin(s). | The question number(s |
|--------------------------|--|-----------------------|
| | and the state of t | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | I | |
| | D | |



Copyright Information

OCR is committed to seeking permission to reproduce all third-party content that it uses in its assessment materials. OCR has attempted to identify and contact all copyright holders whose work is used in this paper. To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced in the OCR Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download from our public website (www.ocr.org.uk) after the live examination series.

If OCR has unwittingly failed to correctly acknowledge or clear any third-party content in this assessment material, OCR will be happy to correct its mistake at the earliest possible opportunity.

For queries or further information please contact the Copyright Team, First Floor, 9 Hills Road, Cambridge CB2 1GE.

OCR is part of the Cambridge Assessment Group; Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.