

# OCR

Oxford Cambridge and RSA

## Friday 23 June 2017 – Morning

### A2 GCE MATHEMATICS (MEI)

4754/01A Applications of Advanced Mathematics (C4) Paper A

#### QUESTION PAPER

Candidates answer on the Printed Answer Book.

#### OCR supplied materials:

- Printed Answer Book 4754/01A
- MEI Examination Formulae and Tables (MF2)

#### Other materials required:

- Scientific or graphical calculator

**Duration:** 1 hour 30 minutes



#### INSTRUCTIONS TO CANDIDATES

These instructions are the same on the Printed Answer Book and the Question Paper.

- The Question Paper will be found inside the Printed Answer Book.
- Write your name, centre number and candidate number in the spaces provided on the Printed Answer Book. Please write clearly and in capital letters.
- **Write your answer to each question in the space provided in the Printed Answer Book.** If additional space is required, you should use the lined page(s) at the end of the Printed Answer Book. The question number(s) must be clearly shown.
- Use black ink. HB pencil may be used for graphs and diagrams only.
- Read each question carefully. Make sure you know what you have to do before starting your answer.
- Answer **all** the questions.
- Do **not** write in the barcodes.
- You are permitted to use a scientific or graphical calculator in this paper.
- Final answers should be given to a degree of accuracy appropriate to the context.

#### INFORMATION FOR CANDIDATES

This information is the same on the Printed Answer Book and the Question Paper.

- The number of marks is given in brackets [ ] at the end of each question or part question on the Question Paper.
- You are advised that an answer may receive **no marks** unless you show sufficient detail of the working to indicate that a correct method is being used.
- The total number of marks for this paper is **72**.
- The Printed Answer Book consists of **16** pages. The Question Paper consists of **4** pages. Any blank pages are indicated.
- This paper will be followed by **Paper B: Comprehension**.

#### INSTRUCTION TO EXAMS OFFICER/INVIGILATOR

- Do not send this Question Paper for marking; it should be retained in the centre or recycled. Please contact OCR Copyright should you wish to re-use this document.

## Section A (36 marks)

- 1 (i) Express  $\frac{5-x}{(2-x)(1+x)}$  in partial fractions. [3]
- (ii) Hence or otherwise find the first 3 terms of the binomial expansion of  $\frac{5-x}{(2-x)(1+x)}$  in ascending powers of  $x$ . [5]

- 2 The equation of a line is  $\mathbf{r} = \begin{pmatrix} -1 \\ 0 \\ 1 \end{pmatrix} + \lambda \begin{pmatrix} 2 \\ -3 \\ 1 \end{pmatrix}$  and the equation of a plane is  $3x + 4y - z = 17$ .

(i) Find the coordinates of the point of intersection of the line and the plane. [4]

(ii) Find the acute angle between the line and the normal to the plane. [4]

- 3 Fig. 3 shows the curve  $y = \sqrt{1 + e^{2x}}$ .




Fig. 3

The value of  $\int_{-1}^1 \sqrt{1 + e^{2x}} dx$  is to be estimated using the trapezium rule.  $T_2$  and  $T_4$  are the estimates obtained from the trapezium rule using 2 strips and 4 strips respectively.

(i) Explain whether  $T_4$  is greater or less than  $T_2$ . [2]

(ii) Evaluate  $T_4$ , giving your answer to 3 significant figures. [4]

- 4 Vectors  $\mathbf{u}$  and  $\mathbf{v}$  are given by  $\mathbf{u} = \mathbf{i} - 7\mathbf{j} - 2\mathbf{k}$  and  $\mathbf{v} = a\mathbf{i} + b\mathbf{j} + 5\mathbf{k}$ , where  $a$  and  $b$  are constants.

Find  $a$  and  $b$  given that the magnitude of  $\mathbf{v}$  is  $\sqrt{27}$  and that  $\mathbf{u}$  and  $\mathbf{v}$  are perpendicular. [6]

- 5 Solve the equation  $4 \tan \theta \tan 2\theta = 1$ , for  $0^\circ < \theta < 180^\circ$ . [4]

- 6 The number of bacteria in a population at time  $t$  is denoted by  $P$ . The rate of increase of  $P$  is proportional to the square root of  $P$ .

(i) Write down a differential equation relating  $P$ , the time  $t$ , and a constant of proportionality  $k$ . [1]

(ii) Verify that  $P = (A + Bt)^2$ , where  $A$  and  $B$  are constants, satisfies the differential equation, and find  $k$  in terms of  $B$ . [3]

## Section B (36 marks)

- 7 The curve shown in Fig. 7 passes through the origin and satisfies the differential equation

$$\frac{dy}{dx} = \frac{9x}{4(y+3)}.$$




Fig. 7

- (i) Show by integration that the equation of the curve is  $9x^2 - 4y^2 - 24y = 0$ . [5]

The finite region bounded by the curve and the line  $y = 2$  is rotated through  $180^\circ$  about the  $y$ -axis.

- (ii) Find the volume of the solid of revolution generated, giving your answer as an exact multiple of  $\pi$ . [4]

- (iii) Use the substitutions  $x = 2 \tan \theta$  and  $y = 3(\sec \theta - 1)$

(A) to verify that  $9x^2 - 4y^2 - 24y = 0$ ,

(B) to show that  $\frac{9x}{4(y+3)}$  can be expressed as  $k \sin \theta$ , where  $k$  is a constant to be found.

Hence find the exact gradient of the curve at the point with  $x$ -coordinate 2. [9]

8 Fig. 8 shows the curve with parametric equations

$$x = \cos 2\theta, \quad y = \cos \theta + 2 \sin \theta, \quad \text{for } -\pi < \theta \leq \pi.$$

The curve intersects the  $x$ -axis at A, and the points B and C have maximum  $x$ - and  $y$ -coordinates respectively.




Fig. 8

- (i) Find the value of  $\theta$  corresponding to the point B. Hence find the coordinates of the point B. [3]
- (ii) Express  $\cos \theta + 2 \sin \theta$  in the form  $R \cos(\theta - \alpha)$ , where  $R > 0$  and  $0 < \alpha < \frac{1}{2}\pi$ . [5]
- (iii) Hence find the coordinates of the points A and C. [5]

The angle  $\beta$  is the angle between the tangents to the curve at A.

- (iv) Find  $\frac{dy}{dx}$  in terms of  $\theta$ . Hence, assuming the scales of the  $x$ - and  $y$ -axes are equal, find  $\beta$ , giving the answer in radians correct to 2 decimal places. [You may assume the curve is symmetrical about the  $x$ -axis.] [5]

END OF QUESTION PAPER

**OCR**  
Oxford Cambridge and RSA

**Copyright Information**

OCR is committed to seeking permission to reproduce all third-party content that it uses in its assessment materials. OCR has attempted to identify and contact all copyright holders whose work is used in this paper. To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced in the OCR Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download from our public website ([www.ocr.org.uk](http://www.ocr.org.uk)) after the live examination series.

If OCR has unwittingly failed to correctly acknowledge or clear any third-party content in this assessment material, OCR will be happy to correct its mistake at the earliest possible opportunity.

For queries or further information please contact the Copyright Team, First Floor, 9 Hills Road, Cambridge CB2 1GE.

OCR is part of the Cambridge Assessment Group; Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.

**Friday 23 June 2017 – Morning**

**A2 GCE MATHEMATICS (MEI)**

**4754/01B Applications of Advanced Mathematics (C4) Paper B: Comprehension**

**QUESTION PAPER**

Candidates answer on the Question Paper.


**OCR supplied materials:**

- Insert (inserted)
- MEI Examination Formulae and Tables (MF2)

**Other materials required:**

- Scientific or graphical calculator

**Duration: Up to 1 hour**



Candidate forename		Candidate surname	
-----------------------	--	----------------------	--

Centre number						Candidate number				
---------------	--	--	--	--	--	------------------	--	--	--	--

**INSTRUCTIONS TO CANDIDATES**

- The Insert will be found inside this document.
- Write your name, centre number and candidate number in the boxes above. Please write clearly and in capital letters.
- Use black ink. HB pencil may be used for graphs and diagrams only.
- Answer **all** the questions.
- Read each question carefully. Make sure you know what you have to do before starting your answer.
- Write your answer to each question in the space provided. Additional paper may be used if necessary but you must clearly show your candidate number, centre number and question number(s).
- Do **not** write in the bar codes.
- The Insert contains the text for use with the questions.
- You are permitted to use a scientific or graphical calculator in this paper.
- Final answers should be given to a degree of accuracy appropriate to the context.

**INFORMATION FOR CANDIDATES**

- The number of marks is given in brackets [ ] at the end of each question or part question.
- You may find it helpful to make notes and do some calculations as you read the passage.
- You are **not** required to hand in these notes with your question paper.
- You are advised that an answer may receive **no marks** unless you show sufficient detail of the working to indicate that a correct method is being used.
- The total number of marks for this paper is **18**.
- This document consists of **8** pages. Any blank pages are indicated.

1 State the set of values of  $x_0$  for which the iteration

$$x_{n+1} = 2.5x_n(1-x_n)$$

(i) converges to a single non-zero number, [1]

(ii) has all terms from  $x_1$  onwards equal to zero. [1]

<b>1(i)</b>	
<b>1(ii)</b>	

- 2 (i) Use the algebraic method indicated in lines 68 to 70 to find the equilibrium point of the iteration

$$x_{n+1} = 1.6x_n(1-x_n). \quad [2]$$

- (ii) Show that the iteration

$$x_{n+1} = x_n^2 + 2$$

does not have any points of equilibrium. [2]

<b>2(i)</b>	
<b>2(ii)</b>	

- 3 One of the assumptions for the model used for the population of squirrels in the text was that there are no predators.

An alternative model is proposed in which predators kill a fixed number of squirrels each year.

An iterative equation for this model is given by

$$x_{n+1} = kx_n(1-x_n) - 0.25.$$

In the table below  $x_0$  is taken to be 0.55 and four different values are considered for  $k$ .

- (i) Complete as many of the empty cells as you need to in order to establish the outcomes for these values of  $k$ .

- (ii) Comment on what the table tells you for each of the four values of  $k$ .

[6]

<b>3(i)</b>	$x_{n+1} = kx_n(1-x_n) - 0.25$				
		$k = 2$	$k = 3$	$k = 4$	$k = 5$
	$x_0$	0.55	0.55	0.55	0.55
	$x_1$	0.245	0.4925	0.74	0.9875
	$x_2$				
	$x_3$				
	$x_4$				
	$x_5$				
	$x_6$				
	$x_7$				
	$x_8$				
	$x_9$				
$x_{10}$					
...	...	...	...	...	



- 4 (i) Table 3 gives the first four points of bifurcation of the iteration

$$x_{n+1} = kx_n(1 - x_n).$$

Feigenbaum's Constant is 4.6692 correct to 5 significant figures. Using this value for the ratio of the interval lengths, estimate the values of  $k$  for the next two points of bifurcation. [3]

- (ii) (A) Find,  $S$ , the sum to infinity of the geometric series

$$1 + \frac{1}{4.6692} + \left(\frac{1}{4.6692}\right)^2 + \left(\frac{1}{4.6692}\right)^3 + \dots$$

[2]

- (B) Using certain figures from Table 3, a value of  $k$  is estimated to be

$$k = 3.5644 + 0.0203 \times S.$$

State what happens at this value of  $k$ .

[1]

<b>4(i)</b>	

<b>4(ii)(A)</b>	
<b>4(ii)(B)</b>	

**END OF QUESTION PAPER**

**PLEASE DO NOT WRITE ON THIS PAGE**

---

**OCR**  
Oxford Cambridge and RSA

**Copyright Information**

OCR is committed to seeking permission to reproduce all third-party content that it uses in its assessment materials. OCR has attempted to identify and contact all copyright holders whose work is used in this paper. To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced in the OCR Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download from our public website ([www.ocr.org.uk](http://www.ocr.org.uk)) after the live examination series.

If OCR has unwittingly failed to correctly acknowledge or clear any third-party content in this assessment material, OCR will be happy to correct its mistake at the earliest possible opportunity.

For queries or further information please contact the Copyright Team, First Floor, 9 Hills Road, Cambridge CB2 1GE.

OCR is part of the Cambridge Assessment Group; Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.

# OCR

Oxford Cambridge and RSA

**Friday 23 June 2017 – Morning**

**A2 GCE MATHEMATICS (MEI)**

**4754/01B** Applications of Advanced Mathematics (C4) Paper B: Comprehension

**INSERT**

**Duration:** Up to 1 hour



## **INFORMATION FOR CANDIDATES**

- This Insert contains the text for use with the questions.
- This document consists of **8** pages. Any blank pages are indicated.

## **INSTRUCTION TO EXAMS OFFICER/INVIGILATOR**

- Do not send this Insert for marking; it should be retained in the centre or recycled. Please contact OCR Copyright should you wish to re-use this document.

## Feigenbaum's Constant

### A population model

A remote island has a population of squirrels. There are no predators so that the population is limited only by the supply of food and death from natural causes.

How would the number of squirrels be expected to change from year to year?

5

A simple model involves the following variables.

- Time is measured in years and the year number is denoted by  $n$ . In this article, year  $n$  is also described as 'this year' and year  $n+1$  as 'next year'.
- The size of the squirrel population is given as the proportion of the maximum possible population; the population is given at the start of each year. The population in year  $n$  is denoted by  $x_n$  where  $0 \leq x_n \leq 1$ . Similarly, in year  $n+1$  the population is  $x_{n+1}$ .
- A parameter  $k$  is a measure of the reproductivity of the squirrels, and so determines the rate of growth of their population.

10

The model involves the following assumptions.

- The number of squirrels next year,  $x_{n+1}$ , is jointly proportional to the number this year,  $x_n$ , and to the quantity  $(1-x_n)$  which represents the food available.
- The parameter  $k$  is the constant of proportionality.

15

The model can be expressed by the iterative equation

$$x_{n+1} = kx_n(1-x_n).$$

This is a general model for a population in a restricted environment without predators. The animals do not have to be squirrels.

20

### Different values of $k$

To investigate this model, it is first necessary to choose values for  $k$  and for the initial population,  $x_0$ . Table 1 gives the first ten values of  $x_n$  for four particular values of  $k$  and with  $x_0 = 0.5$ . The numbers given in the table have been truncated; many more figures were used in calculating them.

25

	$x_{n+1} = kx_n(1-x_n)$			
	$k = 0.6$	$k = 1.5$	$k = 3.3$	$k = 4.5$
$x_0$	0.5	0.5	0.5	0.5
$x_1$	0.15	0.375	0.825	1.125
$x_2$	0.0765	0.3515...	0.4764...	-0.6328...
$x_3$	0.0423...	0.3419...	0.8231...	-4.649...
$x_4$	0.0243...	0.3375...	0.4803...	-118.2...
$x_5$	0.0142...	0.3354...	0.8237...	-
$x_6$	0.0084...	0.3343...	0.4791...	-
$x_7$	0.0050...	0.3338...	0.8235...	-
$x_8$	0.0029...	0.3335...	0.4795...	-
$x_9$	0.0017...	0.3334...	0.8236...	-
$x_{10}$	0.0010...	0.3333...	0.4794...	-
...	...	...	...	...

30

35

**Table 1**

40

The different values of  $k$  used in Table 1 lead to four different outcomes for the population in this model.

- When  $k = 0.6$ ,  $x_n \rightarrow 0$ . Eventually the population dies out.
- When  $k = 1.5$ ,  $x_n \rightarrow 0.3333... = \frac{1}{3}$ . The population attains a stable equilibrium level.
- When  $k = 3.3$ , the population alternates between a high value of 0.823... and a low value of 0.479... in successive years.
- When  $k = 4.5$ , the population appears to become negative and so to have died out. In the first year the population exceeds the limit of 1 and so the model has broken down.

45

In each case in Table 1 the value of  $x_0$  was taken to be 0.5, the middle of the possible values of  $x_0$ . If you try other starting values you will find that the final outcomes are the same for any values of  $x_0$  between, but not including, 0 and 1.

50

Overall the model suggests that having too few or too many young can both be fatal for the population.

## Possible outcomes

This iteration can be used as a population model, but it can also be thought of as a mathematical iteration in its own right with an interesting variety of possible outcomes.

At this stage it is helpful to extend the notation used to include the letter  $x$ . This denotes the value, or values, of  $x_n$  as  $n$  tends to infinity. 55

In Fig. 2, these limiting values,  $x$ , are plotted for values of  $k$  between 0 and 3.6. For larger values of  $k$  there are no limiting values. It is assumed that  $0 < x_0 < 1$ .




Fig. 2

Fig. 2 shows that, for this iteration, five different types of outcome are possible according to the value of  $k$ . These are described below. 60

### *Tending to zero*

For  $0 \leq k \leq 1$ , the iteration converges to zero.

### *Converging to a single non-zero value*

For  $1 < k < 3$ , the iteration converges to a non-zero value between 0 and 1. 65

An example of this is given in Table 1 for  $k = 1.5$ . The iteration is found to converge to  $\frac{1}{3}$ . This value may be described as an *equilibrium point*.

It can be found algebraically. Denoting it by  $x$  gives the equation

$$x = 1.5x(1-x)$$

which has a solution of  $x = 0$  or  $\frac{1}{3}$ .

70

### ***Oscillating***

Table 1 shows that for  $k = 3.3$ ,  $x_n$  oscillates between two values. There is a range of values of  $k$  for which this occurs.

- For  $k = 2.99$ ,  $x_n$  converges slowly to 0.6655... . At  $k = 3$ , it starts to oscillate. After 5000 iterations the low value is 0.6633... and the high value is 0.6699... . This is a cycle of length 2.

75

Thus it is found, for example by experiment using a spreadsheet, that the smallest value of  $k$  for which the iteration oscillates is 3. Such a value of  $k$  where the iteration splits is called a *point of bifurcation*.

However,  $k = 3$  is not the only point of bifurcation. At  $k = 3.449...$  there is a further point of bifurcation at which the cycle of length 2 becomes a cycle of length 4.

- For  $k = 3.5$ , the four values of  $x$  are 0.3828..., 0.8269..., 0.5008... and 0.8749... .

80

Another point of bifurcation occurs at  $k = 3.544...$  . At this point the length of the cycle goes up to 8. Further points of bifurcation give cycles of length 16, then 32, then 64 and so on.

### ***Chaos***

The pattern of cycles of increasing length does not continue indefinitely as  $k$  increases. For larger values of  $k$ , for example  $k = 3.8$ , the outcomes have no pattern, and so the situation is described as *chaos*. It is very difficult to distinguish between chaos and a long cycle; a sophisticated computer program is required.

85

A feature of chaos is that the iteration remains bounded. The values of  $x_n$  are always between 0 and 1.

### ***Unbounded outcomes***

For values of  $k \geq 4$ , the outcomes cease to be bounded. An example of this occurs in the final column of Table 1, where the value of  $k$  is 4.5.

90

### Feigenbaum's Constant

To summarise, from  $k = 1$  to 3 the iteration  $x_{n+1} = kx_n(1-x_n)$  converges to a single non-zero value. There is then a point of bifurcation and this is followed by further points of bifurcation. It is evident from Fig. 2 that the intervals between successive points of bifurcation become progressively shorter.

95

Information about these intervals is given in Table 3. The numbers in this table have been rounded to 4 decimal places.

Cycle length	Boundary values of $k$		Interval	$\frac{\text{Previous interval}}{\text{This interval}}$
1	1	3	2	–
2	3	3.4495	0.4495	4.4494
4	3.4495	3.5441	0.0946	4.7516
8	3.5441	3.5644	0.0203	4.6601
16	3.5644	...	...	...

100

**Table 3**

The right hand column gives the ratio by which the length of this interval decreases with successive cycles. The three values of this ratio in Table 3 are close together.

105

In the 1970s, Mitchell Feigenbaum started to investigate this ratio. He discovered that similar patterns of bifurcation are found with many other iterations; examples include  $x_{n+1} = x_n^2 + k$  and  $x_{n+1} = k \sin(\pi x_n)$ .

He also discovered that the ratio tends to a definite limit and that this has the same value for all iterations that show this pattern of bifurcation.

110

His work was conducted to a very high level of accuracy and covered many more cycles than the small number considered here. The limited power of computers in those days meant that it was an enormous undertaking.

Feigenbaum was immediately convinced of the importance of his discovery.

*'I called my parents that evening and told them I had discovered something truly remarkable, that, when I had understood it, would make me a famous man.'*

115

He is indeed now a famous man and the ratio he discovered is called Feigenbaum's Constant. It has now been found to over 1000 figures; the first ten of these are 4.669 201 609.

**BLANK PAGE**

**Copyright Information**

OCR is committed to seeking permission to reproduce all third-party content that it uses in its assessment materials. OCR has attempted to identify and contact all copyright holders whose work is used in this paper. To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced in the OCR Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download from our public website ([www.ocr.org.uk](http://www.ocr.org.uk)) after the live examination series.

If OCR has unwittingly failed to correctly acknowledge or clear any third-party content in this assessment material, OCR will be happy to correct its mistake at the earliest possible opportunity.

For queries or further information please contact the Copyright Team, First Floor, 9 Hills Road, Cambridge CB2 1GE.

OCR is part of the Cambridge Assessment Group; Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.