

# Mark Scheme (Results)

January 2016

International GCSE Mathematics A  
4MA0/4HR

## **Edexcel and BTEC Qualifications**

Edexcel and BTEC qualifications come from Pearson, the world's leading learning company. We provide a wide range of qualifications including academic, vocational, occupational and specific programmes for employers. For further information, please visit our website at [www.edexcel.com](http://www.edexcel.com).

Our website subject pages hold useful resources, support material and live feeds from our subject advisors giving you access to a portal of information. If you have any subject specific questions about this specification that require the help of a subject specialist, you may find our Ask The Expert email service helpful.

[www.edexcel.com/contactus](http://www.edexcel.com/contactus)

## **Pearson: helping people progress, everywhere**

Our aim is to help everyone progress in their lives through education. We believe in every kind of learning, for all kinds of people, wherever they are in the world. We've been involved in education for over 150 years, and by working across 70 countries, in 100 languages, we have built an international reputation for our commitment to high standards and raising achievement through innovation in education. Find out more about how we can help you and your students at: [www.pearson.com/uk](http://www.pearson.com/uk)

January 2016

Publications Code UG043268

All the material in this publication is copyright

© Pearson Education Ltd 2016

## General Marking Guidance

- All candidates must receive the same treatment. Examiners must mark the first candidate in exactly the same way as they mark the last.
- Mark schemes should be applied positively. Candidates must be rewarded for what they have shown they can do rather than penalised for omissions.
- Examiners should mark according to the mark scheme not according to their perception of where the grade boundaries may lie.
- There is no ceiling on achievement. All marks on the mark scheme should be used appropriately.
- All the marks on the mark scheme are designed to be awarded. Examiners should always award full marks if deserved, i.e. if the answer matches the mark scheme. Examiners should also be prepared to award zero marks if the candidate's response is not worthy of credit according to the mark scheme.
- Where some judgement is required, mark schemes will provide the principles by which marks will be awarded and exemplification may be limited.
- When examiners are in doubt regarding the application of the mark scheme to a candidate's response, the team leader must be consulted.
- Crossed out work should be marked UNLESS the candidate has replaced it with an alternative response.
- **Types of mark**
  - M marks: method marks
  - A marks: accuracy marks
  - B marks: unconditional accuracy marks (independent of M marks)
- **Abbreviations**
  - cao – correct answer only
  - ft – follow through
  - isw – ignore subsequent working
  - SC - special case
  - oe – or equivalent (and appropriate)
  - dep – dependent
  - indep – independent
  - eeoo – each error or omission

- **No working**  
 If no working is shown then correct answers normally score full marks  
 If no working is shown then incorrect (even though nearly correct) answers score no marks.
- **With working**  
 If there is a wrong answer indicated on the answer line always check the working in the body of the script (and on any diagrams), and award any marks appropriate from the mark scheme.  
 If it is clear from the working that the “correct” answer has been obtained from incorrect working, award 0 marks.  
 Any case of suspected misread loses A (and B) marks on that part, but can gain the M marks.  
 If working is crossed out and still legible, then it should be given any appropriate marks, as long as it has not been replaced by alternative work.  
 If there is a choice of methods shown, then no marks should be awarded, unless the answer on the answer line makes clear the method that has been used.  
 If there is no answer on the answer line then check the working for an obvious answer.
- **Ignoring subsequent work**  
 It is appropriate to ignore subsequent work when the additional work does not change the answer in a way that is inappropriate for the question: eg. Incorrect cancelling of a fraction that would otherwise be correct.  
 It is not appropriate to ignore subsequent work when the additional work essentially makes the answer incorrect eg algebra.  
 Transcription errors occur when candidates present a correct answer in working, and write it incorrectly on the answer line; mark the correct answer.
- **Parts of questions**  
 Unless allowed by the mark scheme, the marks allocated to one part of the question CANNOT be awarded in another.

<b>International GCSE Maths</b>				
<b>Apart from Questions 6e, 14d and 24 (where the mark scheme states otherwise), the correct answer, unless clearly obtained by an incorrect method, should be taken to imply a correct method.</b>				
<b>Q</b>	<b>Working</b>	<b>Answer</b>	<b>Mark</b>	<b>Notes</b>
<b>1 (a)</b>	$(40 \div 16) \times 240$ oe			M1 for a fully correct method
		600	2	A1
<b>(b)</b>	$(600 \div 120) \times 16$ oe			M1 for a fully correct method
		80	2	A1
<b>(c)</b>	$240 \div 150$ <b>or</b> $150 : 240$ oe			M1
		1.6 oe	2	A1
				<b>Total 6 marks</b>

<b>2 (a) (i)</b>		57	1	B1
<b>(ii)</b>		<u>Corresponding angles</u>	1	B1 For correct reason
<b>(b)</b>	$(5 - 2) \times 180$ <b>or</b> $3 \times 180$ <b>or</b> $(2 \times 5 - 4) \times 90$ <b>or</b> $6 \times 90$ <b>or</b> $360 + 180$ <b>or</b> 540			M1 for correct method to find total of angles in a pentagon <b>or</b>
	'540' - (86+142+72+115) oe			M1 (dep) fully correct method to find y
		125	3	A1 cao
	<b>Alternative method (exterior angles)</b> $360 - ("94" + "38" + "108" + "65") (=55)$ <b>or</b> $360 - 305 (=55)$			M1 if just values seen then condone one error in exterior angles
	$180 - "55"$			M1 (dep) fully correct method to find y
		125	3	A1 cao
				<b>Total 5 marks</b>

<b>3</b>	$\frac{3 \times 5}{20} + \frac{4 \times 4}{20}$ or $\frac{15}{20} + \frac{16}{20}$			M1 for any pair of correct fractions with denominator a multiple of 20
		$\frac{31}{20}$	2	A1 dependent on M1
	<b>Alternative</b> $0.75 + 0.8 = 1.55$			M1
		$1\frac{55}{100}$		A1 dependent on M1
				<b>Total 2 marks</b>

<b>4</b>		5, 10, 20, 25, 50, 100	2	B2 If not B2 then  B1 for at least 3 correct values and no incorrect values <b>or</b> all correct values with only 1 incorrect value
				<b>Total 2 marks</b>

<b>5</b>	$48 \div 8 (=6)$			M1 width of rectangle
	$(8 + "6") \times 2 (=28)$			M1 perimeter
	$"28" \div 4 (=7)$			M1 length of side
		49	4	A1
				<b>Total 4 marks</b>

<b>6</b>	<b>(a)</b>		$x^9$	1	
	<b>(b)</b>		$y^6$	1	
	<b>(c)</b>	$8d+12-6d+10$ oe	$2d+22$	2	M1 for 3 terms with correct signs or 4 terms without signs A1 for $2d+22$ <b>or</b> $2(d+11)$
	<b>(d)</b>	eg. $9y-5y=2+3$ <b>or</b> $4y=5$			M1 for a correct equation with terms in $y$ on one side and numbers on the other.
			1.25	2	A1 for 1.25 or $\frac{5}{4}$ or $1\frac{1}{4}$
	<b>(e)</b>	$7x-1=5x$			M1 multiplying $x$ by 5 (seen as part of an equation) or showing $\frac{7}{5}x - \frac{1}{5} = x$
		eg. $7x-5x=1$ <b>or</b> $2x=1$ <b>or</b> $\frac{7}{5}x - x = \frac{1}{5}$			M1 for isolating terms in $x$
			$\frac{1}{2}$ oe	3	A1 for $\frac{1}{2}$ or 0.5 dep on M1 scored
					<b>Total 9 marks</b>

<b>7</b>	<b>(a)</b>		15 - 19	1	B1
	<b>(b)</b>	$2 \times 1 + 7 \times 5 + 12 \times 6 + 17 \times 10 + 22 \times 8$ or $2 + 35 + 72 + 170 + 176$ or 455			M2 Freq $\times$ midpoint values stated or evaluated with intention to add (condone any two errors in midpoints or frequencies).  If not M2 then award M1 for all products $t \times f$ (and $t$ is consistently within the interval, including end values) and intention to add (condone two errors).
		$\frac{2 \times 1 + 7 \times 5 + 12 \times 6 + 17 \times 10 + 22 \times 8}{30}$ or "455" $\div$ 30			M1 (dep on at least M1) for division by 30
			15.2	4	A1 accept 15.166... rounded or truncated to 4 or more sig figs Accept 15 with working (15 without working gains M0A0) NB: accept 2.25 as mid-point for mid-interval value of 1 <sup>st</sup> class (gives mean 15.175)
					<b>Total 5 marks</b>

<b>8</b>	$0.16x = 192$ or $16\% = 192$ oe or $\frac{192}{16} (=12)$			3	M1
	$\frac{192}{0.16}$ or $\frac{192}{16} \times 100$ oe				M1
		1200			A1 cao
					<b>Total 3 marks</b>



<b>9</b>	$1\frac{24}{60}$ oe <b>or</b> 1.4 <b>or</b> 84			B1 for changing time to a decimal or to minutes
	$\frac{725}{1.4}$ oe <b>or</b> $\frac{725}{84} \times 60$			M1 allow $725 \div 1.24$
		518	3	A1 for 518 or 517.857...
				<b>Total 3 marks</b>

<b>10</b>	(a)		1 590 000	1	B1 cao
	(b)	$2.97 \times 10^6 - 4.22 \times 10^5$ oe <b>or</b> $4.22 \times 10^5 - 2.97 \times 10^6$ oe <b>or</b> $\pm 2548000$			M1
			$\pm 2.548 \times 10^6$	2	A1 allow $\pm 2.5 \times 10^6$ or $\pm 2.55 \times 10^6$ if working shown.
	(c)	$1.25 \times 10^6 \div 4.22 \times 10^5$ oe <b>or</b> 2.96... <b>or</b> $\frac{625}{211}$			M1
			3	2	A1 cao
					<b>Total 5 marks</b>

11	$\pi \times (20 - 2 \times 4)$ oe <b>or</b> $\pi \times 12$ oe <b>or</b> $2 \times \pi \times 6$ <b>or</b> 37.6... <b>or</b> 37.7 <b>or</b>			3	M1 for a correct method to find the circumference or half of the circumference
	$\frac{1}{2} \times \pi \times (20 - 2 \times 4)$ oe <b>or</b> $\frac{1}{2} \times \pi \times 12$ oe <b>or</b> $\pi \times 6$ <b>or</b> 18.8...				
	$4 + 10 + 20 + 10 + 4 + \frac{1}{2} \times "37.6\dots"$ <b>or</b> $4 + 10 + 20 + 10 + 4 + "18.8\dots"$				M1 (dep on previous M1) for complete method
			66.8		A1 66.8 – 66.9
					<b>Total 3 marks</b>

12	$(4x^2 + 6x + 6x + 9) - (4x^2 - 6x - 6x + 9)$			M1 any one correct expansion	M1 for $((2x + 3) + (2x - 3))((2x + 3) - (2x - 3))$
	$4x^2 + 6x + 6x + 9 - 4x^2 + 6x + 6x - 9$			M1 for dealing correctly with the negative sign	M1 for $(2x + 3 + 2x - 3)(2x + 3 - 2x + 3)$
		24x	3	A1	
					<b>Total 3 marks</b>

<b>13</b>	(a)	eg. $22 \times \frac{24}{20}$ <b>or</b> $22 \times 1.2$			M1 for complete method <b>or</b> correct scale factor (may be seen within an equation) eg. 20/24 or 24/20 or 1.2 or 0.83...oe
			26.4	2	A1
	(b)	eg. $28.2 - 28.2 \div "1.2"$ <b>or</b> $28.2 \div 6$ oe			M1ft for a complete method ft from "1.2" used in (a) which must come from a correct method
			4.7	2	A1
					<b>Total 4 marks</b>

<b>14</b>	(a)		-10, (4), 6, 2, (-2), 0, (14)	2	B2 Award B1 for any 2 correct.
	(b)	(-3, -10), (-2, 4), (-1, 6), (0, 2), (1, -2), (2, 0), ((3, 14),	Correct curve	2	B2 For the correct smooth curve.  B1 for at least 5 points plotted correctly; ft from table for plotting only provided at least B1 scored in (a).
	(c)	$1^3 - 6 \times 1 + m = 0$ <b>or</b> $1 - 6 + m = 0$	5	1	B1
	(d)				M1 $y = x - 3$ drawn
			-2.7 to -2.9	2	A1 ft from graph (dep on M1)
					<b>Total 7 marks</b>

<b>15</b>	eg. $x = 0.2666\dots$ and $10x = 2.666\dots$ <b>or</b> $10x = 2.666\dots$ and $100x = 26.66\dots$			M1	Two correct decimals that, when subtracted, will leave a non-recurring value with intention to subtract. Must see a recurring symbol or 3+ sixes.
		show	2	A1	Must see a fraction prior to seeing $\frac{4}{15}$
	<b>Alternative method</b>				
	eg. $x = 0.0666\dots$ and $100x = 6.666$ with subtraction			M1	for conversion of 0.0666... to a fraction
			2	A1	for conclusion with all steps seen
					<b>Total 2 marks</b>

<b>16</b>	$Q = \frac{"k"}{t^2}$			3	M1 $k$ must be a letter not a number
	$320 = \frac{"k"}{0.5^2}$ or " $k$ " = 80				M1 for substitution (implies first M1)
		$Q = \frac{80}{t^2}$			A1 Award 3 marks for $Q = \frac{"k"}{t^2}$ and " $k$ " = 80 stated anywhere
					<b>Total 3 marks</b>

<b>17</b>	<b>(a)</b>		0.4, 0.9, 0.3, 0.7 in appropriate places	2	B2 All correct (B1 for any one correct)
	<b>(b)</b>	0.6 × 0.1			M1
			0.06	2	A1 or 6% or $\frac{3}{50}$ oe
	<b>(c)</b>	(0.6 × '0.9') + ('0.4' × '0.7')			M2ft Award M1 for one correct product
			0.82	3	A1 or 82% or $\frac{41}{50}$ oe
					<b>Total 7 marks</b>

<b>18</b>	<b>(a)</b>	1cm × 1cm square = 2.5 people or 1 large square = 10 people or 4.8 on axis corresponding to top of 20-25 bar or Correct scale marked on vertical axis or 10 small squares = 1 person oe		2	M1
			31		A1
	<b>(b)</b>	Freq densities: 38/10 (=3.8), 63/15 (=4.2), 24/30 (=0.8)		2	M1 2 correct frequency densities <b>or</b> 1 of the 3 bars correctly drawn
					A1 All 3 bars correctly drawn
					<b>Total 4 marks</b>

<b>19</b>	$4g - 9eg = 7 - 3e$ or $3e - 7 = 9eg - 4g$			M1	Correctly collecting terms in $g$ on one side and everything else on the other.
	$g(4 - 9e) = 7 - 3e$ or $3e - 7 = g(9e - 4)$			M1	Factorising $g(4-9e)$ or $g(9e-4)$
		$g = \frac{7-3e}{4-9e}$ or $g = \frac{3e-7}{9e-4}$	3	A1	
				<b>Total 3 marks</b>	

<b>20</b>	$\frac{3(2x + 5) - 6(x + 2)}{(x + 2)(2x + 5)}$			M1	For expressing both fractions correctly with a common denominator. Allow as two separate fractions. Condone one error in numerator expansion.
	$\frac{6x + 15 - 6x - 12}{(x + 2)(2x + 5)}$			M1	For removing brackets correctly in a correct single fraction. Allow denominator to be expanded or 2 brackets.
		$\frac{3}{(x + 2)(2x + 5)}$	3	A1	$\frac{3}{2x^2 + 9x + 10}$ or
				<b>Total 3 marks</b>	

21	(BD <sup>2</sup> or AC <sup>2</sup> =) 15 <sup>2</sup> + 15 <sup>2</sup> or $\sqrt{450}$ or $15\sqrt{2}$ or 21.2 ...			M1 A correct statement to find diagonal of base
	eg. " $\sqrt{450}$ " <sup>2</sup> = 12 <sup>2</sup> + 12 <sup>2</sup> - 2×12×12×cosBED or $\sin BEX = \frac{0.5\sqrt{450}}{12}$ or $\cos EBD = \frac{0.5\sqrt{450}}{12}$			M1 dep for use of cosine rule – correct statement in any form or correct trig statement to find angle BED or angle BEX or angle EBD or angle EDB  <b>NB:</b> Any multi-step method must be fully correct
	eg $DEB = \cos^{-1}\left(\frac{12^2 + 12^2 - 450}{2 \times 12 \times 12}\right)$  or $DEB = 2 \times \sin^{-1}\left(\frac{0.5\sqrt{450}}{12}\right)$  or $DEB = 180 - 2 \times \cos^{-1}\left(\frac{0.5\sqrt{450}}{12}\right)$  or $DEB = 55.7\dots$ or $56^\circ$ (from ambiguous case of Sine Rule)			M1 Complete correct method to find angle DEB
		124	4	A1 answer in range 124 – 124.3
				<b>Total 4 marks</b>

22	$a^2 + a\sqrt{b} + a\sqrt{b} + b$ <b>or</b> $a^2 + a\sqrt{b} + a\sqrt{b} + (\sqrt{b})^2$			M1 Correct expansion
		6		A1 For $a$
		13	3	A1 For $b$
				<b>Total 3 marks</b>

23	$0.5 \times 12 \times 14 \times \sin x = 72$			M1 For a correct equation for the area including 12, 14, $\sin x$ and 72
	$\sin x = \frac{72}{84}$			M1 For $\sin x = \frac{72}{84}$ oe
		$59^\circ$		A1
		$121^\circ$	4	A1
				<b>Total 4 marks</b>



24	$x^2 + (3x + 2)^2 = 20$		6	M1 for elimination of one variable $\left(\frac{y-2}{3}\right)^2 + y^2 = 20$
	$x^2 + 9x^2 + 6x + 6x + 4 = 20$			M1 indep for correct expansion ie. $9x^2 + 6x + 6x + 4$ <b>or</b> $\frac{y^2 - 2y - 2y + 4}{9}$
	$10x^2 + 12x - 16 = 0$ $(5x^2 + 6x - 8 = 0)$			A1 Correct simplified 3 part quadratic equation (may not be equated to zero) $10y^2 - 4y - 176 = 0$ <b>or</b> $5y^2 - 2y - 88 = 0$
	eg. $(10x - 8)(x + 2)$ <b>or</b> $(5x - 4)(2x + 4)$ <b>or</b> $(5x - 4)(x + 2)$ <b>or</b> $\frac{-12 \pm \sqrt{12^2 - 4 \times 10 \times -16}}{2 \times 10}$ <b>oe</b>			M1 for correct factorisation or substitution into quadratic formula ft (dep on at least 1 previous M1 scored) a 3 term quadratic for this mark only eg. $(5y - 22)(y + 4) = 0$
				A1 for <b>both</b> x values (or <b>both</b> y values)
		$x = \frac{4}{5}, y = \frac{22}{5}$ <b>or</b> $x = -2, y = -4$		A1 for both solutions with x and y values correctly paired
				<b>Total 6 marks</b>

Pearson Education Limited. Registered company number 872828  
with its registered office at 80 Strand, London WC2R 0RL