Vrite your name here Surname	Othe	r names
Pearson Edexcel GCE	Centre Number	Candidate Number
Chemist		
Advanced Subsid Unit 1: The Core P	iary	mistry
Advanced Subsid	iary Principles of Che	mistry Paper Reference 6CH01/01

Instructions

- Use **black** ink or ball-point pen.
- Fill in the boxes at the top of this page with your name, centre number and candidate number.
- Answer **all** questions.
- Answer the questions in the spaces provided there may be more space than you need.

Information

- The total mark for this paper is 80.
- The marks for **each** question are shown in brackets – use this as a guide as to how much time to spend on each question.
- Questions labelled with an asterisk (*) are ones where the quality of your written communication will be assessed
 you should take particular care with your spelling, punctuation and grammar, as well as the clarity of expression, on these questions.
- A Periodic Table is printed on the back cover of this paper.

Advice

- Read each question carefully before you start to answer it.
- Keep an eye on the time.
- Try to answer every question.
- Check your answers if you have time at the end.



Turn over 🕨





SECTION A

Answer ALL the questions in this section. You should aim to spend no more than 20 minutes on this section. For each question, select one answer from A to D and put a cross in the box \boxtimes . If you change your mind, put a line through the box \bigotimes and then mark your new answer with a cross \boxtimes .

1 Which molecule does **not** exhibit *E*/*Z* isomerism?





2 What is the systematic name for the compound with the following formula?



A 2-methyl-4-ethylpentane

- B 2-ethyl-4-methylpentane
- C 2,4-dimethylhexane

D 3,5-dimethylhexane

(Total for Question 2 = 1 mark)





5		ch of the following equations has the correct state symbols for the reaction of the hydrochloric acid with magnesium oxide?	
	\mathbf{X}	A MgO(s) + 2HCl(aq) \rightarrow MgCl ₂ (s) + H ₂ O(l)	
	\mathbf{X}	B MgO(s) + 2HCl(aq) \rightarrow MgCl ₂ (aq) + H ₂ O(l)	
		$C MgO(s) + 2HCI(I) \rightarrow MgCI_2(s) + H_2O(I)$	
	X	D MgO(s) + 2HCl(l) \rightarrow MgCl ₂ (aq) + H ₂ O(l)	
_		(Total for Question 5 = 1 mar	k)
6		question is about the reaction between sodium carbonate solution and dilute c acid.	
		$Na_2CO_3(aq) + 2HNO_3(aq) \rightarrow 2NaNO_3(aq) + CO_2(g) + H_2O(I)$	
	(a) \	What is the ionic equation for this reaction?	
			1)
		A Na ₂ CO ₃ (aq) + 2H ⁺ (aq) \rightarrow 2Na ⁺ (aq) + CO ₂ (g) + H ₂ O(l) B Na ⁺ (aq) = 100 ⁻ (a x) = 100 ⁻ (a x)	
		3 Na ⁺ (aq) + NO ₃ ⁻ (aq) \rightarrow NaNO ₃ (aq) 5 $O^{2^{-}}(a) \rightarrow O^{2^{-}}(a) \rightarrow O^{2^{-}}(a)$	
		$CO_3^{2-}(aq) + 2H^+(aq) \rightarrow CO_2(g) + H_2O(I)$	
	X	D $\text{CO}_3^{2-}(\text{aq}) + 2\text{HNO}_3(\text{aq}) \rightarrow 2\text{NO}_3^{-}(\text{aq}) + \text{CO}_2(\text{g}) + \text{H}_2\text{O}(\text{I})$	
		What is the volume of carbon dioxide produced from the complete reaction of 0.10 mol of nitric acid at room temperature and pressure?	
	[1 mol of any gas occupies 24 dm ³ at room temperature and pressure.]	
		A 1.2 dm ³	1)
		3 1.8 dm ³	
	_	$C = 2.4 \text{dm}^3$	
	_	D $3.6 \mathrm{dm^3}$	
	١	What volume of sodium carbonate solution of concentration 0.500 mol dm ⁻³ , would be needed to completely react with 25.0 cm ³ of nitric acid of concentration 0.250 mol dm ⁻³ ?	
	\mathbf{X}	A 6.25 cm ³	1)
	\mathbf{X}	3 12.50 cm ³	
	\mathbf{X}	C 18.75 cm ³	
	\mathbf{X}	D 25.00 cm ³	
	_	(Total for Question 6 = 3 mark	s)

7	Lithiu	m reacts with water to produce hydrogen.	
		$Li(s) + H_2O(I) \rightarrow LiOH(aq) + \frac{1}{2}H_2(g)$	
		an experiment, 0.069 g (0.01 mol) of lithium produced 90 cm ³ of hydrogen at om temperature and pressure. What is the percentage yield of hydrogen?	
	[1	mol of any gas occupies 24 dm ³ at room temperature and pressure.]	
	A	45%	(1)
	B	60%	
	🛛 C	75%	
	D 🛛	90%	
		hich of the following is not a possible reason for the yield being less than 0%?	
	🖾 A	Some oil remained on the surface of the lithium.	(1)
	B	Hydrogen gas is very soluble in water.	
	🛛 C	A layer of oxide was present on the surface of the lithium.	
	D 🛛	Some of the hydrogen gas escaped collection.	
		(Total for Question 7 = 2 ma	rks)
8		nany moles of atoms are present in 240 cm ³ of carbon dioxide at room erature and pressure?	
	[1 mol	l of any gas occupies 24 dm ³ at room temperature and pressure.]	
	A 🛛	0.010	
	B	0.020	
	🖾 C	0.024	
	🖾 D	0.030	
		(Total for Question 8 = 1 m	ark)



9	Wh	at i	s the percen	tage by mass of nit	rogen in ammoniur	n nitrate, NH₄NO₃?	
	[Mc	olar	masses/g m	nol ⁻¹ : N = 14.0; H =	1.0; O = 16.0]		
	×	A	14.0%				
	\times	B	17.5%				
	\mathbf{X}	С	28.0%				
	×	D	35.0%				
						(Total for Questio	n 9 = 1 mark)
10	The	e fir	st five succe	ssive ionization ene	rgies for an elemen	t J, in kJ mol⁻¹, are	
			1st	2nd	3rd	4th	5th
			738	1450	7733	10543	13630
	The	e fo	rmula of the	compound of chlo	rine with element J	is	
			JCI				
	X	В	JCl ₂				
	×	С	JCl₃				
	X	D	J_2CI_3				
					("	Total for Question	10 = 1 mark)
11			of the follow nts of Period		order of increasing r	nelting temperatur	e of
	×	A	Na, Mg, Al,	Si			
	X	B	Na, Mg, Si, J	AI			
	X	C	Si, Na, Mg, J	AI			
	\times	D	Si, Al, Mg, N	la			
					("	Total for Question	11 = 1 mark)

P 4 4 8 7 9 A 0 6 2 4

12 Which one of the following elements undergoes the change in electronic configuration shown when it forms the stated ion?							
	Atom 1s ² 2s ² 2p ⁶ 3s ² 3p ³ Ion 1s ² 2s ² 2p ⁶ 3s ² 3p ⁶						
🖾 A	B to B ³⁺						
B	AI to AI ³⁺						
🖂 C	N to N ³⁻						
D	P to P ³⁻						
		(Total for Question 12 = 1 mark)					
13 Which	of the following compo	unds has the most polarized anion ?					
A	Na ₂ O						
B	MgO						
🖾 C	K ₂ O						
D	CaO						
		(Total for Question 13 = 1 mark)					
		7					







15 Calcul	late the entha	lpy change, in kJ mol⁻¹, for	the reaction	
		$H_2(g) + \frac{1}{2}O_2(g)$	\rightarrow H ₂ O(g)	
DATA:				
		Bond	Bond enthalpy / kJ mol ⁻¹	
		Н—Н	+436	
		0=0	+498	
		H—O	+464	
🖾 A	-243			
B	-6			
🛛 C	+6			
🛛 D	+221			
			(Total for Question	15 = 1 mark)
16 The cl	hemical prope	erties of an element are det	rermined by its	
	electronic st			
🛛 C	relative ator			
🖾 D	number of p	protons plus neutrons.		
			(Total for Question	16 = 1 mark)
			TOTAL FOR SECTION A	. = 20 MARKS
				9





BLANK PAGE





*(f) One of the uses of mass spectrometers is for the detection of banned substances, such as anabolic steroids, in a blood or urine sample taken from competitors in sports events.	
 Suggest two precautions that are necessary to ensure that the result of any analysis would be valid. 	
	(2)
(ii) These substances can give competitors an unfair advantage. Suggest why the	
use of these substances may be of concern to the user.	(1)
(g) Suggest one other use for mass spectrometers.	
	(1)
(Total for Question 17 = 16 ma	rkc)
	1K5)



18	Alkenes and cycloalkanes have the	he same ge	neral form	ula, but react very differently	
	with halogens.				
	(a) Give the general formula that	t applies to	both alke	nes and cycloalkanes.	(1)
	(b) Using structural formulae, co				
	alkene of your own choice, co bromine.	ontaining fe	ewer than	four carbon atoms, with liquid	
	Name the product.				(3)
		+	Br ₂	\rightarrow	
				Name:	



*(c) An example of an alkene with six carbon atoms is 2-methylpent-1-ene. It reacts with chlorine by means of an electrophilic addition reaction. The diagram below shows a student's attempt at drawing the mechanism for this reaction.

	H CH3 H CH3 H CH3 H CH3 H CH3 H CH3 H CH3 H CH3 H CH3 G C CH2-CH2-CH2-CH2-CH2-CH2-CH2-CH2-CH2-CH2-	CH3
Error	(i) Identify the three errors in this student's drawing of the mechanism.	(3)
Error 2	2	
Error 3	3	
	(ii) The structure of the carbocation intermediate is correctly drawn. Explain why the positive charge is on the carbon atom shown.	(1)



(d) There are five possible cycloalkanes, each containing five carbon atoms. Three of the isomers are given below. Complete the other two boxes, by adding the skeletal formulae of the other two structural isomers. (2) (e) Define the term **structural isomerism**. (1) (f) Another example of a cycloalkane is cyclobutane. This compound, like other cycloalkanes, can also react with chlorine. The overall reaction of cyclobutane with chlorine is as follows: $C_4H_8 + CI_2 \rightarrow C_4H_7CI + HCI$ (i) This reaction can occur at room temperature and pressure. What further condition is needed for this reaction to take place? (1) (ii) Using the appropriate arrows, complete the equation for the initiation step of the reaction mechanism for the reaction of chlorine with cyclobutane. (2) $CI \longrightarrow CI \longrightarrow$



 (iii) Using molecular formulae, write equations for the two propagation steps of this mechanism. First propagation step: 	(2)
Second propagation step:	
(iv) Name the type of bond fission which occurs in these propagation steps.	(1)
(v) There are also termination steps in this mechanism. Explain how these differ from the other steps in the mechanism and why these result in the reaction ending.	(2)
(g) If the reaction with cyclobutane is carried out with an excess of chlorine, how are the products of the reaction affected?	(1)
(Total for Question 18 = 20 ma	ırks)
	17

 19 Sodium and chlorine react together to produce sodium chloride. The bonding in the product is different from that in both of the reactants. Evidence for the type of bonding present can be obtained in a number of different ways. (a) Draw the electron density map for a chlorine molecule to show covalent bonding 	g. (1)
(b) Sodium chloride is ionically bonded. What is meant by the term ionic bond ?	(1)
 (c) Electrolysis is an experiment which you could carry out in a school or college laboratory on an aqueous solution of sodium chloride, to provide evidence for the presence of ionic bonding. Draw a labelled diagram of the apparatus that you would use for this experiment indicating how your results would show that the bonding was ionic. 	



(d) Chlorine gains an electron when it reacts with sodium to form sodium chloride. (i) Draw the dot and cross diagram of a chloride ion showing outer electrons o	
(ii) Give the formula of an ion from Period 3 that is isoelectronic with the chlori	de ion. (1)
(e) Sodium and sodium chloride can both be good conductors of electricity.	
Under what conditions do these substances conduct electricity?	
Compare the method of conductivity in each case.	(3)
(Total for Question 19 = 10 r	marks)

20 The reaction of calcium oxide with hydrochloric acid is an exothermic reaction.

 $CaO(s) + 2HCI(aq) \rightarrow CaCI_2(aq) + H_2O(I)$

In an experiment to investigate this reaction, the following procedure was carried out.

- 1. 50.0 cm³ of hydrochloric acid, concentration 2.0 mol dm⁻³ (an excess), was pipetted into a polystyrene cup and the initial temperature measured using a thermometer with 0.5 °C graduations.
- 2. 1.46 g of calcium oxide powder was weighed out and added to the acid. The mixture was stirred and the maximum temperature measured.

Maximum temperature / °C	35.0
Initial temperature / °C	19.5

(a) Calculate the enthalpy change, in joules, for the quantities in this experiment. Assume that the specific heat capacity of the solution is $4.18 \text{ J g}^{-1} \text{ C}^{-1}$.

Use the expression:

energy transferred in joules = $50.0 \times$ specific heat capacity \times temperature change

(1)

(b) Using your answer from (a), calculate the molar enthalpy change for the reaction between calcium oxide and hydrochloric acid. Include a sign and units in your answer.

(2)



(c)		e standard molar enthalpy change for the reaction between calcium oxide and drochloric acid is $-196.8 \text{ kJ mol}^{-1}$.	
	(i)	Suggest three reasons why the calculated value in part (b) is different from this value.	(3)
Reaso	n 1		(5)
Reaso	n 2		
Reaso	n 3		
	(ii)	Using the standard enthalpy change of –196.8 kJ mol ⁻¹ , calculate the minimum mass of calcium oxide that would be needed to raise the temperature of 250 cm ³ of hydrochloric acid (an excess) by 25.0 °C.	(3)



(d) The reaction of calcium carbonate with hydrochloric acid has the following standard molar enthalpy change.

$$\Delta H^{\ominus} = -18.8 \text{ kJ mol}^{-1}$$

This value can be used, with the enthalpy change for the reaction of calcium oxide with hydrochloric acid, to determine the enthalpy change for the thermal decomposition of calcium carbonate. This cannot be measured directly.

(i) Complete the Hess energy cycle below by adding the missing arrow and entities.

Use the cycle, and the standard enthalpy change for the reaction of calcium oxide and hydrochloric acid (–196.8 kJ mol⁻¹), to determine the standard enthalpy change for the decomposition of calcium carbonate.



Enthalpy change = kJ mol⁻¹

(4)





P 4 4 8 7 9 A 0 2 3 2 4

1				-				5			m	4	LC LC	9	7	0 (8)
4											י	r	2	>	-	(18)
0			Kev			1.0 hydrogen 1					(13)	(14)	(15)	(16)	(21)	4.0 He helium
0.6	Г	relat	relative atomic mass	mass							10.8	12.0	14.0	16.0	19.0	20.2
		atc	atomic symbol	pol							B	υ	z	0	Ŀ	Ne
lithium beryllium 3 4	E	atomic	name atomic (proton) number	number							boron 5	carbon 6	nitrogen 7	oxygen 8	fluorine 9	neon 10
23.0 24.3	T]							27.0	28.1	31.0	32.1	35.5	39.9
											AI	Si	٩.	S	บ	Ar
sodium magnesium 11 12	m (3)	(4)	(2)	(9)	(2)	(8)	(6)	(10)	(11)	(12)	aluminium 13	silicon 14	phosphorus 15	sulfur 16	chlorine 17	argon 18
39.1 40.1	45.0	47.9	50.9	52.0	54.9	55.8	58.9	58.7	63.5	65.4	69.7	72.6	74.9	79.0	79.9	83.8
Ca		ï		ں د		Fe	ပိ	Ż	Cu	Zn	Ga	Ge	As	Se	Br	ĸ
potassium calcium 19 20	n scandium 21	titanium 77	vanadium 73	chromium 74	ma	iron 26	cobalt 77	nickel 28	29	zinc 30	gallium 31	germanium 32	arsenic 33	selenium 34	bromine 35	krypton 36
85.5 87.6	—	91.2	92.9	95.9	[98]	101.1	102.9	106.4	107.9	112.4	114.8	118.7	121.8	127.6	126.9	131.3
Rb Sr	≻	Zr	qN	Wo	Ч	Ru	Rh	РЧ	Ag	Cd	Ľ	Sn	Sb	Te	_	Xe
rubidium strontium 37 38	m yttrium 39	zirconium 40	niobium 41	molybdenum 42	technetium 43	ruthenium 44	rhodium 45	palladium 46	silver 47	cadmium 48	indium 49	tin 50	antimony 51	tellurium 52	iodine 53	xenon 54
132.9 137.3	138.9	178.5	180.9	183.8	186.2	190.2	192.2	195.1	197.0	200.6	204.4	207.2	209.0	[209]	[210]	[222]
		Hf	Ta	≯	Re	0s	<u> </u>	Pt	Αu	Hg	F	Pb	Bi	Ъо	At	Rn
caesium barium 55 56	r lanthanum 57	hafnium 72	tantalum 73	tungsten 74	rhenium 75	osmium 76	iridium 77	platinum 78	gold 79	mercury 80	thallium 81	lead 82	bismuth 83	polonium 84	astatine 85	radon 86
[223] [226]		[261]		[266]	_	[277]	[268]		[272]							
Fr Radium	AC*	Rf ntherfordium	Db	Sg seahoroium	Bh bohrium		Mt	DS damstadtium	Rg		nents with	atomic nui but not f	tomic numbers 112-116 hav	116 have b Iticated	Elements with atomic numbers 112-116 have been reported but not fully authenticated	ed
_		104	105	106		108	109	110	111					5		
		140	141	144	[147]	150	152	157	159	163	165	167	169	173	175	
* Lanthanide series	ries	Ce	Pr	PN	Pm	Sm	Eu			Dy	Ю	Er	Tm	Υb	Lu	
* Actinide series	Ş	cerium 58	praseodymium 59	praseodymium neodymium promethium 59 60 61	promethium 61	samarium 62	europium 63	gadolinium 64	terbium 65	dysprosium 66	holmium 67	erbium 68	thulium 69	ytterbium 70	lutetium 71	
		232	[231]		[237]	[242]	[243]	[247]	[245]	[251]	[254]	[253]	[256]	[254]	[257]	
					ЧN	Pu	Am		BĶ		Es	Fm	PW	No	Ļ	
		thorium 90	protactinium 01	uraniur 00	n neptunium plutonium ar 03 04	plutonium 0.4	americium 05	aurium 0,6	berkelium 9.7		californium einsteinium	fermium 100	mendelevium	nobelium	lawrencium	
				77	57	74	<i>ر د</i>		71		77	202	101	1 701	CU1	

The Periodic Table of Elements

P 4 4 8 7 9 A 0 2 4 2 4