


2.

Figure 1



The curve with equation $y = \frac{1}{3(1+2x)}$, $x > -\frac{1}{2}$, is shown in Figure 1.

The region bounded by the lines $x = -\frac{1}{4}$, $x = \frac{1}{2}$, the x -axis and the curve is shown shaded in Figure 1.

This region is rotated through 360 degrees about the x -axis.

(a) Use calculus to find the exact value of the volume of the solid generated.

(5)

Figure 2




Figure 2 shows a paperweight with axis of symmetry AB where $AB = 3$ cm. A is a point on the top surface of the paperweight, and B is a point on the base of the paperweight. The paperweight is geometrically similar to the solid in part (a).

(b) Find the volume of this paperweight.

(2)



6. (a) Given that $y = 2^x$, and using the result $2^x = e^{x \ln 2}$, or otherwise, show that $\frac{dy}{dx} = 2^x \ln 2$. (2)

(b) Find the gradient of the curve with equation $y = 2^{(x^2)}$ at the point with coordinates (2,16). (4)



8.

$$I = \int_0^5 e^{\sqrt{3x+1}} dx.$$

- (a) Given that $y = e^{\sqrt{3x+1}}$, complete the table with the values of y corresponding to $x = 2, 3$ and 4 .

x	0	1	2	3	4	5
y	e^1	e^2				e^4

(2)

- (b) Use the trapezium rule, with all the values of y in the completed table, to obtain an estimate for the original integral I , giving your answer to 4 significant figures.

(3)

- (c) Use the substitution $t = \sqrt{3x+1}$ to show that I may be expressed as $\int_a^b kte^t dt$, giving the values of a, b and k .

(5)

- (d) Use integration by parts to evaluate this integral, and hence find the value of I correct to 4 significant figures, showing all the steps in your working.

(5)



