

Please write clearly in	n block capitals.
Centre number	Candidate number
Surname	
Forename(s)	
Candidate signature	I declare this is my own work.

A-level PHYSICS

Paper 3
Section B Astrophysics

Friday 5 June 2020

Afternoon

Materials

For this paper you must have:

- a pencil and a ruler
- a scientific calculator
- a Data and Formulae Booklet.

Time allowed: The total time for both sections of this paper is 2 hours. You are advised to spend approximately 50 minutes on this section.

Instructions

- Use black ink or black ball-point pen.
- Fill in the boxes at the top of this page.
- Answer all questions.
- You must answer the questions in the spaces provided. Do not write outside the box around each page or on blank pages.
- If you need extra space for your answer(s), use the lined pages at the end of this book. Write the question number against your answer(s).
- Do all rough work in this book. Cross through any work you do not want to be marked.
- Show all your working.

Information

- The marks for questions are shown in brackets.
- The maximum mark for this paper is 35.
- You are expected to use a scientific calculator where appropriate.
- A Data and Formulae Booklet is provided as a loose insert.

For Examiner's Use			
Question	Mark		
1			
2			
3			
4			
TOTAL			



	2	
	Section B	
	Answer all questions in this section.	
0 1.1	Draw a ray diagram for a Cassegrain telescope. Your diagram should show the paths of two rays up to the eyepiece lens. The rays should initially be parallel to the principal axis.	[2 marks]
		_ principal axis
0 1.2	A spacecraft passes Pluto at a distance of $12500\mathrm{km}$. The telescope on boan aperture of diameter $0.21\mathrm{m}$ and operates at a wavelength of $450\mathrm{nm}$.	
	Discuss whether this telescope is suitable for studying a crater with a diamental approximately $1\ km$ on Pluto.	eter of [3 marks]



0 1.3	The Hubble telescope has an aperture of diameter 2.4 m.	(
	Compare the collecting power of the Hubble telescope with the telescope on the spacecraft in Question 01.2 . [2 marks]	
0 1.4	An astrophysicist had to decide whether to use a reflecting telescope or a refracting telescope on the spacecraft in Question 01.2 .	
	Discuss which type of telescope to use. [3 marks]	
		<u> </u>

Turn over ▶



0 2

Table 1 summarises some information about four stars in the constellation Cassiopeia.

Table 1

Name	Colour	Apparent magnitude	Distance / ly
Caph	white	2.3	55
Ruchbah	blue/white	2.7	99
Schedar	orange	2.2	228
Tsih	blue	2.2	610

0 2.1	Which star has the highest surface temperature? Tick (\checkmark) one box.	[1 mark]
	Caph	
	Ruchbah	
	Schedar	
	Tsih	




0 2 . 2

Figure 1 shows the intensity received at Earth from two of the stars, plotted against wavelength.

The effect of absorption by the Earth's atmosphere is not shown.

Figure 1



Discuss what information can be found from **Figure 1** about the temperature and colour of these stars.

Support your answer with suitable calculations.

[4 marks]

0	uestion 2 cont	inues on the r	next nage	





0 2.3	State which star in Table 1 is dimmest on the absolute magnitude scale.	[1 mark]	Oi
0 2.4	Calculate the absolute magnitude of Schedar.	[3 marks]	
	absolute magnitude =		
0 2.5	Tsih has a mass over 15 times the mass of the Sun. Tsih may eventually collapse to form a black hole.		
	Calculate the radius of the event horizon for a black hole with a mass 15 tir the Sun.	mes that of [2 marks]	
	radius =	m	_



7 0 3 Type 1a supernovae can be used as standard candles. 0 3 State what is meant by a standard candle. [1 mark] 0 3 . 2 Sketch on **Figure 2** the light curve for a type 1a supernova. Annotate your graph with suitable scales and a unit for time. [3 marks] Figure 2 absolute magnitude time / Question 3 continues on the next page

Turn over ▶

Do not write outside the



0 3.3	Measurements of type 1a supernovae are used to find a value for the Hubble constant.			
	The distance from Earth is known for many type 1a supernovae.			
	Describe how these values of distance are used, with other data, to find the Hubb constant.			
	Your answer should include:			
	 the other data needed and how these data are used the graph plotted, including appropriate units for the axes how the Hubble constant is obtained and any limitations on the result. [6 marks]			



Do not write outside the box
10

Turn over ▶



Do not write outside the

0 4

 $\textbf{Table 2} \ \text{gives data about the supergiant star Melnick } 34 \ \text{and the Sun}.$

Table 2


Name	Radius / m	Surface temperature / K
Melnick 34	1.4×10^{10}	53 000
Sun	7.0×10^{8}	5 700

	Melnick 34	1.4×10^{10}	53 000	
	Sun	7.0×10^{8}	5 700	
0 4.1	Calculate power power	output of Melnick 34 er output of the Sun		[2 marks]
0 4.2	Discuss why the dangerous for life		answer =ar in the local part of our gala	xy could be [2 marks]

4

END OF QUESTIONS







Question number	Additional page, if required. Write the question numbers in the left-hand margin.



Question number	Additional page, if required. Write the question numbers in the left-hand margin.



Question number	Additional page, if required. Write the question numbers in the left-hand margin.



Question number	Additional page, if required. Write the question numbers in the left-hand margin.



16 Do not write outside the There are no questions printed on this page DO NOT WRITE ON THIS PAGE ANSWER IN THE SPACES PROVIDED

Copyright information

For confidentiality purposes, all acknowledgements of third-party copyright material are published in a separate booklet. This booklet is published after each live examination series and is available for free download from www.aqa.org.uk.

Permission to reproduce all copyright material has been applied for. In some cases, efforts to contact copyright-holders may have been unsuccessful and AQA will be happy to rectify any omissions of acknowledgements. If you have any queries please contact the Copyright Team.

Copyright © 2020 AQA and its licensors. All rights reserved



